

Gushel–Mukai varieties: moduli

Olivier Debarre

*Université Paris-Diderot, PSL Research University
CNRS, École normale supérieure
Département de Mathématiques et Applications
45 rue d'Ulm, 75230 Paris Cedex 05, France
olivier.debarre@ens.fr*

Alexander Kuznetsov*

*Algebraic Geometry Section, Steklov Mathematical Institute
of Russian Academy of Sciences, 8 Gubkin str.
Moscow 119991, Russia
akuznet@mi-ras.ru*

Received 14 June 2019

Accepted 23 October 2019

Published 7 February 2020

We describe the moduli stack of Gushel–Mukai varieties as a global quotient stack and its coarse moduli space as the corresponding GIT quotient. The construction is based on a comprehensive study of the relation between this stack and the stack of so-called Lagrangian data defined in our previous works; roughly speaking, we show that the former is a generalized root stack of the latter. As an application, we define the period map for Gushel–Mukai varieties and construct some complete nonisotrivial families of smooth Gushel–Mukai varieties. In an appendix, we describe a generalization of the root stack construction used in our approach to the moduli stack.

Keywords: Gushel–Mukai varieties; moduli stacks; moduli spaces; period map; root stacks.

Mathematics Subject Classification 2010: 14D22, 14D23, 14J10, 14J45, 14J30, 14J35, 14J40, 14D07

1. Introduction

This paper is the third in the series started in [6, 7] and devoted to the investigation of Gushel–Mukai (GM) varieties defined over a field \mathbf{k} of characteristic zero. These varieties are positive-dimensional, dimensionally transverse intersections

$$X = \text{CGr}(2, V_5) \cap \mathbf{P}(W) \cap Q,$$

*Corresponding author.

where $\text{CGr}(2, V_5) \subset \mathbf{P}(\mathbf{k} \oplus \wedge^2 V_5)$ is the cone over the Plücker embedding of the Grassmannian of two-dimensional vector subspaces in a \mathbf{k} -vector space V_5 of dimension 5, $\mathbf{P}(W) \subset \mathbf{P}(\mathbf{k} \oplus \wedge^2 V_5)$ is a projective space of dimension $n + 4$, and Q is a quadric hypersurface in $\mathbf{P}(W)$. We have

$$\dim(X) = n \in \{1, \dots, 6\}.$$

Various geometric characterizations of GM varieties can be found in [6, Sec. 2.3]; for instance, [6, Theorem 2.16] shows that smooth GM varieties of dimension $n \geq 3$ are exactly the Fano varieties of Picard rank 1, coindex 3, and degree 10.

In [6], we described the set of isomorphism classes of all GM varieties. In particular, we associated with each GM variety what we called a **GM data set**. Roughly speaking, it is a collection $(W, V_6, V_5, \mu, \mathbf{q})$, where W , V_6 , and V_5 are \mathbf{k} -vector spaces of respective dimensions $n + 5$, 6, and 5, with $V_5 \subset V_6$, and

$$\mu : W \rightarrow \wedge^2 V_5 \quad \text{and} \quad \mathbf{q} : V_6 \rightarrow \text{Sym}^2 W^\vee$$

are \mathbf{k} -linear maps. The map μ is the composition of the embedding $W \hookrightarrow \mathbf{k} \oplus \wedge^2 V_5$ coming from the definition of X with the projection onto the second summand, whereas the map \mathbf{q} is obtained by identifying V_6 with the space of quadratic equations of X in $\mathbf{P}(W)$. Under this identification, the hyperplane $V_5 \subset V_6$ corresponds to the space of Plücker quadrics defining $\text{CGr}(2, V_5)$ in $\mathbf{P}(\mathbf{k} \oplus \wedge^2 V_5)$.

There are two types of smooth GM varieties: ordinary and special, distinguished by the injectivity or the noninjectivity of the map μ . When the field \mathbf{k} is quadratically closed, there is a natural bijection between the set of isomorphism classes of special GM varieties of dimension n and the set of ordinary GM varieties of dimension $n - 1$ [6, Lemma 2.33]. On the other hand, special GM varieties can be obtained from ordinary GM varieties of the same dimension by specialization (except in the case $n = 6$, when there are no ordinary GM varieties). However, they behave in a slightly different way and provide various complications to the theory.

The first main result of [6], Theorem 2.9, provides a bijection between the set of isomorphism classes of GM varieties of dimension n and an appropriate subset of the set of isomorphism classes of GM data sets. After introducing in Secs. 3.1 and 3.2 the stacks of GM varieties and of GM data, we present in Sec. 3.3 a version of this bijection that works for families and promotes the bijection of sets of isomorphism classes to an isomorphism of moduli stacks (see Theorem 3.7).

The second main result of [6], Theorem 3.6, relates GM data sets of ordinary GM varieties to so-called **Lagrangian data sets**. These consist of triples (V_6, V_5, A) , where V_6 is a vector space of dimension 6, $V_5 \subset V_6$ is a hyperplane, and $A \subset \wedge^3 V_6$ is a subspace which is Lagrangian for the $\det(V_6)$ -valued symplectic form on $\wedge^3 V_6$ given by wedge product. Theorem 3.6 of [6] establishes a bijection between ordinary GM data sets of dimension n and Lagrangian data sets with $\dim(A \cap \wedge^3 V_5) = 5 - n$, as well as, (if \mathbf{k} is quadratically closed) using the bijection between ordinary and special GM data sets, a bijection between special GM data sets of dimension n and Lagrangian data sets such that $\dim(A \cap \wedge^3 V_5) = 6 - n$.

A nice feature of this bijection, proved in [6, Theorem 3.16], is that the smoothness of the GM variety associated with a Lagrangian data set (V_6, V_5, A) only depends on A : when $n \geq 3$, the corresponding GM variety is smooth if and only if A has no decomposable vectors, that is,

$$\mathbf{P}(A) \cap \mathrm{Gr}(3, V_6) = \emptyset,$$

the intersection being taken inside $\mathbf{P}(\wedge^3 V_6)$.

The main goal of this paper is to combine all these constructions into a single construction of the moduli stack of smooth GM varieties. In other words, we find analogs of the above constructions that work for “mixed” families (with both ordinary and special varieties as members). The main difficulty is that the GM/Lagrangian data sets bijection of [6] does not work with these “mixed” families. Nevertheless, we define in Sec. 3.5 the stack of Lagrangian data that classifies all Lagrangian data sets (V_6, V_5, A) such that

$$\dim(A \cap \wedge^3 V_5) \in \{5 - n, 6 - n\}$$

and such that A has no decomposable vectors. We observe in Sec. 4.1 (see Proposition 4.1) that the natural family version of the construction from [6, Theorem 3.6] that associates with a family of GM data $(S, \mathscr{W}, \mathscr{V}_6, \mathscr{V}_5, \mu, \mathbf{q})$ over a scheme S a family of Lagrangian data $(S, \mathscr{V}_6, \mathscr{V}_5, \mathscr{A})$ is still well defined and gives a morphism of stacks. However, this morphism cannot be an isomorphism for the following simple reason.

For any family of GM data $(S, \mathscr{W}, \mathscr{V}_6, \mathscr{V}_5, \mu, \mathbf{q})$ over a scheme S , we define in Sec. 3.4 a closed subset $S_{\mathrm{GM}, \mathrm{spe}} \subset S$ corresponding to points of S that parameterize GM data sets of special varieties and endow it with a natural scheme structure (Definition 3.11). Similarly, given a family of Lagrangian data $(S, \mathscr{V}_6, \mathscr{V}_5, \mathscr{A})$, we consider the closed subset $S_{\mathrm{Lag}, \mathrm{spe}} \subset S$ corresponding to points of S such that $\dim(A \cap \wedge^3 V_5) = 6 - n$ and endow it with a natural scheme structure (Definition 3.19). An important consequence of Proposition 4.1 is that although the special loci of an S -family of GM data and of the associated S -family of Lagrangian data are the same set-theoretically, they have different scheme structures: the ideal of $S_{\mathrm{Lag}, \mathrm{spe}}$ is the square of the ideal of $S_{\mathrm{GM}, \mathrm{spe}}$. Consequently, if we start with a family of Lagrangian data such that the ideal of its special locus is not a square, there is no corresponding family of GM data!

However, we prove in Sec. 4.2 that the inverse construction can be made when the ideal of the subscheme $S_{\mathrm{Lag}, \mathrm{spe}} \subset S$ for an S -family of Lagrangian data is a square and $S_{\mathrm{Lag}, \mathrm{spe}}$ is a Cartier divisor in S (this second condition seems to be of technical nature but we do not know how to make the inverse construction without it). This is the central construction of the paper. It is based on two vector bundle constructions which we develop in Sec. 2 and which are interesting by themselves.

The first is the *canonical factorization construction* of Proposition 2.8: given a morphism of vector bundles $\varphi : \mathscr{E} \rightarrow \mathscr{F}$ of generic rank r over a scheme S , such that the rank of φ is everywhere at least $r - 1$ and the degeneration scheme of φ is

a Cartier divisor $D \subset S$, we construct a canonical factorization

$$\mathcal{E} \rightarrow \mathcal{E}_1 \xrightarrow{\varphi_1} \mathcal{F}_1 \hookrightarrow \mathcal{F},$$

where \mathcal{E}_1 and \mathcal{F}_1 are vector bundles of rank r , the first map is an epimorphism, the last map is a fiberwise monomorphism, and the map φ_1 is an embedding of coherent sheaves whose cokernel is a line bundle on D .

Given a family of Lagrangian data $(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$ such that $S_{\text{Lag, spe}}$ is a Cartier divisor in S , we apply in Sec. 4.2 this construction to the composition

$$\varphi : \mathcal{A} \hookrightarrow \bigwedge^3 \mathcal{V}_6 \xrightarrow{\lambda_3} \bigwedge^2 \mathcal{V}_5 \otimes (\mathcal{V}_6/\mathcal{V}_5),$$

(where the second map is induced by the natural projection $\lambda : \mathcal{V}_6 \rightarrow \mathcal{V}_6/\mathcal{V}_5$) and obtain a factorization

$$\mathcal{A} \rightarrow \mathcal{W}' \xrightarrow{\varphi_1} \mathcal{W}'' \hookrightarrow \bigwedge^2 \mathcal{V}_5 \otimes (\mathcal{V}_6/\mathcal{V}_5). \tag{1}$$

The cokernel of the morphism φ_1 is supported on the Cartier divisor $D = S_{\text{Lag, spe}}$. Assuming that this divisor can be written as

$$D = 2E,$$

where E is also a Cartier divisor, we find a unique vector bundle \mathcal{W} such that the morphism φ_1 factors as

$$\mathcal{W}' \xrightarrow{\varphi'} \mathcal{W} \xrightarrow{\varphi''} \mathcal{W}'' ,$$

where both φ' and φ'' are embeddings of sheaves whose cokernels are line bundles on E . We prove in Proposition 4.6 that $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$, where μ is the composition

$$\mathcal{W} \xrightarrow{\varphi''} \mathcal{W}'' \hookrightarrow \bigwedge^2 \mathcal{V}_5 \otimes (\mathcal{V}_6/\mathcal{V}_5)$$

and the map \mathbf{q} will be defined below, is a family of GM data corresponding to a smooth family of GM varieties, whose associated family of Lagrangian data is equivalent to $(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$.

The construction of the map $\mathbf{q} : \mathcal{V}_6 \rightarrow \text{Sym}^2 \mathcal{W}^\vee$ is carried out in three steps (there is actually an extra line bundle twist on the target of \mathbf{q} , but we will ignore it here for simplicity). First, we define a map

$$\mathbf{q}_{\mathcal{A}} : \mathcal{V}_6 \rightarrow \text{Sym}^2 \mathcal{A}^\vee$$

by an explicit formula (35), which is just a family version of a formula used in the proof of [6, Theorem 3.6]. After that we observe that the kernel of the epimorphism $\mathcal{A} \rightarrow \mathcal{W}'$ is contained in the kernel of $\mathbf{q}_{\mathcal{A}}$, hence there is a morphism

$$\mathbf{q}' : \mathcal{V}_6 \rightarrow \text{Sym}^2 \mathcal{W}'^\vee \tag{2}$$

induced by $\mathbf{q}_{\mathcal{A}}$. The last step is the construction of a map

$$\mathbf{q} : \mathcal{V}_6 \rightarrow \text{Sym}^2 \mathcal{W}^\vee \tag{3}$$

such that $(\text{Sym}^2 \varphi'^T) \circ \mathbf{q} = \mathbf{q}'$; it uses the second vector bundle construction from Sec. 2.

This second construction is explained in Sec. 2.5; we call it the *Hecke transform* of a family of quadratic forms. It starts from a morphism $\mathcal{V} \rightarrow \text{Sym}^2 \mathcal{E}^\vee$ of vector bundles over a scheme S (viewed as a family of quadrics in $\mathbf{P}_S(\mathcal{E})$ parameterized by $\mathbf{P}_S(\mathcal{V})$), a double Cartier divisor $D = 2E$ on S , and a line subbundle $\mathcal{K} \subset \mathcal{E}|_D$ contained in the kernel of all quadratic forms (restricted to D). We define a new vector bundle $\tilde{\mathcal{E}}$ on S by the exact sequence

$$0 \rightarrow \tilde{\mathcal{E}}^\vee \xrightarrow{\epsilon} \mathcal{E}^\vee \rightarrow \mathcal{K}^\vee|_E \rightarrow 0$$

and check in Proposition 2.12 that there is a unique family of quadratic forms $\mathcal{V} \rightarrow \text{Sym}^2 \tilde{\mathcal{E}}^\vee$ such that the original family of quadratic forms is the composition of this family with $\text{Sym}^2 \epsilon$.

We apply this construction to the family of quadratic forms \mathbf{q}' from (2), with $\mathcal{V} = \mathcal{V}_6$ and $\mathcal{E} = \mathcal{W}'$, taking $D = S_{\text{Lag, spe}}$ and $\mathcal{K} = \text{Ker}(\varphi_1|_D : \mathcal{W}'|_D \rightarrow \mathcal{W}''|_D)$. The corresponding Hecke transform $\widetilde{\mathcal{W}'}$ is just \mathcal{W} , so Proposition 2.12 provides the required family \mathbf{q} of quadratic forms on \mathcal{W} as in (3). We also prove in Proposition 2.12 that there is a canonical direct sum decomposition

$$\mathcal{W}|_E \simeq (\mathcal{W}'|_E)/(\mathcal{K}|_E) \oplus (\mathcal{K}|_E)(E)$$

which is orthogonal for all quadrics in the family $\mathbf{q}|_E$ and which recovers the canonical direct sum decomposition of [6, Proposition 2.30] for special GM data sets. This observation is essential for proving that the constructed family of GM varieties is smooth.

In Sec. 5, we use the constructions of Sec. 4 to provide a description of the moduli stack of smooth GM varieties as a global quotient stack. We first fix a vector space V_6 of dimension 6 and consider the scheme

$$\overline{\mathbf{S}}_n = \left\{ (A, V_5) \in \text{LGr}(\wedge^3 V_6) \times \mathbf{P}(V_6^\vee) \mid \begin{array}{l} \dim(A \cap \wedge^3 V_5) \in \{5 - n, 6 - n\} \text{ and} \\ A \text{ has no decomposable vectors} \end{array} \right\}.$$

The first condition $\dim(A \cap \wedge^3 V_5) \geq 5 - n$ is closed, while the other two conditions $\dim(A \cap \wedge^3 V_5) \leq 6 - n$ and “ A has no decomposable vectors” are open, so $\overline{\mathbf{S}}_n \subset \text{LGr}(\wedge^3 V_6) \times \mathbf{P}(V_6^\vee)$ is a locally closed subscheme. When $n \in \{3, 4, 5\}$, this scheme contains a closed subscheme \mathbf{S}_{n-1} and its open complement \mathbf{S}_n , defined by the conditions that $\dim(A \cap \wedge^3 V_5)$ equals $6 - n$ and $5 - n$, respectively, while $\overline{\mathbf{S}}_6 = \mathbf{S}_5$. The fibers of the projection $\mathbf{S}_n \rightarrow \text{LGr}(\wedge^3 V_6)$ over a point $[A]$ are just the strata of the Eisenbud–Popescu–Walter stratification of $\mathbf{P}(V_6^\vee)$ associated with A (see [15, Sec. 2] or Sec. 5.1) and the fibers of $\overline{\mathbf{S}}_n$ are unions of these strata. In particular, the schemes $\overline{\mathbf{S}}_6$ and $\overline{\mathbf{S}}_5$ are smooth, and for $n \in \{3, 4\}$, one has $\mathbf{S}_{n-1} = \text{Sing}(\overline{\mathbf{S}}_n)$. Both strata \mathbf{S}_{n-1} and \mathbf{S}_n are *Lagrangian intersection loci*.

The construction of the moduli stack of smooth GM varieties of dimension n goes as follows. Assume $n \in \{3, 4, 5\}$ (the case $n = 6$ is slightly different and we skip it in this introduction). It was proved in [8] that, if a certain divisibility condition holds in the group $\text{Pic}(\overline{\mathbf{S}}_n)$ (in fact it does not, but we will go back to this point

later), there is a double covering

$$\tilde{S}_n \rightarrow \overline{S}_n$$

branched over S_{n-1} such that \tilde{S}_n is smooth. Note that $\text{codim}_{\overline{S}_n}(S_{n-1}) = 6 - n$, so for $n \leq 4$, this is not a classical double covering branched over a hypersurface. We consider the quotient stack

$$\widehat{S}_n := \tilde{S}_n / \mu_2$$

with respect to the involution of the double covering (this is the *canonical stack* of \overline{S}_n in the terminology of [24]). This is a smooth Deligne–Mumford stack and the natural $\text{PGL}(V_6)$ -action on the scheme \overline{S}_n lifts to a $\text{PGL}(V_6)$ -action on \widehat{S}_n . Our main theorem, Theorem 5.11, states that there is an isomorphism

$$\mathfrak{M}_n^{\text{GM}} \simeq \widehat{S}_n / \text{PGL}(V_6)$$

between the moduli stack $\mathfrak{M}_n^{\text{GM}}$ of smooth GM varieties of dimension $n \in \{3, 4, 5\}$ and the quotient stack $\widehat{S}_n / \text{PGL}(V_6)$.

The main step in the proof of this theorem is the construction of a family of smooth GM varieties over the stack \widehat{S}_n or, more precisely, over a certain scheme $\widehat{\widehat{S}}_n$ that provides a covering of \widehat{S}_n in the smooth topology (the morphism $\widehat{\widehat{S}}_n \rightarrow \widehat{S}_n$ is actually a \mathbb{G}_m -torsor). There is also a double covering map from $\widehat{\widehat{S}}_n$ to a certain \mathbb{G}_m -torsor over \overline{S}_n which is branched over the preimage of S_{n-1} . Consequently, pulling back from \overline{S}_n , we construct on $\widehat{\widehat{S}}_n$ a family of Lagrangian data $(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$ with trivial $\mathcal{V}_6 = V_6 \otimes \mathcal{O}$ and (pullbacks of) tautological bundles \mathcal{V}_5 and \mathcal{A} . The Lagrangian special locus of this family is the scheme-theoretic preimage of S_{n-1} , that is, the preimage of the branch locus of the double covering, hence its ideal is the square of the ideal of a certain smooth subscheme in $\widehat{\widehat{S}}_n$. Considering the blow up $\beta : \mathbf{S} \rightarrow \widehat{\widehat{S}}_n$ of this subscheme, we arrive at the situation of Sec. 4.2.

Applying Proposition 4.6, we obtain a family $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ of GM data. We check that this family is the pullback by β of a family of GM data on the scheme $\widehat{\widehat{S}}_n$. Moreover, this family is equivariant with respect to the natural action of the algebraic group

$$G_n = \text{GL}(V_6) / \mu_{3(5-n)}$$

and thus descends to a family of GM data over

$$\widehat{\widehat{S}}_n / G_n \simeq \widehat{S}_n / \text{PGL}(V_6). \tag{4}$$

This construction provides a morphism from the quotient stack $\widehat{S}_n / \text{PGL}(V_6)$ to the moduli stack of GM data. For the construction in the opposite direction, we use the much simpler procedure of Proposition 4.1 and a universal property of the stack $\widehat{\widehat{S}}_n$ proved in Proposition A.6. Combining these two constructions, we obtain an isomorphism between the moduli stack of smooth GM varieties and the global quotient stack (4).

To deal with the fact that the double covering $\widetilde{\mathcal{S}}_n \rightarrow \overline{\mathcal{S}}_n$ does not exist (since the required divisibility condition does not hold in $\text{Pic}(\overline{\mathcal{S}}_n)$), we note that the divisibility condition holds locally over $\overline{\mathcal{S}}_n$, so the double covering exists locally. One can obtain the stack $\widehat{\mathcal{S}}_n$ by gluing the quotient stacks of the local coverings, as in the standard construction of the root stack. Alternatively, one can construct the stack $\widehat{\mathcal{S}}_n$ directly (see Appendix A). After that, the construction goes as explained above.

To conclude this introduction, we mention that the global quotient stack description of Theorem 5.11 gives, via GIT, a construction of the coarse moduli space for smooth GM varieties (Theorem 5.15). This provides a foundation for the period map of GM varieties that was discussed in [7] (see Proposition 6.1). We also use our results to construct in Sec. 6.2 several examples of complete nonisotrivial families of smooth GM varieties.

2. Preliminaries on Vector Bundles

All schemes are over a fixed field \mathbf{k} .

We first discuss some aspects of the theory of vector bundles on possibly nonreduced schemes. Most of the material in Secs. 2.1 and 2.2 is well known but we collect it for the reader's convenience. The results of Secs. 2.3, 2.4, and 2.5 seem to be new and are essential for our treatment of the stack of GM varieties.

2.1. Epimorphisms and fiberwise monomorphisms

Let S be a scheme. By a *point of S* , we mean a \mathbf{K} -point $s : \text{Spec}(\mathbf{K}) \rightarrow S$ for some field \mathbf{K} . A *geometric point of S* is a \mathbf{K} -point with \mathbf{K} algebraically closed.

A vector bundle \mathcal{E} on S is a locally free sheaf of \mathcal{O}_S -modules of constant finite rank. Given a \mathbf{K} -point s of S , we let \mathcal{E}_s be the \mathbf{K} -vector space $\mathcal{E} \otimes_{\mathcal{O}_s} \mathbf{K}$, the *fiber* of \mathcal{E} at s .

Lemma 2.1. *A morphism $\varphi : \mathcal{E} \rightarrow \mathcal{F}$ between vector bundles on the scheme S is surjective if and only if, for every geometric point s of S , the induced linear map $\varphi_s : \mathcal{E}_s \rightarrow \mathcal{F}_s$ between fibers is surjective.*

Proof. Let \mathcal{C} be the cokernel of φ . Since the tensor product functor is right exact, we have, for each point s of S , an exact sequence

$$\mathcal{E}_s \xrightarrow{\varphi_s} \mathcal{F}_s \rightarrow \mathcal{C}_s \rightarrow 0.$$

By Nakayama's lemma, $\mathcal{C} = 0$ if and only if $\mathcal{C}_s = 0$ for every geometric point s of S . \square

We say that φ is a *fiberwise monomorphism* if, for every geometric point s of S , the morphism $\varphi_s : \mathcal{E}_s \rightarrow \mathcal{F}_s$ is a monomorphism.

Lemma 2.2. *A morphism $\varphi : \mathcal{E} \rightarrow \mathcal{F}$ between vector bundles on a scheme S is a fiberwise monomorphism if and only if the dual map $\varphi^\vee : \mathcal{F}^\vee \rightarrow \mathcal{E}^\vee$ is surjective.*

Proof. Since $(\mathcal{E}^\vee)_s = (\mathcal{E}_s)^\vee$, $(\mathcal{F}^\vee)_s = (\mathcal{F}_s)^\vee$, and $(\varphi^\vee)_s = (\varphi_s)^\vee$, the result follows from Lemma 2.1. \square

Epimorphisms and fiberwise monomorphisms enjoy the following nice properties.

Lemma 2.3. *Let $\varphi : \mathcal{E} \rightarrow \mathcal{F}$ be a morphism between vector bundles on a scheme S .*

If φ is surjective, $\text{Ker}(\varphi)$ is a vector bundle and the natural map $\text{Ker}(\varphi) \rightarrow \mathcal{E}$ is a fiberwise monomorphism.

If φ is a fiberwise monomorphism, $\text{Coker}(\varphi)$ is a vector bundle and the natural map $\mathcal{F} \rightarrow \text{Coker}(\varphi)$ is surjective.

Proof. The kernel is locally free since this is a local property and the kernel of an epimorphism of projective modules over a ring is projective. Furthermore, since \mathcal{F} is locally free, we have $\text{Tor}_1(\mathcal{F}, \mathbf{K}) = 0$ for any \mathbf{K} -point s of S , hence the sequence

$$0 \rightarrow \text{Ker}(\varphi)_s \rightarrow \mathcal{E}_s \rightarrow \mathcal{F}_s \rightarrow 0$$

is exact. By definition, the map $\text{Ker}(\varphi) \rightarrow \mathcal{E}$ is therefore a fiberwise monomorphism. The second part of the lemma follows by duality. \square

An effective Cartier divisor on a scheme is a subscheme locally defined by a regular function which is *not a zero divisor*.

Lemma 2.4. *Let S be a scheme, let $i : D \hookrightarrow S$ be an effective Cartier divisor, let \mathcal{E} be a vector bundle on S , and let \mathcal{F} be a vector bundle on D . If $\varphi : \mathcal{E} \rightarrow i_*\mathcal{F}$ is surjective, $\text{Ker}(\varphi)$ is a vector bundle on S .*

Proof. We may assume that D is nonempty. Since D is a Cartier divisor, the projective dimension as an \mathcal{O}_S -module of \mathcal{O}_D , hence also of $i_*\mathcal{F}$, is 1. Therefore, the projective dimension of $\text{Ker}(\varphi)$ is 0, so $\text{Ker}(\varphi)$ is locally free. \square

2.2. Degeneration schemes

Let $\varphi : \mathcal{E} \rightarrow \mathcal{F}$ be a morphism between vector bundles on a scheme S . For every nonnegative integer k , it induces a morphism

$$\wedge^k \varphi : \wedge^k \mathcal{E} \rightarrow \wedge^k \mathcal{F}$$

locally given by the $k \times k$ -minors of a matrix of regular functions defining φ . The rank of φ is the smallest integer r such that $\wedge^{r+1} \varphi = 0$ (identically on S). In particular, $\varphi = 0$ if and only if its rank is 0.

Given any nonnegative integer k , we define the rank- k degeneration scheme of φ as the zero locus on S of the morphism $\wedge^{k+1} \varphi$. If the rank of φ is r , we abbreviate its rank- $(r-1)$ degeneration scheme to just degeneration scheme (the degeneration scheme of the zero morphism is empty).

The morphism $\varphi : \mathcal{E} \rightarrow \mathcal{F}$ is generically surjective if its rank on every irreducible component of S equals the rank of \mathcal{F} . The cokernel of a generically surjective morphism φ is a torsion sheaf supported on the degeneration scheme of φ .

If φ has rank r , for any \mathbf{K} -point s of S , the rank of the \mathbf{K} -linear map φ_s is at most r . The converse may however not be true: if $S = \text{Spec}(\mathbf{k}[x]/x^2)$, $\mathcal{E} = \mathcal{F} = \mathcal{O}_S$, and $\varphi = x$, the rank of φ is 1 and φ is generically surjective, but $\varphi_s = 0$ at all points s of S . Its degeneration scheme is S_{red} .

Lemma 2.5. *Let $\varphi : \mathcal{E} \rightarrow \mathcal{F}$ be a morphism of positive rank r between vector bundles on a scheme S . Assume that $\bigwedge^{r-1} \varphi_s$ does not vanish for any geometric point s of S .*

The degeneration scheme of φ then equals the scheme-theoretic support of the sheaf $\text{Coker}(\varphi)$, that is, the subscheme defined by the annihilator ideal of $\text{Coker}(\varphi)$. Moreover, the sheaf $\text{Coker}(\varphi)$ is isomorphic to the pushforward of a line bundle on this subscheme.

Proof. For any \mathbf{K} -point s of S , one of the $(r - 1) \times (r - 1)$ -minors of φ does not vanish in \mathbf{K} , hence it spans the local ring $\mathcal{O}_{S,s}$; this means that the first Fitting ideal of $\text{Coker}(\varphi)$ (generated by the $(r - 1) \times (r - 1)$ -minors of φ) is trivial [10, Corollary-Definition 20.4]. By [10, Proposition 20.7], the zeroth Fitting ideal (which defines the degeneration scheme of φ) is then equal to the annihilator of $\text{Coker}(\varphi)$.

To prove the second part, we base change to the support of $\text{Coker}(\varphi)$. By [10, Corollary 20.5], the first Fitting ideal of $\text{Coker}(\varphi)$ is still trivial, while the zeroth Fitting ideal is equal to zero; [10, Proposition 20.8] then implies that $\text{Coker}(\varphi)$ is a line bundle. □

Lemma 2.5 can also be proved by the argument of Proposition 2.8.

Lemma 2.6. *Let $\varphi : \mathcal{E} \rightarrow \mathcal{F}$ be a morphism between vector bundles of rank r on a scheme S . Assume that the degeneration scheme of φ is a Cartier divisor D on S .*

We have $\text{Ker}(\varphi) = 0$ and $\text{Coker}(\varphi)$ is supported scheme-theoretically on D .

Proof. Let $\mathcal{L} := \det(\mathcal{E}^\vee) \otimes \det(\mathcal{F})$. By definition of the degeneration scheme, the ideal of D is generated by $\det(\varphi)$, so the assumption that D be a Cartier divisor means that $\det(\varphi)$, viewed as a section of \mathcal{L} , is not a zero divisor. Consider the diagram

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & \text{Ker}(\varphi) & \longrightarrow & \mathcal{E} & \xrightarrow{\varphi} & \mathcal{F} & \longrightarrow & \text{Coker}(\varphi) & \longrightarrow & 0 \\
 & & & & \downarrow \det(\varphi) & \swarrow \hat{\varphi} & \downarrow \det(\varphi) & & & & \\
 0 & \longrightarrow & \text{Ker}(\varphi) \otimes \mathcal{L} & \longrightarrow & \mathcal{E} \otimes \mathcal{L} & \xrightarrow{\varphi} & \mathcal{F} \otimes \mathcal{L} & \longrightarrow & \text{Coker}(\varphi) \otimes \mathcal{L} & \longrightarrow & 0,
 \end{array}$$

where $\hat{\varphi}$ is the composition

$$\mathcal{F} \simeq \bigwedge^{r-1} \mathcal{F}^\vee \otimes \det(\mathcal{F}) \xrightarrow{\bigwedge^{r-1} \varphi^\vee} \bigwedge^{r-1} \mathcal{E}^\vee \otimes \det(\mathcal{F}) \simeq \mathcal{E} \otimes \mathcal{L},$$

that is, $\widehat{\varphi}$ is the adjoint morphism of φ . In particular, the diagram commutes. It follows that the morphism induced by $\det(\varphi)$ on $\text{Ker}(\varphi)$ and $\text{Coker}(\varphi)$ is zero, hence both sheaves are supported on D . Since \mathcal{E} is torsion free, it follows that $\text{Ker}(\varphi) = 0$. \square

Lemma 2.7. *Let $\varphi : \mathcal{E} \rightarrow \mathcal{E}'$ be a morphism between vector bundles on a scheme S , which is surjective on the complement of an effective Cartier divisor, and let \mathcal{F} be a vector bundle on S . Then the map $\text{Hom}(\mathcal{E}', \mathcal{F}) \xrightarrow{\circ\varphi} \text{Hom}(\mathcal{E}, \mathcal{F})$ is injective. In other words, if a morphism $\psi : \mathcal{E} \rightarrow \mathcal{F}$ factors through \mathcal{E}' , such a factorization is unique.*

Proof. Let \mathcal{C} be the cokernel of φ . By the left exactness of the Hom functor, it is enough to show that $\text{Hom}(\mathcal{C}, \mathcal{F}) = 0$. The question is local, so we may assume $\mathcal{F} = \mathcal{O}_S$. Moreover, since the degeneration scheme is contained in an effective Cartier divisor, we may assume that \mathcal{C} is annihilated by a regular function f on S which is not a zero divisor. But the image of any morphism $\mathcal{C} \rightarrow \mathcal{O}_S$ is then annihilated by f , hence is zero. \square

If we do not assume that the degeneration scheme of φ is contained in an effective Cartier divisor, the conclusion of Lemma 2.7 may not hold. For example, let $S = \text{Spec}(\mathbf{k}[x, y]/(xy, y^2))$, $\mathcal{E} = \mathcal{O}_S \oplus \mathcal{O}_S$, $\mathcal{E}' = \mathcal{F} = \mathcal{O}_S$, and $\varphi = (x, y)$. Consider the nonzero map $\mathcal{E}' \rightarrow \mathcal{F}$ given by y ; its composition with φ is zero. The degeneration scheme of φ is defined by the maximal ideal (x, y) ; it is a Weil divisor, but not a Cartier divisor.

2.3. Canonical factorization

The following canonical factorization of a morphism between vector bundles seems to be little known, but it will be crucial for our construction.

Proposition 2.8. *Let $\varphi : \mathcal{E} \rightarrow \mathcal{F}$ be a morphism of positive rank r between vector bundles on a scheme S . Assume that its degeneration scheme is a Cartier divisor D on S and that $\bigwedge^{r-1} \varphi_s$ does not vanish for any geometric point s of S . There is a unique factorization*

$$\varphi : \mathcal{E} \rightarrow \mathcal{E}_1 \xrightarrow{\varphi_1} \mathcal{F}_1 \hookrightarrow \mathcal{F}$$

such that

- \mathcal{E}_1 and \mathcal{F}_1 are vector bundles of rank r ,
- the map $\mathcal{E} \rightarrow \mathcal{E}_1$ is an epimorphism,
- the map $\mathcal{F}_1 \hookrightarrow \mathcal{F}$ is a fiberwise monomorphism,
- the map φ_1 is a monomorphism and its cokernel is a line bundle on D .

Proof. For any such factorization, φ_1 is injective by Lemma 2.6, hence \mathcal{E}_1 is the image of φ and \mathcal{F}_1 is the dual of the image of φ^\vee , so the uniqueness is clear. It is

therefore enough to prove the proposition locally so we may assume that \mathcal{E} and \mathcal{F} are trivial vector bundles and that φ is given by a matrix of regular functions on S .

Let s be a geometric point on S . The rank of φ_s is either r or $r - 1$. If it is r , one of the $r \times r$ -minors of φ is nonzero at s , hence is invertible in a neighborhood of s . Restricting to such a neighborhood and considering appropriate bases for the fibers \mathcal{E}_s and \mathcal{F}_s , we may assume that the minor corresponds to the first r vectors in each basis. In other words, the matrix has the form

$$\varphi = \begin{pmatrix} \varphi_{1,1} & \varphi_{1,2} \\ \varphi_{2,1} & \varphi_{2,2} \end{pmatrix},$$

where $\varphi_{1,1}$ is a square matrix of size r with invertible determinant. The matrix $\varphi_{1,1}$ is therefore invertible and, upon multiplying it by its inverse, we may assume that it is the identity matrix I_r . Applying elementary transformations to rows and columns, we may assume that $\varphi_{1,2} = \varphi_{2,1} = 0$. The entries of the matrix $\varphi_{2,2}$ are then $(r + 1) \times (r + 1)$ -minors of the matrix φ , hence they all vanish. In a neighborhood of s , the map φ can therefore be written as a composition of the epimorphism of \mathcal{E} onto the trivial vector bundle of rank r (corresponding to the first r basis vectors) and its fiberwise monomorphism into \mathcal{F} . In particular, φ_1 is an isomorphism.

If the rank of φ_s is $r - 1$, restricting to a neighborhood of s and choosing bases of \mathcal{E}_s and \mathcal{F}_s appropriately, we may assume that φ is in the form as above, but where now $\varphi_{1,1} = I_{r-1}$ and $\varphi_{1,2} = \varphi_{2,1} = 0$. Again, the entries of $\varphi_{2,2}$ are the $r \times r$ -minors of φ , hence they generate the ideal generated by the equation f of the Cartier divisor D . Therefore, we can write $\varphi_{2,2} = f\varphi'_{2,2}$; the ideal generated by the entries of $\varphi'_{2,2}$ is trivial, hence the matrix $\varphi'_{2,2}$ vanishes nowhere.

On the other hand, the 2×2 -minors of the matrix $\varphi_{2,2}$ are equal to (some) $(r + 1) \times (r + 1)$ -minors of φ , hence they all vanish identically; therefore (recall that f is not a zero divisor), the same is true for the matrix $\varphi'_{2,2}$. Applying the same arguments as above, we can assume the matrix $\varphi_{2,2}$ has top left entry f and all other entries 0. In a neighborhood of s , the map φ can thus be written as the composition of the epimorphism of \mathcal{E} onto the trivial vector bundle $\mathcal{O}^{\oplus r}$ (corresponding to the first r basis vectors), a map φ_1 given by the diagonal matrix $\text{diag}(1, \dots, 1, f)$, and a fiberwise monomorphism from $\mathcal{O}^{\oplus r}$ into \mathcal{F} . In particular, φ_1 is a monomorphism and its cokernel is (locally) the structure sheaf of D . \square

2.4. Families of quadratic forms and residual families

In this section, we assume that the characteristic of the base field \mathbf{k} is not 2. Let \mathcal{E} and \mathcal{V} be vector bundles of respective ranks r and k on a scheme S and let

$$\mathbf{q} : \mathcal{V} \rightarrow \text{Sym}^2 \mathcal{E}^\vee \tag{5}$$

be a morphism of sheaves. We may think of \mathbf{q} as a family $\mathcal{E} \otimes \mathcal{E} \rightarrow \mathcal{V}^\vee$ of quadratic forms on \mathcal{E} with values in \mathcal{V}^\vee .

When \mathcal{V} has rank 1, we define the discriminant subscheme $\text{Disc}(\mathbf{q}) \subset S$ of the family \mathbf{q} as the degeneration scheme of the associated morphism

$$\mathcal{E} \xrightarrow{\mathbf{q}} \mathcal{E}^\vee \otimes \mathcal{V}^\vee$$

of rank- r vector bundles, that is, the zero locus of the induced section $\det(\mathbf{q})$ of the line bundle $\det(\mathcal{E}^\vee)^{\otimes 2} \otimes (\mathcal{V}^\vee)^{\otimes r}$.

Let $i : D \hookrightarrow S$ be an effective Cartier divisor. Let $\mathcal{K} \subset i^*\mathcal{E}$ be a line subbundle which is contained in the kernel of the quadratic forms $i^*\mathbf{q}$. In other words, the composition

$$i^*\mathcal{V} \xrightarrow{i^*\mathbf{q}} \text{Sym}^2(i^*\mathcal{E}^\vee) \rightarrow i^*\mathcal{E}^\vee \otimes \mathcal{K}^\vee$$

vanishes (when \mathcal{V} has rank 1, this implies that D is contained in the discriminant $\text{Disc}(\mathbf{q})$). In this situation, we construct a family of quadratic forms on the line bundle \mathcal{K} over D as follows.

Let $\widetilde{\mathcal{K}} \subset \mathcal{E}$ be a local extension over an open subset of S of the line subbundle $\mathcal{K} \subset i^*\mathcal{E}$. By postcomposition, the family of quadratic forms \mathbf{q} induces a map $\mathcal{V} \rightarrow \text{Sym}^2\mathcal{E}^\vee \rightarrow \text{Sym}^2\widetilde{\mathcal{K}}^\vee$ which by our assumption vanishes on the divisor D , hence factors through a map

$$\mathcal{V}(D) \rightarrow \text{Sym}^2\widetilde{\mathcal{K}}^\vee.$$

Restricting it to D , we get a map

$$\mathcal{V}(D)|_D \rightarrow \text{Sym}^2\mathcal{K}^\vee$$

which we call the residual family of quadratic forms.

Lemma 2.9. *The residual family of quadratic forms on D is independent of the choices made.*

If the rank of \mathcal{V} is 1 and $\text{Disc}(\mathbf{q}) = D$ as subschemes of S , the rank of \mathbf{q} on D is equal to $r - 1$ and the residual family of quadratic forms vanishes nowhere on D .

Proof. Let s_0 be a section of \mathcal{E} extending locally a section generating \mathcal{K} . Any other extension can be written as $s_0 + ts$ for some section s of \mathcal{E} , where t is a local equation of the divisor D . The evaluation of \mathbf{q} on this section is equal to

$$\mathbf{q}(s_0 + ts, s_0 + ts) = \mathbf{q}(s_0, s_0) + 2t\mathbf{q}(s_0, s) + t^2\mathbf{q}(s, s).$$

The factorization through $\mathcal{V}(D)$ is then given by

$$\frac{1}{t}(\mathbf{q}(s_0, s_0) + 2t\mathbf{q}(s_0, s) + t^2\mathbf{q}(s, s)) \Big|_{t=0} = \left(\frac{1}{t}\mathbf{q}(s_0, s_0) + 2\mathbf{q}(s_0, s) \right) \Big|_{t=0}.$$

It remains to note that $\mathbf{q}(s_0, s)$ vanishes on D since s_0 is in the kernel of $i^*\mathbf{q}$.

Let us choose local trivializations of \mathcal{E} and \mathcal{V} such that \mathbf{q} corresponds to a k -tuple of symmetric matrices $(q_{ij}^\alpha)_{1 \leq i, j \leq r}$ of regular functions on S (where $\alpha \in \{1, \dots, k\}$) and the section s_0 (defining a local extension $\widetilde{\mathcal{K}}$ of \mathcal{K}) corresponds

to the first basis vector of \mathcal{E} . The condition that \mathcal{K} be in the kernel of $i^*\mathbf{q}$ then means

$$q_{1i}^\alpha = t\bar{q}_{1i}^\alpha \quad \text{for all } 1 \leq i \leq r \text{ and all } 1 \leq \alpha \leq k, \tag{6}$$

and the residual quadric is just $\bar{q}_{11}^\alpha|_D$.

When the rank of \mathcal{V} is one (so we have just one symmetric matrix (q_{ij})), we have

$$\det(q_{ij}) \equiv t\bar{q}_{11} \det(q_{i,j})_{2 \leq i,j \leq r} \pmod{t^2}.$$

The condition $\text{Disc}(\mathbf{q}) = D$ implies that the last factor $\det(q_{i,j})_{2 \leq i,j \leq r}$ is invertible (hence the rank of \mathbf{q} on D is $r - 1$) and \bar{q}_{11} is invertible on D too, so that the residual family of quadratic forms vanishes nowhere. \square

2.5. Hecke transform of a family of quadratic forms

We continue working in the set-up of the previous section. We keep the assumption $\text{char}(\mathbf{k}) \neq 2$. The construction presented here (which we call Hecke transform) is a generalization of the construction from [22, Lemma 1.14].

Definition 2.10. An effective Cartier divisor D on a scheme S is a **double** if there is an effective Cartier divisor E on S such that $D = 2E$, that is, the ideal of D is the square of the ideal of E .

Assume that the effective Cartier divisor D considered in the previous section is a double and write $D = 2E$. For any line subbundle $\mathcal{K} \subset \mathcal{E}|_D$, we set

$$\mathcal{K}_E := \mathcal{K}|_E, \quad \mathcal{K}_E^\vee := \mathcal{K}^\vee|_E, \quad \mathcal{E}_D := \mathcal{E}|_D, \quad \mathcal{E}_E := \mathcal{E}|_E.$$

Since \mathcal{K} is a line bundle on D and E is a Cartier divisor on S , the kernel of the natural epimorphism $\mathcal{E}^\vee \rightarrow \mathcal{K}_E^\vee$ is a vector bundle (Lemma 2.4). We denote by $\tilde{\mathcal{E}}$ its dual, so that there is an exact sequence

$$0 \rightarrow \tilde{\mathcal{E}}^\vee \rightarrow \mathcal{E}^\vee \rightarrow \mathcal{K}_E^\vee \rightarrow 0 \tag{7}$$

of sheaves on S , whose dual sequence can be written as

$$0 \rightarrow \mathcal{E} \rightarrow \tilde{\mathcal{E}} \rightarrow \mathcal{K}_E(E) \rightarrow 0. \tag{8}$$

The following lemma will be very useful later. We keep the notation of Sec. 2.4.

Lemma 2.11. *Assume $D = 2E$. Let $\mathcal{K} \subset \mathcal{E}_D$ be a line subbundle contained in the kernel of $\mathbf{q}|_D$. Define the vector bundle $\tilde{\mathcal{E}}$ by the exact sequence (7). The family of quadratic forms $\mathbf{q} : \mathcal{V} \rightarrow \text{Sym}^2 \mathcal{E}^\vee$ then factors through $\text{Sym}^2 \tilde{\mathcal{E}}^\vee$ in a unique way.*

Proof. We can use a representation of \mathbf{q} by a matrix (q_{ij}^α) as in the proof of Lemma 2.9 with the same conventions on the coordinates, assuming in particular that (6) holds. Let u be an equation of E in S , so that the equation of D can be

written as $t = u^2$. The sequence (7) can then be written in local coordinates on S as

$$0 \rightarrow \mathcal{O}_S^{\oplus r} \xrightarrow{\text{diag}(u, 1, \dots, 1)} \mathcal{O}_S^{\oplus r} \longrightarrow \mathcal{O}_E \rightarrow 0. \tag{9}$$

The factorization condition that we want to prove just means that q_{11}^α is divisible by u^2 and $q_{12}^\alpha, \dots, q_{1r}^\alpha$ are divisible by u ; since $t = u^2$, this follows from (6). The uniqueness of the factorization follows from Lemma 2.7 applied to the symmetric square of (8). \square

We denote by $\tilde{\mathbf{q}} : \mathcal{V} \rightarrow \text{Sym}^2 \tilde{\mathcal{E}}^\vee$ the induced map and call it the Hecke transform of \mathbf{q} with respect to the line subbundle $\mathcal{K} \subset \mathcal{E}_D$.

Proposition 2.12. *Let \mathcal{E} and \mathcal{V} be vector bundles of respective ranks r and k on a scheme S . Let $\mathbf{q} : \mathcal{V} \rightarrow \text{Sym}^2 \mathcal{E}^\vee$ be a family of quadratic forms on \mathcal{E} with values in \mathcal{V}^\vee . Assume finally that there exist a double Cartier divisor $D = 2E$ and a line subbundle $\mathcal{K} \subset \mathcal{E}_D$ on D which is contained in the kernel of $\mathbf{q}|_D$. Let $\tilde{\mathbf{q}} : \mathcal{V} \rightarrow \text{Sym}^2 \tilde{\mathcal{E}}^\vee$ be the Hecke transform of \mathbf{q} with respect to \mathcal{K} .*

- (a) *The restriction of the sequence (8) to E splits and gives a canonical direct sum decomposition*

$$\tilde{\mathcal{E}}_E \simeq (\mathcal{E}_E / \mathcal{K}_E) \oplus \mathcal{K}_E(E) \tag{10}$$

of the restriction $\tilde{\mathcal{E}}_E$ of $\tilde{\mathcal{E}}$ to E .

- (b) *The summands of (10) are mutually orthogonal with respect to the quadratic form $\tilde{\mathbf{q}}|_E$. Moreover, the restriction of $\tilde{\mathbf{q}}|_E$ to the first summand of (10) is induced by $\mathbf{q}|_E$ and the restriction of $\tilde{\mathbf{q}}|_E$ to the second summand is the residual family of quadratic forms for \mathbf{q} .*

Proof. Consider the vector bundle $\hat{\mathcal{E}}$ on S defined as the dual of the kernel of the natural map $\mathcal{E}^\vee \rightarrow \mathcal{K}^\vee$. We have an exact sequence

$$0 \rightarrow \mathcal{E} \rightarrow \hat{\mathcal{E}} \rightarrow \mathcal{K}(D) \rightarrow 0. \tag{11}$$

By construction, the embedding $\mathcal{E} \rightarrow \hat{\mathcal{E}}$ factors as $\mathcal{E} \rightarrow \tilde{\mathcal{E}} \rightarrow \hat{\mathcal{E}}$ and the map $\tilde{\mathcal{E}} \rightarrow \hat{\mathcal{E}}$ fits into the exact sequence

$$0 \rightarrow \tilde{\mathcal{E}} \rightarrow \hat{\mathcal{E}} \rightarrow \mathcal{K}_E(2E) \rightarrow 0. \tag{12}$$

Restricting (8), (12), and (11) to E , we obtain exact sequences

$$0 \rightarrow \mathcal{K}_E \rightarrow \mathcal{E}_E \rightarrow \tilde{\mathcal{E}}_E \rightarrow \mathcal{K}_E(E) \rightarrow 0, \tag{13}$$

$$0 \rightarrow \mathcal{K}_E(E) \rightarrow \tilde{\mathcal{E}}_E \rightarrow \hat{\mathcal{E}}_E \rightarrow \mathcal{K}_E(2E) \rightarrow 0, \tag{14}$$

$$0 \rightarrow \mathcal{K}_E \rightarrow \mathcal{E}_E \rightarrow \hat{\mathcal{E}}_E \rightarrow \mathcal{K}_E(2E) \rightarrow 0. \tag{15}$$

Since the composition of the middle arrows of (13) and (14) is the middle arrow of (15), the composition $\mathcal{K}_E(E) \rightarrow \tilde{\mathcal{E}}_E \rightarrow \mathcal{K}_E(E)$ is an isomorphism, hence the sheaf $\mathcal{K}_E(E)$ is a direct summand of $\tilde{\mathcal{E}}_E$. The sequence (13) identifies the other summand with $\mathcal{E}_E/\mathcal{K}_E$. This proves (a).

Let us prove (b). The first map in (8) restricted to E can be written as a composition

$$\mathcal{E}_E \twoheadrightarrow \mathcal{E}_E/\mathcal{K}_E \hookrightarrow \tilde{\mathcal{E}}_E.$$

Taking the symmetric square of these morphisms and dualizing, we see that the map $\mathrm{Sym}^2 \tilde{\mathcal{E}}_E^\vee \rightarrow \mathrm{Sym}^2 \mathcal{E}_E^\vee$ can be written as the composition

$$\mathrm{Sym}^2 \tilde{\mathcal{E}}_E^\vee \rightarrow \mathrm{Sym}^2 (\mathcal{E}_E/\mathcal{K}_E)^\vee \hookrightarrow \mathrm{Sym}^2 \mathcal{E}_E^\vee.$$

By Lemma 2.11, the quadric $\mathbf{q}|_E$ (considered as a map from \mathcal{V} to $\mathrm{Sym}^2 \mathcal{E}_E^\vee$) factors through $\mathrm{Sym}^2 \tilde{\mathcal{E}}_E^\vee$ as $\tilde{\mathbf{q}}|_E$. Hence it a fortiori factors through the middle term. Such a factorization is nothing but the induced family of quadratic forms on $\mathcal{E}_E/\mathcal{K}_E$ and the image of $\tilde{\mathbf{q}}|_E$ is the restriction of $\mathbf{q}|_E$ to $\mathcal{E}_E/\mathcal{K}_E$. These two families of quadratic forms therefore coincide.

We now show that the summands in (10) are mutually orthogonal and that the restriction of $\tilde{\mathbf{q}}|_E$ to the summand $\mathcal{K}_E(E)$ is given by the residual family of quadratic forms. The question is local, so we can assume that \mathcal{E} , \mathcal{K} , and \mathcal{V} are trivialized as in the proof of Lemma 2.9. Under these assumptions, the sequence (7) can be rewritten as in (9). This means that the matrix of $\tilde{\mathbf{q}}$ is

$$\begin{pmatrix} q_{11}^\alpha/u^2 & q_{12}^\alpha/u & \cdots & q_{1r}^\alpha/u \\ q_{21}^\alpha/u & q_{22}^\alpha & \cdots & q_{2r}^\alpha \\ \vdots & \vdots & \ddots & \vdots \\ q_{r1}^\alpha/u & q_{r2}^\alpha & \cdots & q_{rr}^\alpha \end{pmatrix}.$$

In particular, its restriction to the summand $\mathcal{K}_E(E)$ is given by the regular function $(q_{11}^\alpha/u^2)|_E = (q_{11}^\alpha/t)|_E$, which is the residual family of quadratic forms. Moreover, if we set $\bar{q}_{1i}^\alpha := q_{1i}^\alpha/t$ for all $i > 1$ as in (6), we have $(q_{1i}^\alpha/u)|_E = (u\bar{q}_{1i}^\alpha)|_E = 0$, hence the summands in (10) are mutually orthogonal. \square

3. The Moduli Stack of Smooth GM Varieties

In this section, we introduce the stack of GM varieties and the closely related stacks of GM and Lagrangian data. We mostly work in the étale topology, but one can also work with the fppf topology. From now on, we assume that the characteristic of the base field \mathbf{k} is zero.

3.1. The stack of GM varieties

We start with a definition of the stack of GM varieties.

Definition 3.1. A family of smooth polarized GM varieties of dimension n over a scheme S is a pair $(\mathcal{X} \rightarrow S, \mathcal{H})$, where

- $\pi_{\mathcal{X}} : \mathcal{X} \rightarrow S$ is a smooth and proper morphism,
- $\mathcal{H} \in \text{Pic}_{\mathcal{X}/S}(S)$ is a relative $\pi_{\mathcal{X}}$ -ample divisor class,

such that for every geometric point s of S ,

- the pair $(\mathcal{X}_s, \mathcal{H}|_{\mathcal{X}_s})$ is a smooth polarized GM variety of dimension n in the sense of [6, Definition 2.1].

A morphism of families of GM varieties from $(\mathcal{X} \rightarrow S, \mathcal{H})$ to $(\mathcal{X}' \rightarrow S', \mathcal{H}')$ is a pair (f, φ) giving a Cartesian square

$$\begin{array}{ccc}
 \mathcal{X} & \xrightarrow{\varphi} & \mathcal{X}' \\
 \pi_{\mathcal{X}} \downarrow & & \downarrow \pi_{\mathcal{X}'} \\
 S & \xrightarrow{f} & S'
 \end{array} \tag{16}$$

such that $\mathcal{H} = \varphi^* \mathcal{H}'$ in the relative Picard group $\text{Pic}_{\mathcal{X}/S}(S)$.

Families of smooth polarized GM varieties of dimension n form a category fibered in groupoids over the category Sch/\mathbf{k} of schemes over \mathbf{k} ; we denote it by $\mathfrak{M}_n^{\text{GM}}$.

Smooth GM varieties exist in each dimension $n \in \{1, \dots, 6\}$. A GM variety of dimension 1 is a Clifford general curve of genus 6 (see [6, Proposition 2.12]), a GM variety of dimension 2 is a Brill–Noether general polarized K3 surface of genus 6 (see [6, Proposition 2.13]), and their moduli stacks are well studied. Accordingly, we will concentrate in this paper on GM varieties of dimension $n \in \{3, 4, 5, 6\}$. There is then an isomorphism

$$\text{Pic}_{\mathcal{X}/S} \simeq \underline{\mathbf{Z}}$$

between étale sheaves (see [6, Lemma 2.29]); over a connected scheme S , there is therefore a unique choice of a relative divisor class \mathcal{H} .

Proposition 3.2 ([13, Proposition A.2]). *For $n \in \{2, \dots, 6\}$, the fibered category $\mathfrak{M}_n^{\text{GM}}$ is a smooth and irreducible Deligne–Mumford stack of dimension $25 - (5 - n)(6 - n)/2$. It is of finite type over \mathbf{Q} with affine diagonal of finite type.*

We call $\mathfrak{M}_n^{\text{GM}}$ the moduli stack of smooth polarized GM varieties of dimension n . We will see later (Theorem 5.11) that $\mathfrak{M}_n^{\text{GM}}$ is separated.

3.2. The stack of GM data

GM data sets over a field were defined in [6, Definition 2.5]. It will be convenient to change the definition slightly as follows.

Definition 3.3. A normalized family of GM data of dimension n over a scheme S is a collection $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$, where

- \mathcal{W} , \mathcal{V}_6 , and \mathcal{V}_5 are vector bundles on S of respective ranks $n + 5$, 6, and 5,
- $\mathcal{V}_5 \hookrightarrow \mathcal{V}_6$ is a fiberwise monomorphism,
- $\mu : \mathcal{W} \rightarrow \wedge^2 \mathcal{V}_5 \otimes (\mathcal{V}_6/\mathcal{V}_5)$ and
- $\mathbf{q} : \mathcal{V}_6 \rightarrow \text{Sym}^2 \mathcal{W}^\vee \otimes \det(\mathcal{V}_5) \otimes (\mathcal{V}_6/\mathcal{V}_5)^{\otimes 2}$ are morphisms between vector bundles,

such that the diagram

$$\begin{array}{ccc}
 \mathcal{V}_5 \otimes \text{Sym}^2 \mathcal{W} & \hookrightarrow & \mathcal{V}_6 \otimes \text{Sym}^2 \mathcal{W} \\
 \text{Sym}^2 \mu \downarrow & & \downarrow \mathbf{q} \\
 \mathcal{V}_5 \otimes \text{Sym}^2 (\wedge^2 \mathcal{V}_5) \otimes (\mathcal{V}_6/\mathcal{V}_5)^{\otimes 2} & \xrightarrow{\wedge} & \det(\mathcal{V}_5) \otimes (\mathcal{V}_6/\mathcal{V}_5)^{\otimes 2}
 \end{array} \tag{17}$$

commutes (the bottom arrow is given by wedge product).

A morphism of normalized families of GM data $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ and $(S', \mathcal{W}', \mathcal{V}'_6, \mathcal{V}'_5, \mu', \mathbf{q}')$ over schemes S and S' is a morphism $f : S \rightarrow S'$ and isomorphisms

$$\varphi_V : \mathbf{P}_S(\mathcal{V}_6) \xrightarrow{\sim} \mathbf{P}_{S'}(f^* \mathcal{V}'_6) \quad \text{and} \quad \varphi_W : \mathbf{P}_S(\mathcal{W}) \xrightarrow{\sim} \mathbf{P}_{S'}(f^* \mathcal{W}')$$

such that $\varphi_V(\mathbf{P}_S(\mathcal{V}_5)) = \mathbf{P}_{S'}(f^* \mathcal{V}'_5)$ and the following diagrams commute:

$$\begin{array}{ccc}
 \mathbf{P}_S(\mathcal{W}) & \xrightarrow{\varphi_W} & \mathbf{P}_{S'}(f^* \mathcal{W}') \\
 \downarrow \mu & & \downarrow \mu' \\
 \mathbf{P}_S(\wedge^2 \mathcal{V}_5) & \xrightarrow{\wedge^2 \varphi_V} & \mathbf{P}_{S'}(f^* \wedge^2 \mathcal{V}'_5)
 \end{array}
 \quad \text{and} \quad
 \begin{array}{ccc}
 \mathbf{P}_S(\mathcal{V}_6) & \xrightarrow{\varphi_V} & \mathbf{P}_{S'}(f^* \mathcal{V}'_6) \\
 \downarrow \mathbf{q} & & \downarrow \mathbf{q}' \\
 \mathbf{P}_S(\text{Sym}^2 \mathcal{W}^\vee) & \xleftarrow{\text{Sym}^2 \varphi_W^\vee} & \mathbf{P}_{S'}(f^* \text{Sym}^2 \mathcal{W}'^\vee)
 \end{array}$$

It is sometimes convenient to express the commutativity of (17) as the equality

$$\mathbf{q}(v)(w_1, w_2) = v \wedge \mu(w_1) \wedge \mu(w_2) \quad \text{on } \mathcal{V}_5 \otimes \text{Sym}^2 \mathcal{W}. \tag{18}$$

Families of normalized GM data of dimension n form a category fibered in groupoids over the category Sch/\mathbf{k} of schemes over \mathbf{k} ; we denote it by $\mathfrak{M}_n^{\text{GM-data}}$.

Remark 3.4. One could alternatively define morphisms of families of GM data to be triples $(f, \tilde{\varphi}_V, \tilde{\varphi}_W)$, where f is a morphism $S \rightarrow S'$ and

$$\tilde{\varphi}_V : \mathcal{V}_6 \xrightarrow{\sim} f^* \mathcal{V}'_6 \quad \text{and} \quad \tilde{\varphi}_W : \mathcal{W} \xrightarrow{\sim} f^* \mathcal{W}'$$

are isomorphisms compatible with the subbundle $\mathcal{V}_5 \subset \mathcal{V}_6$ and the morphisms μ and \mathbf{q} . This also defines a category fibered in groupoids over Sch/\mathbf{k} , which we denote by $\widetilde{\mathfrak{M}}_n^{\text{GM-data}}$ and call the category of linearized GM data.

Denoting by $\widetilde{\text{Aut}}(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ the automorphism group scheme in $\widetilde{\mathfrak{M}}_n^{\text{GM-data}}$, we obtain an embedding of group schemes

$$\begin{aligned} \mathbb{G}_m(S) &\hookrightarrow \widetilde{\text{Aut}}(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q}) \\ u &\mapsto (\tilde{\varphi}_V = u, \tilde{\varphi}_W = u^3). \end{aligned} \tag{19}$$

We have

$$\text{Aut}(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q}) \simeq \widetilde{\text{Aut}}(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q}) / \mathbb{G}_m(S),$$

which essentially means that the fibered category of GM data is the *rigidification* [1, 2] of the fibered category of linearized GM data with respect to the embeddings (19).

This observation implies that any morphism in $\mathfrak{M}_n^{\text{GM-data}}$ over $f : S \rightarrow S'$ can be locally over S' lifted to a morphism in $\widetilde{\mathfrak{M}}_n^{\text{GM-data}}$ (and such a lifting is unique up to composition with the action of $\mathbb{G}_m(S)$). We will frequently use these liftings.

Lemma 3.5. *The fibered categories $\mathfrak{M}_n^{\text{GM-data}}$ and $\widetilde{\mathfrak{M}}_n^{\text{GM-data}}$ are stacks over Sch/\mathbf{k} .*

Proof. For the fibered category $\widetilde{\mathfrak{M}}_n^{\text{GM-data}}$, this is a consequence of the fact that quasicohherent sheaves form a stack in the fppf topology: a family of linearized GM data is a collection of quasicohherent sheaves and morphisms between them that satisfy some properties that are stable under base change.

For the fibered category $\mathfrak{M}_n^{\text{GM-data}}$, use [1, Theorem 5.1.5]. □

3.3. Equivalence of stacks

The main result of this section is a relation between the stack $\mathfrak{M}_n^{\text{GM}}$ of smooth polarized GM varieties and an open substack of the stack $\mathfrak{M}_n^{\text{GM-data}}$ of normalized GM data. To define this substack, we use the notion of a GM intersection associated with a GM data set defined in [6, (2.8)].

Definition 3.6. A family $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ of normalized GM data of dimension n over a scheme S is **smooth** if for each geometric point s in S , the GM intersection

$$\bigcap_{v \in \mathcal{V}_{6,s}} \{\mathbf{q}(v) = 0\} \subset \mathbf{P}(\mathcal{W}_s)$$

corresponding to the GM data set $(\mathcal{W}_s, \mathcal{V}_{6,s}, \mathcal{V}_{5,s}, \mu_s, \mathbf{q}_s)$ is a smooth GM variety of dimension n .

By [6, Lemma 2.8], a GM intersection corresponding to a GM data set of dimension n is a smooth GM variety of dimension n if and only if the GM intersection has dimension n and is smooth. As we will see in the proof of Lemma 3.8, n is the expected dimension of the corresponding GM intersection, hence the condition for the GM intersection to be a smooth GM variety of dimension n is open. Therefore,

families of smooth normalized GM data of dimension n are classified by an open substack of $\mathfrak{M}_n^{\text{GM-data}}$.

Theorem 3.7. *For each $n \in \{2, \dots, 6\}$, the stack of polarized GM varieties $\mathfrak{M}_n^{\text{GM}}$ is equivalent to the open substack of $\mathfrak{M}_n^{\text{GM-data}}$ classifying families of smooth normalized GM data of dimension n .*

Proof. Let $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ be a smooth family of normalized GM data over a scheme S and let $\pi_{\mathbf{P}_S(\mathcal{W})} : \mathbf{P}_S(\mathcal{W}) \rightarrow S$ be the natural projection. We have

$$\begin{aligned} & H^0(\mathbf{P}_S(\mathcal{W}), \pi_{\mathbf{P}_S(\mathcal{W})}^*(\mathcal{V}_6^\vee \otimes \det(\mathcal{V}_5) \otimes (\mathcal{V}_6/\mathcal{V}_5)^{\otimes 2}) \otimes \mathcal{O}_{\mathbf{P}_S(\mathcal{W})}(2)) \\ & \simeq H^0(S, \mathcal{V}_6^\vee \otimes \det(\mathcal{V}_5) \otimes (\mathcal{V}_6/\mathcal{V}_5)^{\otimes 2} \otimes \pi_{\mathbf{P}_S(\mathcal{W})*} \mathcal{O}_{\mathbf{P}_S(\mathcal{W})}(2)) \\ & \simeq H^0(S, \mathcal{V}_6^\vee \otimes \det(\mathcal{V}_5) \otimes (\mathcal{V}_6/\mathcal{V}_5)^{\otimes 2} \otimes \text{Sym}^2 \mathcal{W}^\vee) \\ & \simeq \text{Hom}(\mathcal{V}_6, \text{Sym}^2 \mathcal{W}^\vee \otimes \det(\mathcal{V}_5) \otimes (\mathcal{V}_6/\mathcal{V}_5)^{\otimes 2}). \end{aligned}$$

Thus, the morphism \mathbf{q} can be thought of as a global section

$$\mathbf{q} \in H^0(\mathbf{P}_S(\mathcal{W}), \pi_{\mathbf{P}_S(\mathcal{W})}^*(\mathcal{V}_6^\vee \otimes \det(\mathcal{V}_5) \otimes (\mathcal{V}_6/\mathcal{V}_5)^{\otimes 2}) \otimes \mathcal{O}_{\mathbf{P}_S(\mathcal{W})}(2)). \tag{20}$$

Consider the closed subscheme $\mathcal{X} \subset \mathbf{P}_S(\mathcal{W})$ defined as the zero locus of this global section. Define the morphism $\pi_{\mathcal{X}} : \mathcal{X} \rightarrow S$ as the restriction of the projection $\pi_{\mathbf{P}_S(\mathcal{W})} : \mathbf{P}_S(\mathcal{W}) \rightarrow S$ and the polarization \mathcal{H} on \mathcal{X} as the restriction of the hyperplane class of $\mathbf{P}_S(\mathcal{W})$. Each geometric fiber $(\mathcal{X}_s, \mathcal{H}|_{\mathcal{X}_s})$ is a smooth polarized GM variety of dimension n . Moreover, the map $\pi_{\mathcal{X}} : \mathcal{X} \rightarrow S$ is proper by definition and flat by Lemma 3.8. Since all fibers are smooth (by Definition 3.6), the map $\pi_{\mathcal{X}}$ is also smooth. Thus, $(\mathcal{X} \rightarrow S, \mathcal{H})$ is a family of smooth polarized GM varieties. This construction together with a relative version of [6, Theorem 2.3] implies

$$\mathbf{P}_S(\mathcal{W}) \simeq \mathbf{P}_S((\pi_{\mathcal{X}*} \mathcal{O}_{\mathcal{X}}(\mathcal{H}))^\vee) \quad \text{and} \quad \mathbf{P}_S(\mathcal{V}_6) \simeq \mathbf{P}_S(\pi_{\mathbf{P}_S(\mathcal{W})*} \mathcal{I}_{\mathcal{X}/\mathbf{P}_S(\mathcal{W})}(2)). \tag{21}$$

Similarly, given a morphism $(f, \varphi_W, \varphi_V)$ of families of GM data from $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ to $(S', \mathcal{W}', \mathcal{V}'_6, \mathcal{V}'_5, \mu', \mathbf{q}')$, we consider the isomorphism

$$\mathbf{P}_S(\mathcal{W}) \xrightarrow{\varphi_W} \mathbf{P}_S(f^* \mathcal{W}') \xrightarrow{\sim} \mathbf{P}_{S'}(\mathcal{W}') \times_{S'} S.$$

Since φ_V and φ_W are compatible with \mathbf{q} , it induces a morphism $\varphi : \mathcal{X} \rightarrow \mathcal{X}'$ such that (16) is a Cartesian square. Moreover, by construction, we have $\varphi^* \mathcal{H}' = \mathcal{H}$ in $\text{Pic}_{\mathcal{X}/S}(S)$. Therefore, (f, φ) is a morphism of families of GM varieties.

This means that we have defined a morphism of stacks

$$\zeta : \mathfrak{M}_{n, \text{smooth}}^{\text{GM-data}} \longrightarrow \mathfrak{M}_n^{\text{GM}}$$

from the open substack of $\mathfrak{M}_n^{\text{GM-data}}$ classifying families of smooth normalized GM data of dimension n to the stack $\mathfrak{M}_n^{\text{GM}}$ of smooth polarized GM varieties. It remains to prove that ζ is an isomorphism of stacks.

Let us check that ζ is faithful: assume that $(f_1, \varphi_{1W}, \varphi_{1V})$ and $(f_2, \varphi_{2W}, \varphi_{2V})$ are morphisms between GM data $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ and $(S', \mathcal{W}', \mathcal{V}'_6, \mathcal{V}'_5, \mu', \mathbf{q}')$ such that the corresponding morphisms (f_1, φ_1) and (f_2, φ_2) between the corresponding families of GM varieties \mathcal{X} and \mathcal{X}' are the same. Set $f := f_1 = f_2$ and $\varphi := \varphi_1 = \varphi_2$. By construction of φ , there is a commutative diagram

$$\begin{array}{ccc} \mathcal{X} & \xrightarrow{\varphi} & \mathcal{X}' \\ \downarrow & & \downarrow \\ \mathbf{P}_S(\mathcal{W}) & \xrightarrow{\varphi_{iW}} \mathbf{P}_S(f^*\mathcal{W}') & \longrightarrow \mathbf{P}(\mathcal{W}'). \end{array}$$

By (21), the fiberwise linear span of \mathcal{X} in $\mathbf{P}_S(\mathcal{W})$ is $\mathbf{P}_S(\mathcal{W})$, hence $\varphi_{1W} = \varphi_{2W}$. Furthermore, there is a commutative diagram

$$\begin{array}{ccc} \mathbf{P}_S(\mathcal{V}_6) & \xrightarrow{\varphi_{iV}} & \mathbf{P}_{S'}(\mathcal{V}'_6) \\ \mathbf{q} \downarrow & & \downarrow \mathbf{q}' \\ \mathbf{P}_S(\mathrm{Sym}^2 \mathcal{W}^\vee) & \xleftarrow{\mathrm{Sym}^2 \varphi_{iW}^\vee} & \mathbf{P}_{S'}(\mathrm{Sym}^2 \mathcal{W}'^\vee) \end{array}$$

in which the vertical arrows are embeddings (again by (21)) and the maps φ_{iV} become isomorphisms after base change to S . Since we already have $\varphi_{1W} = \varphi_{2W}$, the equality $\varphi_{1V} = \varphi_{2V}$ follows. This proves faithfulness.

Next, we check that the morphism of stacks ζ is full. Assume $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ and $(S', \mathcal{W}', \mathcal{V}'_6, \mathcal{V}'_5, \mu', \mathbf{q}')$ are families of smooth normalized GM data, let $(\mathcal{X} \rightarrow S, \mathcal{H})$ and $(\mathcal{X}' \rightarrow S', \mathcal{H}')$ be the corresponding families of GM varieties, and let (f, φ) be a morphism between them. We must show that it comes from a morphism of GM data. By the stack property and the faithfulness proved above, it is enough to prove this locally over S' . Moreover, applying base change along f , we can assume that $S' = S$ and $f = \mathrm{id}_S$. Then $\varphi: \mathcal{X} \rightarrow \mathcal{X}'$ is an isomorphism, so we can identify \mathcal{X} and \mathcal{X}' via φ .

By construction of the morphism ζ , the line bundles $\mathcal{O}_{\mathcal{X}}(\mathcal{H}) = \mathcal{O}_{\mathbf{P}_S(\mathcal{W})}(1)|_{\mathcal{X}}$ and $\mathcal{O}_{\mathcal{X}}(\mathcal{H}') = \mathcal{O}_{\mathbf{P}_S(\mathcal{W}')} (1)|_{\mathcal{X}}$ agree up to the pullback of a line bundle on S . Shrinking S if necessary, we can assume that this line bundle is trivial, so we can choose an isomorphism

$$\varphi_H : \mathcal{O}_{\mathcal{X}}(\mathcal{H}) \xrightarrow{\sim} \mathcal{O}_{\mathcal{X}}(\mathcal{H}').$$

Using the formulas (21), we see that φ_H induces isomorphisms of projective bundles $\varphi_W : \mathbf{P}_S(\mathcal{W}) \xrightarrow{\sim} \mathbf{P}_S(\mathcal{W}')$ and $\varphi_V : \mathbf{P}_S(\mathcal{V}_6) \xrightarrow{\sim} \mathbf{P}_S(\mathcal{V}'_6)$. It is easy to see that these isomorphisms are compatible with the subbundle \mathcal{V}_5 and the morphisms μ and \mathbf{q} , so that $(\mathrm{id}_S, \varphi_V, \varphi_W)$ is an isomorphism of GM data. Moreover, the isomorphism of GM varieties that it induces coincides with the one we started from. This proves fullness.

Finally, we check that ζ is essentially surjective. Given a family $(\mathcal{X} \rightarrow S, \mathcal{H})$ of smooth polarized GM varieties, we need to construct a family of smooth normalized

GM data $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ such that $(\mathcal{X} \rightarrow S, \mathcal{H}) = \zeta(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$. Since we are dealing with stacks and since we already proved that ζ is fully faithful, it is enough to construct this locally over S . So we can assume that \mathcal{H} is the class of a line bundle on \mathcal{X} . Denoting it by $\mathcal{O}_{\mathcal{X}}(\mathcal{H})$ and following the proof of [6, Theorem 2.3], we set

$$\begin{aligned} \widehat{\mathcal{W}} &:= (\pi_{\mathcal{X}*} \mathcal{O}_{\mathcal{X}}(\mathcal{H}))^\vee, & \mathcal{V}_6 &:= \pi_{\mathbf{P}_S(\widehat{\mathcal{W}})*}(\mathcal{I}_{\mathcal{X}}(2)), \\ \mathcal{L} &:= \pi_{\mathcal{X}*}(\wedge^2 \mathcal{U}_{\mathcal{X}} \otimes \mathcal{O}_{\mathcal{X}}(\mathcal{H})). \end{aligned} \tag{22}$$

These are vector bundles of respective ranks $n + 5$, 6, and 1 on S .

To be more precise, we first define the bundle $\widehat{\mathcal{W}}$ by the first equality in (22) and note that the natural rational map $\mathcal{X} \dashrightarrow \mathbf{P}_S(\widehat{\mathcal{W}})$ is regular and a closed embedding (both statements can be verified fiberwise and follow from [6, Theorem 2.3]).

Then, we define the bundle \mathcal{V}_6 by the second equality in (22) (here $\mathcal{I}_{\mathcal{X}}(2)$ is the twist of the ideal sheaf of \mathcal{X} in $\mathbf{P}_S(\widehat{\mathcal{W}})$ by the square of the hyperplane bundle $\mathcal{O}_{\mathbf{P}_S(\widehat{\mathcal{W}})}(1)$).

Finally, we let $\mathcal{U}_{\mathcal{X}}$ be the excess conormal bundle for the closed embedding $\mathcal{X} \rightarrow \mathbf{P}_S(\widehat{\mathcal{W}})$ (see [6, Definition A.1]) and define the line bundle \mathcal{L} by the third equality in (22). By [6, Theorem 2.3] again, there is a natural fiberwise monomorphism $\mathcal{U}_{\mathcal{X}} \rightarrow \pi_{\mathcal{X}}^* \mathcal{V}_6$ inducing a regular map $X \rightarrow \mathrm{Gr}_S(2, \mathcal{V}_6)$ and this map factors through $\mathrm{Gr}_S(2, \mathcal{V}_5)$ for a unique subbundle $\mathcal{V}_5 \subset \mathcal{V}_6$ of rank 5.

Let us renormalize the bundle $\widehat{\mathcal{W}}$ by setting

$$\mathcal{W} := \widehat{\mathcal{W}} \otimes (\mathcal{V}_6/\mathcal{V}_5) \otimes \mathcal{L}.$$

In order to construct the morphisms μ and \mathbf{q} , we use again the construction of [6, Theorem 2.3] in a relative setting, which produces maps

$$\widehat{\mu} : \mathcal{L} \otimes \widehat{\mathcal{W}} \rightarrow \wedge^2 \mathcal{V}_5, \quad \widehat{\mathbf{q}} : \mathcal{V}_6 \rightarrow \mathrm{Sym}^2 \widehat{\mathcal{W}}^\vee, \quad \varepsilon : \det(\mathcal{V}_5) \xrightarrow{\sim} \mathcal{L}^{\otimes 2}.$$

We then set

$$\mu = \widehat{\mu} \otimes \mathrm{id}_{\mathcal{V}_6/\mathcal{V}_5}, \quad \mathbf{q} = \varepsilon^{-1} \circ \widehat{\mathbf{q}}.$$

The relation (18) is equivalent to the relation [6, (2.7)] (with W replaced by $\widehat{\mathcal{W}}$), which is proved in [6, Lemma 2.7]. Thus, we obtain a family of normalized GM data on S . Finally, by [6, Theorem 2.3] again, the family of GM varieties corresponding to this family of GM data is isomorphic to $(\mathcal{X} \rightarrow S, \mathcal{H})$. □

The following lemma was used in the proof of Theorem 3.7.

Lemma 3.8. *Let $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ be a family of normalized GM data over a scheme S . Consider the subscheme $\mathcal{X} \subset \mathbf{P}_S(\mathcal{W})$ defined as the zero locus of the global section (20) and assume that for every geometric point s of S , the fiber \mathcal{X}_s of \mathcal{X} is a smooth GM variety of dimension n . Then $\pi_{\mathcal{X}} : \mathcal{X} \rightarrow S$ is a (flat) family of smooth GM varieties.*

Proof. We only have to check that the morphism $\pi_{\mathcal{X}} : \mathcal{X} \rightarrow S$ is flat. Locally over S , the rational linear projection $\mu : \mathbf{P}_S(\mathcal{W}) \dashrightarrow \mathbf{P}_S(\wedge^2 \mathcal{V}_5)$ can be lifted to a linear closed embedding $\mathbf{P}_S(\mathcal{W}) \rightarrow \mathbf{P}_S(\mathcal{K} \oplus \wedge^2 \mathcal{V}_5)$, where \mathcal{K} is a vector bundle over S . Consider the subscheme $M_{\mathcal{X}} \subset \mathbf{P}_S(\mathcal{W})$ defined as the zero locus of

$$\mathbf{q}|_{\mathcal{V}_5} \in H^0(\mathbf{P}_S(\mathcal{W}), \pi_{\mathbf{P}_S(\mathcal{W})}^*(\mathcal{V}_5^\vee \otimes \det(\mathcal{V}_5) \otimes (\mathcal{V}_6/\mathcal{V}_5)^{\otimes 2}) \otimes \mathcal{O}_{\mathbf{P}_S(\mathcal{W})}(2)).$$

By the commutativity of (17), this subscheme can be represented as

$$M_{\mathcal{X}} = \mathbf{P}_S(\mathcal{W}) \times_{\mathbf{P}_S(\wedge^2 \mathcal{V}_5 \oplus \mathcal{K})} \text{Cone}_{\mathbf{P}_S(\mathcal{K})} \text{Gr}_S(2, \mathcal{V}_5),$$

where $\text{Cone}_{\mathbf{P}_S(\mathcal{K})} \text{Gr}_S(2, \mathcal{V}_5) \subset \mathbf{P}_S(\mathcal{K} \oplus \wedge^2 \mathcal{V}_5)$ is the cone over the relative Grassmannian $\text{Gr}_S(2, \mathcal{V}_5)$ with vertex $\mathbf{P}_S(\mathcal{K})$. Furthermore, on $M_{\mathcal{X}}$, the map \mathbf{q} defines a section

$$\mathbf{q}_{\mathcal{V}_6/\mathcal{V}_5} \in H^0(M_{\mathcal{X}}, \pi_{M_{\mathcal{X}}}^*(\det(\mathcal{V}_5) \otimes (\mathcal{V}_6/\mathcal{V}_5)) \otimes \mathcal{O}_{\mathbf{P}_S(\mathcal{W})}(2)|_{M_{\mathcal{X}}})$$

whose zero locus is the subscheme $\mathcal{X} \subset M_{\mathcal{X}} \subset \mathbf{P}_S(\mathcal{W})$.

Since every fiber of $\mathcal{X} \rightarrow S$ has dimension n , every fiber of $M_{\mathcal{X}}$ has dimension at most $n + 1$. On the other hand, it is the intersection of a codimension-3 subvariety $\text{Cone}_{\mathbf{P}_S(\mathcal{K})} \text{Gr}_S(2, \mathcal{V}_5) \subset \mathbf{P}_S(\wedge^2 \mathcal{V}_5 \oplus \mathcal{K})$ with the linear projective subbundle $\mathbf{P}_S(\mathcal{W})$ of dimension $n + 4$, hence each fiber has dimension at least $n + 1$. Combining these two observations, we see that each fiber of $M_{\mathcal{X}}$ has dimension $n + 1$, hence the intersection (the fiber product) defining $M_{\mathcal{X}}$ is dimensionally transverse.

Let us show that $M_{\mathcal{X}}$ is flat over S . The cone $\text{Cone}_{\mathbf{P}_S(\mathcal{K})} \text{Gr}_S(2, \mathcal{V}_5)$ is flat over S and $M_{\mathcal{X}}$ is cut in it by relative hyperplane sections. Since the intersection is dimensionally transverse, each of these hyperplanes decreases the dimension of fibers by 1, hence they form a regular sequence at every fiber. This implies flatness of $M_{\mathcal{X}}$ over S .

Finally, as observed above, \mathcal{X} is the zero locus of a global section of a line bundle on $M_{\mathcal{X}}$ and each fiber of \mathcal{X} has codimension 1 in the corresponding fiber of $M_{\mathcal{X}}$. Therefore, this global section is not a zero divisor at every fiber, hence \mathcal{X} is also flat over S . □

We can restate Theorem 3.7 as follows.

Corollary 3.9. *For each $n \in \{2, \dots, 6\}$, the stack $\mathfrak{M}_n^{\text{GM}}$ of smooth polarized GM varieties of dimension n is equivalent to the stack $\mathfrak{M}_{n, \text{smooth}}^{\text{GM-data}}$ of smooth normalized GM data of dimension n .*

From now on, we will identify the stacks $\mathfrak{M}_n^{\text{GM}}$ and $\mathfrak{M}_{n, \text{smooth}}^{\text{GM-data}}$ by using the equivalence above. In particular, we will sometimes think of an S -point of the stack $\mathfrak{M}_n^{\text{GM}}$ as a family $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ of smooth normalized GM data over S .

3.4. The ordinary and special substacks

Let $(\mathcal{X} \rightarrow S, \mathcal{H})$ be a family of smooth GM varieties. Consider the corresponding family $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ of GM data. There is a commutative diagram

$$\begin{array}{ccc}
 \mathcal{V}_5 \otimes \det(\mathcal{V}_5)^\vee \otimes ((\mathcal{V}_6/\mathcal{V}_5)^\vee)^{\otimes 2} \otimes \mathcal{W} & & \\
 \mu \downarrow & \searrow \mathbf{q} & \\
 \Lambda^2 \mathcal{V}_5^\vee \otimes (\mathcal{V}_6/\mathcal{V}_5)^\vee & \xrightarrow{\mu^T} & \mathcal{W}^\vee
 \end{array} \tag{23}$$

where the vertical arrow is defined as the composition of the map μ with the canonical morphism $\mathcal{V}_5 \otimes \det(\mathcal{V}_5)^\vee \otimes \Lambda^2 \mathcal{V}_5 \rightarrow \Lambda^2 \mathcal{V}_5^\vee$. The commutativity of (23) follows from the commutativity of (17).

Lemma 3.10. *The cokernels of the diagonal maps in (23) are isomorphic. They are line bundles on a closed subscheme of S .*

Proof. Let us show that the arrow μ in (23) is surjective. By Lemma 2.1, this can be done pointwise, so it is enough to consider the case where S is the spectrum of an algebraically closed field \mathbf{K} . Then, $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ is just a normalized GM data set $(W, V_6, V_5, \mu, \mathbf{q})$ over \mathbf{K} .

Assume that the map μ is not surjective. Trivializing $\det(V_5)$ and V_6/V_5 for simplicity, we can rewrite the nonsurjectivity condition as follows: there is an element $\xi \in \Lambda^2 V_5$ such that the subspace $W_0 = \text{Im}(\mu) \subset \Lambda^2 V_5$ is orthogonal to the space $V_5 \wedge \xi \subset \Lambda^3 V_5$, that is,

$$V_5 \wedge \xi \subset W_0^\perp.$$

The space $\mathbf{P}(V_5 \wedge \xi)$ contains quite a lot of decomposable vectors — if ξ is decomposable, $\mathbf{P}(V_5 \wedge \xi) \simeq \mathbf{P}^2$ consists of decomposable vectors only, while if ξ has rank 4, $\mathbf{P}(V_5 \wedge \xi) \simeq \mathbf{P}^4$ contains a \mathbf{P}^3 of decomposable vectors. But by [6, Proposition 3.13], the space W_0^\perp is equal to $A \cap \Lambda^3 V_5$, where $A \subset \Lambda^3 V_6$ is the Lagrangian subspace associated with X by [6, Theorem 3.6]. By [6, Theorem 3.16], for smooth GM varieties of dimension $n \geq 3$, it contains no decomposable vectors, and for smooth GM surfaces, the set of decomposable vectors that it contains has dimension at most 1 [6, Remark 3.17]. The arrow μ in (23) is therefore surjective.

The isomorphism between the cokernels of the horizontal arrows then follows by abstract nonsense. Finally, the rank of the cokernel sheaves is at most 1 by [6, Proposition 2.28]. Therefore, they are line bundles on a subscheme of S by Lemma 2.5. □

Definition 3.11. Given a GM data set over a scheme S , consider the cokernel sheaf

$$\mathcal{E} := \text{Coker}(\mu^T) \simeq \text{Coker}(\mathbf{q}) \tag{24}$$

discussed in Lemma 3.10 and denote by $S_{\text{GM,spe}} \subset S$ its (closed) scheme-theoretic support (we call it the GM-special locus) and by $\mathcal{I}_{\text{GM,spe}} \subset \mathcal{O}_S$ its ideal.

By Lemma 2.5, the scheme $S_{\text{GM,spe}}$ is the degeneration scheme for both morphisms μ^T and \mathbf{q} in (23). We further define

$$S_{\text{GM,ord}} := S \setminus S_{\text{GM,spe}}$$

to be the open complement of $S_{\text{GM,spe}}$ in S (we call it the GM-ordinary locus).

Lemma 3.12. *For each $n \in \{2, \dots, 6\}$, there is an open substack $\mathfrak{M}_{n,\text{ord}}^{\text{GM}} \subset \mathfrak{M}_n^{\text{GM}}$ and a closed substack $\mathfrak{M}_{n,\text{spe}}^{\text{GM}} \subset \mathfrak{M}_n^{\text{GM}}$ such that*

$$\begin{aligned} \mathfrak{M}_{n,\text{ord}}^{\text{GM}}(S) &= \{(\mathcal{X} \rightarrow S, \mathcal{H}) \in \mathfrak{M}_n^{\text{GM}}(S) \mid S_{\text{spe}} = \emptyset\}, \\ \mathfrak{M}_{n,\text{spe}}^{\text{GM}}(S) &= \{(\mathcal{X} \rightarrow S, \mathcal{H}) \in \mathfrak{M}_n^{\text{GM}}(S) \mid S_{\text{spe}} = S\}. \end{aligned}$$

Moreover, $\mathfrak{M}_{n,\text{ord}}^{\text{GM}}$ is the open complement of the closed substack $\mathfrak{M}_{n,\text{spe}}^{\text{GM}} \subset \mathfrak{M}_n^{\text{GM}}$.

Proof. It is enough to prove that the formation of the GM-ordinary $S_{\text{GM,ord}} \subset S$ and GM-special $S_{\text{GM,spe}} \subset S$ loci is functorial in S , that is, that it is compatible with base change. This follows from the fact that the formation of the cokernel sheaf commutes with base change (since the pullback functor is right exact). \square

By [6, Sec. 2.5], the open substack $\mathfrak{M}_{n,\text{ord}}^{\text{GM}} \subset \mathfrak{M}_n^{\text{GM}}$ classifies families of smooth ordinary GM varieties of dimension n , while the closed substack $\mathfrak{M}_{n,\text{spe}}^{\text{GM}} \subset \mathfrak{M}_n^{\text{GM}}$ classifies families of smooth special GM varieties of dimension n .

In the case $n = 2$, consider also the open substack

$$\mathfrak{M}_{2,\text{ord,ss}}^{\text{GM}} \subset \mathfrak{M}_{2,\text{ord}}^{\text{GM}} \tag{25}$$

that classifies strongly smooth ordinary GM surfaces [6, Definition 3.15].

Lemma 3.13. *The stacks $\mathfrak{M}_{n,\text{ord}}^{\text{GM}}$ for $n \in \{2, \dots, 5\}$ and the stacks $\mathfrak{M}_{n,\text{spe}}^{\text{GM}}$ for $n \in \{3, \dots, 6\}$ are smooth Deligne–Mumford stacks of finite type over \mathbf{Q} with affine diagonals of finite type. Moreover,*

$$\dim(\mathfrak{M}_{n,\text{ord}}^{\text{GM}}) = 25 - (5 - n)(6 - n)/2, \quad \dim(\mathfrak{M}_{n,\text{spe}}^{\text{GM}}) = 25 - (6 - n)(7 - n)/2.$$

In particular, for $n \in \{3, \dots, 6\}$, the stack $\mathfrak{M}_{n,\text{spe}}^{\text{GM}}$ has codimension $6 - n$ in $\mathfrak{M}_n^{\text{GM}}$.

For $n \geq 4$, the stack $\mathfrak{M}_{n,\text{spe}}^{\text{GM}}$ is a μ_2 -gerbe over the stack $\mathfrak{M}_{n-1,\text{ord}}^{\text{GM}}$, and the stack $\mathfrak{M}_{3,\text{spe}}^{\text{GM}}$ is a μ_2 -gerbe over the stack $\mathfrak{M}_{2,\text{ord,ss}}^{\text{GM}}$.

Proof. Since $\mathfrak{M}_{n,\text{ord}}^{\text{GM}}$ is an open substack in $\mathfrak{M}_n^{\text{GM}}$, it inherits the properties of the latter established in Proposition 3.2; in particular, it has the same dimension.

We show next that for $n \geq 4$, the stack $\mathfrak{M}_{n,\text{spe}}^{\text{GM}}$ is a μ_2 -gerbe over the stack $\mathfrak{M}_{n-1,\text{ord}}^{\text{GM}}$. Consider the \mathbb{G}_m -gerbes

$$\widetilde{\mathfrak{M}}_{n,\text{spe}}^{\text{GM-data}} \rightarrow \mathfrak{M}_{n,\text{spe}}^{\text{GM}}, \quad \widetilde{\mathfrak{M}}_{n-1,\text{ord}}^{\text{GM-data}} \rightarrow \mathfrak{M}_{n-1,\text{ord}}^{\text{GM}}$$

obtained by passing to linearized data (see Remark 3.4), so that the arrows are rigidification functors with respect to the natural \mathbb{G}_m -actions. We will show that $\widetilde{\mathfrak{M}}_{n,\text{spe}}^{\text{GM-data}}$ is a μ_2 -gerbe over $\widetilde{\mathfrak{M}}_{n-1,\text{ord}}^{\text{GM-data}}$ and that the corresponding μ_2 - and \mathbb{G}_m -actions on objects of $\widetilde{\mathfrak{M}}_{n,\text{spe}}^{\text{GM-data}}$ commute. This will prove that, after passing to \mathbb{G}_m -rigidifications, there is a morphism of stacks $\mathfrak{M}_{n,\text{spe}}^{\text{GM}} \rightarrow \mathfrak{M}_{n-1,\text{ord}}^{\text{GM}}$ which is a μ_2 -gerbe.

To be more precise, we will show that $\widetilde{\mathfrak{M}}_{n,\text{spe}}^{\text{GM-data}}$ is the root stack over $\widetilde{\mathfrak{M}}_{n-1,\text{ord}}^{\text{GM-data}}$ associated with its line bundle $\det(\mathcal{V}_6)$ in the sense of [2, Sec. B.1] (in fact, this is the reason why we pass to stacks of linearized data, since the line bundle $\det(\mathcal{V}_6)$ is just not defined on the stack $\mathfrak{M}_{n-1,\text{ord}}^{\text{GM-data}}$). For this, we check that the groupoid of linearized families $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ of special GM data of dimension n over a scheme S is equivalent to the groupoid of linearized families $(S, \mathcal{W}_0, \mathcal{V}_6, \mathcal{V}_5, \mu_0, \mathbf{q}_0)$ of ordinary GM data of dimension $n - 1$ equipped with a line bundle \mathcal{W}_1 and an isomorphism

$$\mathbf{q}_1 : \text{Sym}^2 \mathcal{W}_1 \xrightarrow{\sim} \det(\mathcal{V}_6).$$

Indeed, given a family of special linearized GM data over a scheme S , we set

$$\mathcal{W}_1 := \text{Ker}(\mathcal{W} \xrightarrow{\mu} \wedge^2 \mathcal{V}_5 \otimes (\mathcal{V}_6/\mathcal{V}_5)).$$

This is a line bundle because, by definition of special GM data and Lemma 3.10, the map μ has constant rank $n + 4$, while \mathcal{W} is a vector bundle of rank $n + 5$. Furthermore, we consider the composition

$$\mathcal{V}_6 \otimes \text{Sym}^2 \mathcal{W}_1 \hookrightarrow \mathcal{V}_6 \otimes \text{Sym}^2 \mathcal{W} \xrightarrow{\mathbf{q}} \det(\mathcal{V}_5) \otimes (\mathcal{V}_6/\mathcal{V}_5)^{\otimes 2} \xrightarrow{\sim} \det(\mathcal{V}_6) \otimes (\mathcal{V}_6/\mathcal{V}_5).$$

By (18), this map vanishes on the subbundle $\mathcal{V}_5 \otimes \text{Sym}^2 \mathcal{W}_1 \subset \mathcal{V}_6 \otimes \text{Sym}^2 \mathcal{W}_1$ hence factors through a morphism $(\mathcal{V}_6/\mathcal{V}_5) \otimes \text{Sym}^2 \mathcal{W}_1 \rightarrow \det(\mathcal{V}_6) \otimes (\mathcal{V}_6/\mathcal{V}_5)$. Twisting it by $(\mathcal{V}_6/\mathcal{V}_5)^\vee$, we obtain a morphism

$$\text{Sym}^2 \mathcal{W}_1 \longrightarrow \det(\mathcal{V}_6)$$

which we denote by \mathbf{q}_1 . It is surjective by Lemma 2.1 and [6, Lemma 2.33], hence an isomorphism by Lemma 2.3, since its source and target are both line bundles. Finally, by [6, Proposition 2.30], there is a canonical direct sum decomposition $\mathcal{W} \simeq \mathcal{W}_0 \oplus \mathcal{W}_1$ (orthogonal with respect to all quadrics). We denote by μ_0 and \mathbf{q}_0 the restrictions of μ to \mathcal{W}_0 and of \mathbf{q} to $\mathcal{V}_6 \otimes \text{Sym}^2 \mathcal{W}_0$. By [6, Lemma 2.33], $(S, \mathcal{W}_0, \mathcal{V}_6, \mathcal{V}_5, \mu_0, \mathbf{q}_0)$ is a linearized family of ordinary GM data of dimension $n - 1$. This defines a functor between the groupoids (the action of the functor on morphisms is obvious).

Conversely, assume we are given a family $(S, \mathcal{W}_0, \mathcal{V}_6, \mathcal{V}_5, \mu_0, \mathbf{q}_0)$ of ordinary GM data of dimension $n - 1$, a line bundle \mathcal{W}_1 , and an isomorphism of line bundles $\mathbf{q}_1 : \text{Sym}^2 \mathcal{W}_1 \xrightarrow{\sim} \det(\mathcal{V}_6)$. We set

$$\mathcal{W} := \mathcal{W}_0 \oplus \mathcal{W}_1, \quad \mu = (\mu_0, 0), \quad \mathbf{q} = \mathbf{q}_0 \oplus \mathbf{q}_1,$$

where by an abuse of notation, we still denote by \mathbf{q}_1 the composition

$$\begin{aligned} \mathcal{V}_6 \otimes \mathrm{Sym}^2 \mathcal{W}_1 &\rightarrow (\mathcal{V}_6/\mathcal{V}_5) \otimes \mathrm{Sym}^2 \mathcal{W}_1 \xrightarrow{\mathbf{q}_1} (\mathcal{V}_6/\mathcal{V}_5) \otimes \det(\mathcal{V}_6) \\ &\xrightarrow{\sim} \det(\mathcal{V}_5) \otimes (\mathcal{V}_6/\mathcal{V}_5)^{\otimes 2}. \end{aligned}$$

By [6, Lemma 2.33], this is a family of smooth special GM data of dimension n .

It is straightforward to see that the constructed functors are mutually inverse, so the groupoids are equivalent, hence $\widetilde{\mathfrak{M}}_{n,\mathrm{spe}}^{\mathrm{GM-data}}$ is the root stack over $\widetilde{\mathfrak{M}}_{n-1,\mathrm{ord}}^{\mathrm{GM-data}}$ and $\mathfrak{M}_{n,\mathrm{spe}}^{\mathrm{GM}}$ is a μ_2 -gerbe over $\mathfrak{M}_{n-1,\mathrm{ord}}^{\mathrm{GM}}$.

For $n = 3$, the argument is the same; the only difference is that the ordinary GM surface associated with a smooth special GM threefold is automatically strongly smooth (see [6, Sec. 3.4]).

This implies that $\mathfrak{M}_{n,\mathrm{spe}}^{\mathrm{GM}}$ is a smooth Deligne–Mumford stack and gives its dimension. The other properties of $\mathfrak{M}_{n,\mathrm{spe}}^{\mathrm{GM}}$ follow from Proposition 3.2, since it is a closed substack in $\mathfrak{M}_n^{\mathrm{GM}}$. The statement about the codimension is a simple computation. \square

Remark 3.14. The proof of Lemma 3.13 shows that the automorphism group scheme of each object of the stack $\mathfrak{M}_{n,\mathrm{spe}}^{\mathrm{GM}}$ contains the constant group scheme μ_2 and that the morphism of stacks $\mathfrak{M}_{n,\mathrm{spe}}^{\mathrm{GM}} \rightarrow \mathfrak{M}_{n-1,\mathrm{ord}}^{\mathrm{GM}}$ is the μ_2 -rigidification.

3.5. Lagrangian data

In [6, Sec. 3], we explained the relation between GM and Lagrangian data sets. We now define families of Lagrangian data and show that they form a stack.

Definition 3.15. A family of Lagrangian data over a scheme S is a quadruple $(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$, where \mathcal{V}_6 is a vector bundle of rank 6 on S , $\mathcal{V}_5 \subset \mathcal{V}_6$ is a subbundle of corank 1, and $\mathcal{A} \subset \wedge^3 \mathcal{V}_6$ is a Lagrangian subbundle.

A morphism between families of Lagrangian data from $(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$ to $(S', \mathcal{V}'_6, \mathcal{V}'_5, \mathcal{A}')$ is a pair (f, φ) fitting into a Cartesian square

$$\begin{array}{ccc} \mathbf{P}_S(\mathcal{V}_6) & \xrightarrow{\varphi} & \mathbf{P}_{S'}(\mathcal{V}'_6) \\ \downarrow & & \downarrow \\ S & \xrightarrow{f} & S' \end{array}$$

and such that $\varphi(\mathbf{P}_S(\mathcal{V}_5)) = \mathbf{P}_{S'}(\mathcal{V}'_5)$ and $(\wedge^3 \varphi)(\mathbf{P}_S(\mathcal{A})) = \mathbf{P}_{S'}(\mathcal{A}')$.

Families of Lagrangian data form a category fibered in groupoids over the category Sch/\mathbf{k} , which we denote $\mathfrak{M}^{\mathrm{Lag}}$.

Remark 3.16. As for GM data (see Remark 3.4), we can define a category $\widetilde{\mathfrak{M}}^{\mathrm{Lag}}$ of families of linearized Lagrangian data (fibered in groupoids over Sch/\mathbf{k}) with the same objects as in $\mathfrak{M}^{\mathrm{Lag}}$ but with morphisms defined as pairs $(f, \widetilde{\varphi})$ formed by a

morphism $f : S \rightarrow S'$ and an isomorphism $\tilde{\varphi} : \mathcal{V}_6 \xrightarrow{\sim} f^*\mathcal{V}'_6$ such that $\tilde{\varphi}(\mathcal{V}_5) = f^*\mathcal{V}'_5$ and $(\wedge^3 \tilde{\varphi})(\mathcal{A}) = \widetilde{f^*\mathcal{A}'}$.

Denoting by $\text{Aut}(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$ the automorphism group scheme in $\widetilde{\mathfrak{M}}^{\text{Lag}}$, we obtain an embedding of group schemes

$$\mathbb{G}_m(S) \hookrightarrow \widetilde{\text{Aut}}(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A}) \tag{26}$$

that takes an invertible function u to the automorphism given by $\tilde{\varphi} = u$. We have

$$\text{Aut}(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A}) \simeq \widetilde{\text{Aut}}(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A}) / \mathbb{G}_m(S)$$

and the fibered category of Lagrangian data is the rigidification of the fibered category of linearized Lagrangian data with respect to the embeddings (26).

This observation implies that any morphism in $\mathfrak{M}^{\text{Lag}}$ over $f : S \rightarrow S'$ can be locally over S' lifted to a morphism in $\widetilde{\mathfrak{M}}^{\text{Lag}}$ (and such a lifting is unique up to the composition with the action of $\mathbb{G}_m(S)$). In what follows, we will frequently use such a lifting.

The argument of the proof of Lemma 3.5 implies the following.

Lemma 3.17. *The fibered categories $\mathfrak{M}^{\text{Lag}}$ and $\widetilde{\mathfrak{M}}^{\text{Lag}}$ are stacks over Sch/k .*

Given a family of Lagrangian data $(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$, we consider the natural epimorphism

$$\lambda : \mathcal{V}_6 \rightarrow \mathcal{V}_6 / \mathcal{V}_5.$$

For each $p \in \{2, \dots, 6\}$, it extends by the Leibniz rule to an epimorphism

$$\lambda_p : \wedge^p \mathcal{V}_6 \rightarrow \wedge^{p-1} \mathcal{V}_5 \otimes (\mathcal{V}_6 / \mathcal{V}_5)$$

whose kernel is the subbundle $\wedge^p \mathcal{V}_5 \subset \wedge^p \mathcal{V}_6$.

Definition 3.18. We say that a family of Lagrangian data $(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$ has rank k if the composition

$$\varphi : \mathcal{A} \hookrightarrow \wedge^3 \mathcal{V}_6 \xrightarrow{\lambda_3} \wedge^2 \mathcal{V}_5 \otimes (\mathcal{V}_6 / \mathcal{V}_5) \tag{27}$$

has rank at most k and $\wedge^{k-1} \varphi_s$ does not vanish for any geometric point s in S . We say that the family of Lagrangian data avoids decomposable vectors if, for each geometric point s of S , the Lagrangian subspace $\mathcal{A}_s \subset \wedge^3 \mathcal{V}_{6,s}$ contains no decomposable vectors, that is, $\mathbf{P}(\mathcal{A}_s) \cap \text{Gr}(3, \mathcal{V}_{6,s}) = \emptyset$.

The above two conditions define a locally closed substack in $\mathfrak{M}^{\text{Lag}}$ classifying families of Lagrangian data of rank k avoiding decomposable vectors. We denote it by $\mathfrak{M}_k^{\text{Lag}}$. We will show later (Proposition 5.7) that this stack is a global quotient stack.

Finally, we define the special and ordinary loci for Lagrangian data.

Definition 3.19. Given a family $(\mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$ of Lagrangian data on a scheme S , of rank k , we denote by

$$S_{\text{Lag, spe}} \subset S$$

the degeneracy locus of the map φ in (27) and by $\mathcal{I}_{\text{Lag,spe}} \subset \mathcal{O}_S$ its ideal (it is generated by the $k \times k$ -minors of φ). We call $S_{\text{Lag,spe}}$ the Lagrangian special locus of S . Its complement

$$S_{\text{Lag,ord}} = S \setminus S_{\text{Lag,spe}}$$

is called the Lagrangian ordinary locus.

As in the proof of Lemma 3.12, the above construction gives rise to a closed substack $\mathfrak{M}_{k,\text{spe}}^{\text{Lag}} \subset \mathfrak{M}_k^{\text{Lag}}$ of special Lagrangian data and an open substack $\mathfrak{M}_{k,\text{ord}}^{\text{Lag}} \subset \mathfrak{M}_k^{\text{Lag}}$ of ordinary Lagrangian data, such that $\mathfrak{M}_{k,\text{ord}}^{\text{Lag}}$ is the open complement of $\mathfrak{M}_{k,\text{spe}}^{\text{Lag}} \subset \mathfrak{M}_k^{\text{Lag}}$.

4. Relation Between Families of GM and Lagrangian Data

We consider below two constructions relating GM data to Lagrangian data. We pay special attention to the relation between their special loci.

4.1. From families of GM data to families of Lagrangian data

Let $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ be a family of smooth normalized GM data. We construct over the same scheme S an associated family of Lagrangian data avoiding decomposable vectors (Definition 3.18).

Our construction is a relative (and normalized) version of the construction of the proof of [6, Theorem 3.6] (with “the odd part” omitted). We consider the diagram

$$\begin{CD} \mathcal{V}_5 \otimes \mathcal{W} \otimes (\mathcal{V}_6/\mathcal{V}_5)^\vee @>f_1>> \bigwedge^3 \mathcal{V}_5 \oplus (\mathcal{V}_6 \otimes \mathcal{W} \otimes (\mathcal{V}_6/\mathcal{V}_5)^\vee) @>f_2>> \mathcal{W}^\vee \otimes \det(\mathcal{V}_6) \\ @. @VVf_3V @. \\ @. @. \bigwedge^3 \mathcal{V}_6 \end{CD} \tag{28}$$

with morphisms defined by

$$\begin{aligned} f_1(v \otimes w) &= (-v \wedge \mu(w), v \otimes w), \\ f_2(\xi, v \otimes w)(w') &= \xi \wedge \mu(w') + \mathbf{q}(v)(w, w'), \\ f_3(\xi, v \otimes w) &= \xi + v \wedge \mu(w) \end{aligned}$$

(we omit factors corresponding to line bundles, which do not matter here). We have $f_2 \circ f_1 = 0$ by (18) and $f_3 \circ f_1 = 0$. If we set

$$\mathcal{A} := \text{Ker}(f_2)/\text{Im}(f_1), \tag{29}$$

the morphism f_3 induces a morphism $\mathcal{A} \rightarrow \bigwedge^3 \mathcal{V}_6$.

Proposition 4.1. *Let $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ be a family of smooth normalized GM data of dimension $n \in \{3, 4, 5, 6\}$. Define \mathcal{A} by (29). Then $(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$ is a*

family of Lagrangian data of rank $n + 5$ avoiding decomposable vectors. This defines a morphism of stacks

$$\mathbf{a} : \mathfrak{M}_n^{\text{GM}} \rightarrow \mathfrak{M}_{n+5}^{\text{Lag}}$$

$$(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q}) \mapsto (S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A}).$$

Moreover, the Lagrangian and GM special loci in S coincide set-theoretically but not necessarily scheme-theoretically: we have

$$\mathcal{I}_{\text{Lag, spe}} = \mathcal{I}_{\text{GM, spe}}^2,$$

that is, the ideal of the Lagrangian special locus is the square of the ideal of the GM special locus.

Proof. Checking that \mathcal{A} is a vector bundle of rank 10 (it is enough for that to check that f_2 is an epimorphism and that f_1 is a fiberwise monomorphism) and that the map $\mathcal{A} \rightarrow \wedge^3 \mathcal{V}_6$ induced by f_3 is a fiberwise monomorphism can be done pointwise and thus follows from the proof of [6, Theorem 3.6].

We now show that \mathcal{A} has the Lagrangian property, that is, that the composition

$$\mathcal{A} \otimes \mathcal{A} \longrightarrow \wedge^3 \mathcal{V}_6 \otimes \wedge^3 \mathcal{V}_6 \xrightarrow{\wedge} \det(\mathcal{V}_6) \tag{30}$$

vanishes identically. It is not enough to prove this property pointwise, since the scheme S might be nonreduced, but it is enough to check it locally. It will be convenient to compose (30) with the isomorphism $\lambda_6 : \det(\mathcal{V}_6) \xrightarrow{\sim} \det(\mathcal{V}_5) \otimes (\mathcal{V}_6/\mathcal{V}_5)$. We will also use the definition (29) of \mathcal{A} and the fact that the map $\mathcal{A} \rightarrow \wedge^3 \mathcal{V}_6$ is induced by f_3 . The resulting composition

$$\begin{aligned} & (\wedge^3 \mathcal{V}_5 \oplus (\mathcal{V}_6 \otimes \mathcal{W} \otimes (\mathcal{V}_6/\mathcal{V}_5)^\vee)) \otimes (\wedge^3 \mathcal{V}_5 \oplus (\mathcal{V}_6 \otimes \mathcal{W} \otimes (\mathcal{V}_6/\mathcal{V}_5)^\vee)) \\ & \xrightarrow{f_3 \wedge f_3} \det(\mathcal{V}_6) \xrightarrow{\lambda_6} \det(\mathcal{V}_5) \otimes (\mathcal{V}_6/\mathcal{V}_5) \end{aligned}$$

is given by

$$(\xi_1, v_1 \otimes w_1) \otimes (\xi_2, v_2 \otimes w_2) \mapsto \lambda_6((\xi_1 + v_1 \wedge \mu(w_1)) \wedge (\xi_2 + v_2 \wedge \mu(w_2))). \tag{31}$$

It is thus enough to check that (31) vanishes on $\text{Ker}(f_2) \otimes \text{Ker}(f_2)$.

Since ξ_1 and ξ_2 are sections of $\wedge^3 \mathcal{V}_5$, we have $\xi_1 \wedge \xi_2 = 0$. Choosing locally a direct sum decomposition $\mathcal{V}_6 = \mathcal{V}_5 \oplus (\mathcal{V}_6/\mathcal{V}_5)$ and a generator v_0 for the second summand, we can write

$$v_i = v'_i + \lambda(v_i)v_0 \quad \text{with } v'_i \in \mathcal{V}_5,$$

where we think of $\lambda(v_i)$ as of a scalar. We can rewrite the right side of (31) as

$$\begin{aligned} & \lambda_6(\xi_1 \wedge v_2 \wedge \mu(w_2)) + \lambda_6(v_1 \wedge \mu(w_1) \wedge \xi_2) + \lambda_6(v_1 \wedge \mu(w_1) \wedge v_2 \wedge \mu(w_2)) \\ & = -\lambda(v_2)\xi_1 \wedge \mu(w_2) + \lambda(v_1)\xi_2 \wedge \mu(w_1) \\ & \quad + \lambda(v_1)v'_2 \wedge \mu(w_1) \wedge \mu(w_2) - \lambda(v_2)v'_1 \wedge \mu(w_1) \wedge \mu(w_2) \end{aligned}$$

$$\begin{aligned}
 &= -\lambda(v_2)(\xi_1 \wedge \mu(w_2) + \mathbf{q}(v'_1)(w_1, w_2)) + \lambda(v_1)(\xi_2 \wedge \mu(w_1) + \mathbf{q}(v'_2)(w_1, w_2)) \\
 &= -\lambda(v_2)(\xi_1 \wedge \mu(w_2) + \mathbf{q}(v_1)(w_1, w_2)) + \lambda(v_2)\lambda(v_1)\mathbf{q}(v_0)(w_1, w_2) \\
 &\quad + \lambda(v_1)(\xi_2 \wedge \mu(w_1) + \mathbf{q}(v_2)(w_1, w_2)) - \lambda(v_1)\lambda(v_2)\mathbf{q}(v_0)(w_1, w_2) \\
 &= -\lambda(v_2)f_2(\xi_1, v_1 \otimes w_1)(w_2) + \lambda(v_1)f_2(\xi_2, v_2 \otimes w_2)(w_1)
 \end{aligned}$$

(in the first equality, we use the Leibniz rule for λ_6 and the fact that λ vanishes on ξ_i and on $\mu(w_i)$, as well as the relation

$$\lambda_2(v_1 \wedge v_2) = \lambda(v_1)v_2 - \lambda(v_2)v_1 = \lambda(v_1)v'_2 - \lambda(v_2)v'_1;$$

in the second equality, we use (18); in the third equality, we use the definition of v'_i ; and in the last equality, we use the definition of f_2 and cancel out two summands equal to $\pm\lambda(v_1)\lambda(v_2)\mathbf{q}(v_0)(w_1, w_2)$). It follows that the map (31) vanishes identically on the subbundle $\text{Ker}(f_2) \otimes \text{Ker}(f_2)$, hence the induced map vanishes identically on \mathcal{A} .

Consider now the map φ defined by (27). It is induced by the composition of the maps in the top row and the right column of the diagram

$$\begin{array}{ccccc}
 \text{Ker}(f_2) \hookrightarrow & \Lambda^3 \mathcal{V}_5 \oplus (\mathcal{V}_6 \otimes \mathcal{W} \otimes (\mathcal{V}_6/\mathcal{V}_5)^\vee) & \xrightarrow{f_3} & \Lambda^3 \mathcal{V}_6 & \\
 \downarrow & \downarrow (0, \lambda) & & \downarrow \lambda_3 & \\
 \mathcal{A} & \xrightarrow{\nu} & \mathcal{W} & \xrightarrow{\mu} & \Lambda^2 \mathcal{V}_5 \otimes (\mathcal{V}_6/\mathcal{V}_5).
 \end{array} \tag{32}$$

The right square of the diagram is commutative because λ_3 vanishes on $\Lambda^3 \mathcal{V}_5$ and λ_2 vanishes on $\Lambda^2 \mathcal{V}_5$. The arrow $(0, \lambda)$ vanishes on $\text{Im}(f_1) \subset \text{Ker}(f_2)$ hence factors through \mathcal{A} , thus defining the arrow ν . Therefore, we obtain a commutative diagram

$$\begin{array}{ccc}
 \mathcal{A} \hookrightarrow & \Lambda^3 \mathcal{V}_6 & \\
 \downarrow \nu & \searrow \varphi & \downarrow \lambda_3 \\
 \mathcal{W} & \xrightarrow{\mu} & \Lambda^2 \mathcal{V}_5 \otimes (\mathcal{V}_6/\mathcal{V}_5).
 \end{array} \tag{33}$$

The rank of \mathcal{W} is $n + 5$, hence the rank of φ is at most $n + 5$. The fact that it is at least $n + 4$ at each geometric point of S can be verified pointwise and follows from [6, (3.9)]. Also, [6, (3.9), Theorem 3.16] proves that \mathcal{A} has no decomposable vectors (this is the only place where we use the condition $n \geq 3$). Thus, $(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$ is a family of Lagrangian data of rank $n + 5$ avoiding decomposable vectors.

Let us show that the association $\mathbf{a} : \mathfrak{M}_n^{\text{GM}} \rightarrow \mathfrak{M}_{n+5}^{\text{Lag}}$ that takes a family of smooth normalized GM data $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ to the family of Lagrangian data $(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$, where \mathcal{A} is defined by (29), is a morphism of stacks, meaning that it is defined on morphisms.

Assume for simplicity that $f = \text{id}_S$ (the general case reduces to this by base change). A morphism of families of GM data from $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ to $(S, \mathcal{W}', \mathcal{V}'_6, \mathcal{V}'_5, \mu', \mathbf{q}')$ is then given by a pair of isomorphisms of projective bundles $\varphi_V : \mathbf{P}_S(\mathcal{V}_6) \xrightarrow{\sim} \mathbf{P}_S(\mathcal{V}'_6)$ and $\varphi_W : \mathbf{P}_S(\mathcal{W}) \xrightarrow{\sim} \mathbf{P}_S(\mathcal{W}')$ over S . The first can be lifted to an isomorphism

$$\tilde{\varphi}_V : \mathcal{V}_6 \xrightarrow{\sim} \mathcal{V}'_6 \otimes \delta$$

for an appropriate line bundle δ on S . Using compatibility with the morphism μ , we conclude that φ_W lifts to an isomorphism

$$\tilde{\varphi}_W : \mathcal{W} \xrightarrow{\sim} \mathcal{W}' \otimes \delta^{\otimes 3}.$$

It is straightforward to see that the pair $(\tilde{\varphi}_V, \tilde{\varphi}_W)$ defines a morphism from the diagram (28) to the analogous diagram for the family of GM data $(S, \mathcal{W}', \mathcal{V}'_6, \mathcal{V}'_5, \mu', \mathbf{q}')$ twisted by $\delta^{\otimes 3}$. It follows that the morphism $\bigwedge^3 \tilde{\varphi}_V : \bigwedge^3 \mathcal{V}_6 \rightarrow \bigwedge^3 \mathcal{V}'_6 \otimes \delta^{\otimes 9}$ takes \mathcal{A} to $\mathcal{A}' \otimes \delta^{\otimes 3}$. Therefore, we have $\bigwedge^3 \varphi_V(\mathbf{P}_S(\mathcal{A})) = \mathbf{P}_S(\mathcal{A}')$, hence φ_V is a morphism between the associated families of Lagrangian data. This operation is compatible with compositions of morphisms and takes the identity to the identity, hence \mathfrak{a} is a morphism of stacks.

Finally, consider the special locus of the family of Lagrangian data constructed above. Its ideal $\mathcal{I}_{\text{Lag, spe}}$ is generated by the $(n + 5) \times (n + 5)$ -minors of the map φ defined by (27). Because of the factorization in (33) (and since the rank of \mathcal{W} is $n + 5$), every such minor is the product of a minor of ν and a minor of μ of the same size. Consequently, the ideal $\mathcal{I}_{\text{Lag, spe}}$ is the product of two ideals, one generated by the minors of ν and the other generated by the minors of μ . The latter ideal is by definition equal to the ideal $\mathcal{I}_{\text{GM, spe}}$ defining the special GM locus. To finish the proof, we must show that the minors of ν generate the same ideal.

Since the left vertical arrow in (32) is surjective, this ideal coincides with the ideal generated by the minors of the map $\text{Ker}(f_2) \rightarrow \mathcal{W}$, that is, by Lemma 2.5, with the annihilator of the cokernel of this map. Since f_2 is surjective, the cokernel of the map $\text{Ker}(f_2) \xrightarrow{(0, \lambda)} \mathcal{W}$ is isomorphic to the cokernel of the map

$$\bigwedge^3 \mathcal{V}_5 \oplus (\mathcal{V}_6 \otimes \mathcal{W} \otimes (\mathcal{V}_6/\mathcal{V}_5)^\vee) \xrightarrow{(0, \lambda) + f_2} \mathcal{W} \oplus (\mathcal{W}^\vee \otimes \det(\mathcal{V}_6)).$$

Since the map $(0, \lambda)$ is surjective, this sheaf is isomorphic to the cokernel of the map $\text{Ker}(0, \lambda) \xrightarrow{f_2} \mathcal{W}^\vee \otimes \det(\mathcal{V}_6)$. Altogether, this means

$$\text{Coker}(\nu) \simeq \text{Coker}(\bigwedge^3 \mathcal{V}_5 \oplus (\mathcal{V}_5 \otimes \mathcal{W} \otimes (\mathcal{V}_6/\mathcal{V}_5)^\vee) \xrightarrow{(\mu, \mathbf{q})} \mathcal{W}^\vee \otimes \det(\mathcal{V}_6)).$$

By Lemma 3.10, the images of the two components of the map (μ, \mathbf{q}) coincide, hence the cokernel of their sum equals to the cokernel of each of them, that is,

$$\text{Coker}(\nu) \simeq \mathcal{C}, \tag{34}$$

where \mathcal{C} is the cokernel sheaf of the family of GM data as defined in (24). The annihilator of $\text{Coker}(\nu)$ is thus again the ideal $\mathcal{I}_{\text{GM, spe}}$ of the GM special locus. \square

Remark 4.2. The argument of Proposition 4.1 also applies to families of smooth normalized GM data of dimension $n = 2$ such that the corresponding GM varieties are strongly smooth [6, Definition 3.15] ordinary GM surfaces since, by [6, (3.9), Theorem 3.16], the corresponding Lagrangian subspaces have no decomposable vectors. It defines a morphism of stacks $\mathfrak{M}_{2,\text{ord,ss}}^{\text{GM}} \rightarrow \mathfrak{M}_7^{\text{Lag}}$.

We will need some properties of the construction presented above. Consider the family

$$\begin{aligned} \mathbf{q}_{\mathcal{A}} : \mathcal{V}_6 \otimes \text{Sym}^2 \mathcal{A} &\rightarrow \det(\mathcal{V}_5) \otimes (\mathcal{V}_6/\mathcal{V}_5)^{\otimes 2} \\ v \otimes a_1 \otimes a_2 &\mapsto -\lambda_4(v \wedge a_1) \wedge \lambda_3(a_2) \end{aligned} \tag{35}$$

of quadratic forms on \mathcal{A} (the formula is symmetric in a_1 and a_2 by the Lagrangian property of \mathcal{A} ; see the proof of [6, Theorem 3.6] for details).

Lemma 4.3. *The quadratic form on \mathcal{A} defined by (35) is equal to the form induced by \mathbf{q} via the map $\nu : \mathcal{A} \rightarrow \mathcal{W}$.*

Proof. Let v be a local section of \mathcal{V}_6 and let a_1 and a_2 be local sections of \mathcal{A} . Choose a lift of a_i to a local section $(\xi_i, v_i \otimes w_i)$ of

$$\text{Ker}(f_2) \subset \bigwedge^3 \mathcal{V}_5 \oplus (\mathcal{V}_6 \otimes \mathcal{W} \otimes (\mathcal{V}_6/\mathcal{V}_5)^\vee).$$

We have

$$\lambda_4(v \wedge f_3(\xi_1, v_1 \otimes w_1)) = \lambda_4(v \wedge (\xi_1 + v_1 \wedge \mu(w_1))) = \lambda(v)\xi_1 + \lambda_2(v \wedge v_1) \wedge \mu(w_1)$$

and

$$\lambda_3(f_3(\xi_2, v_2 \otimes w_2)) = \lambda_3(\xi_2 + v_2 \wedge \mu(w_2)) = \lambda(v_2)\mu(w_2).$$

Therefore,

$$\begin{aligned} \mathbf{q}_{\mathcal{A}}(v)(a_1, a_2) &= -\lambda_4(v \wedge f_3(\xi_1, v_1 \otimes w_1)) \wedge \lambda_3(f_3(\xi_2, v_2 \otimes w_2)) \\ &= -(\lambda(v)\xi_1 + \lambda_2(v \wedge v_1) \wedge \mu(w_1)) \wedge \lambda(v_2)\mu(w_2) \\ &= -\lambda(v_2)\lambda(v)\xi_1 \wedge \mu(w_2) - \lambda(v_2)\lambda_2(v \wedge v_1) \wedge \mu(w_1) \wedge \mu(w_2). \end{aligned}$$

On the other hand, since $(\xi_1, v_1 \otimes w_1)$ is in the kernel of f_2 , we have the relation $\xi_1 \wedge \mu(w_2) = -\mathbf{q}(v_1)(w_1, w_2)$. Using this and (18), the above equals

$$\begin{aligned} &\lambda(v_2)\lambda(v)\mathbf{q}(v_1)(w_1, w_2) - \lambda(v_2)\lambda_2(v \wedge v_1) \wedge \mu(w_1) \wedge \mu(w_2) \\ &= \mathbf{q}(\lambda(v_2)\lambda(v)v_1 - \lambda(v_2)\lambda_2(v \wedge v_1))(w_1, w_2). \end{aligned}$$

It remains to observe that $\lambda(v_2)\lambda(v)v_1 - \lambda(v_2)\lambda_2(v \wedge v_1) = \lambda(v_1)\lambda(v_2)v$, so that finally

$$\begin{aligned} \mathbf{q}_{\mathcal{A}}(v)(a_1, a_2) &= \lambda(v_1)\lambda(v_2)\mathbf{q}(v)(w_1, w_2) \\ &= \mathbf{q}(v)(\lambda(v_1)w_1, \lambda(v_2)w_2) \\ &= \mathbf{q}(v)(\nu(a_1), \nu(a_2)). \end{aligned}$$

This is precisely the compatibility we were claiming. □

Lemma 4.4. *Assume $n \leq 5$ and that the GM special locus $S_{\text{GM,spe}}$ of a smooth family of normalized GM data is a Cartier divisor. The map (27) of the corresponding family of Lagrangian data then factors as*

$$\begin{array}{ccccc}
 \mathcal{A} & \xrightarrow{\nu} & \mathcal{W} & \xrightarrow{\mu} & \bigwedge^2 \mathcal{V}_5 \otimes (\mathcal{V}_6/\mathcal{V}_5) \\
 & \searrow & \nearrow & \searrow & \nearrow \\
 & & \mathcal{W}' & & \mathcal{W}''
 \end{array}$$

where \mathcal{W}' and \mathcal{W}'' are vector bundles of rank $n + 5$, the left diagonal arrow is an epimorphism, the right diagonal arrow is a fiberwise monomorphism, the inner diagonal arrows are monomorphisms, and their cokernels are line bundles on the subscheme $S_{\text{GM,spe}}$.

Proof. Let \mathcal{W}' be the image of ν (so that the map $\mathcal{A} \rightarrow \mathcal{W}'$ is surjective). We have an exact sequence

$$0 \rightarrow \mathcal{W}' \rightarrow \mathcal{W} \rightarrow \text{Coker}(\nu) \rightarrow 0.$$

By (34) and Lemma 3.10, $\text{Coker}(\nu)$ is a line bundle on $S_{\text{GM,spe}}$. Since $S_{\text{GM,spe}}$ is a Cartier divisor, we conclude that \mathcal{W}' is a vector bundle (Lemma 2.4). Analogously, considering the dual of the map μ , we construct the vector bundle \mathcal{W}'' and the other factorization. □

4.2. From families of Lagrangian data to families of GM data

Starting from this section, we assume $n \leq 5$. Let $(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$ be a family of Lagrangian data of rank $n + 5$ avoiding decomposable vectors. Let $S_{\text{Lag,spe}} \subset S$ be its special locus. Assume additionally that $S_{\text{Lag,spe}}$ is a double Cartier divisor (Definition 2.10), that is,

$$S_{\text{Lag,spe}} = 2E, \tag{36}$$

(equivalently, $\mathcal{I}_{\text{Lag,spe}} = \mathcal{I}_E^2$), where E is an effective Cartier divisor. We will construct from $(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$ a family of smooth normalized GM data on S .

Consider the map (27). By definition, its rank is $n + 5$, it is at least $n + 4$ at every geometric point, and $S_{\text{Lag,spe}}$ is its degeneration scheme. Since $S_{\text{Lag,spe}}$ is a Cartier divisor, Proposition 2.8 applies and implies that the map can be written as a composition

$$\mathcal{A} \twoheadrightarrow \mathcal{W}' \rightarrow \mathcal{W}'' \hookrightarrow \bigwedge^2 \mathcal{V}_5 \otimes (\mathcal{V}_6/\mathcal{V}_5),$$

where \mathcal{W}' and \mathcal{W}'' are vector bundles of rank $n + 5$, the left arrow is an epimorphism, the right arrow is a fiberwise monomorphism, and the middle map is a monomorphism whose cokernel $\mathcal{W}''/\mathcal{W}'$ is a line bundle on $S_{\text{Lag,spe}} = 2E$. Tensoring over \mathcal{O}_{2E} the exact sequence $0 \rightarrow \mathcal{O}_E(-E) \rightarrow \mathcal{O}_{2E} \rightarrow \mathcal{O}_E \rightarrow 0$ with $\mathcal{W}''/\mathcal{W}'$, we obtain

an exact sequence

$$0 \rightarrow (\mathcal{W}''/\mathcal{W}') \otimes_{\mathcal{O}_{2E}} \mathcal{O}_E(-E) \rightarrow \mathcal{W}''/\mathcal{W}' \rightarrow (\mathcal{W}''/\mathcal{W}') \otimes_{\mathcal{O}_{2E}} \mathcal{O}_E \rightarrow 0,$$

where the first and last terms are both line bundles on E . Moreover, this is the unique representation of $\mathcal{W}''/\mathcal{W}'$ as an extension of two line bundles on E .

We define \mathcal{W} as the kernel of the map $\mathcal{W}'' \rightarrow \mathcal{W}''/\mathcal{W}' \rightarrow (\mathcal{W}''/\mathcal{W}') \otimes_{\mathcal{O}_{2E}} \mathcal{O}_E$, so that we have a factorization

$$\varphi : \mathcal{A} \rightarrow \mathcal{W}' \rightarrow \mathcal{W} \rightarrow \mathcal{W}'' \hookrightarrow \wedge^2 \mathcal{V}_5 \otimes (\mathcal{V}_6/\mathcal{V}_5), \tag{37}$$

where \mathcal{W}' , \mathcal{W} , and \mathcal{W}'' are vector bundles of rank $n + 5$, the two middle maps are monomorphisms, and \mathcal{W}/\mathcal{W}' and $\mathcal{W}''/\mathcal{W}$ are line bundles on E . This is the unique factorization of φ with these properties.

We define the map

$$\mu : \mathcal{W} \rightarrow \wedge^2 \mathcal{V}_5 \otimes (\mathcal{V}_6/\mathcal{V}_5)$$

as the composition of the third and the fourth arrows in (37) and the map

$$\nu : \mathcal{A} \rightarrow \mathcal{W}$$

as the composition of the first and the second arrows. With these definitions, we have again a commutative square (33).

Lemma 4.5. *The family of quadratic forms*

$$\mathbf{q}_{\mathcal{A}} : \mathcal{V}_6 \otimes \text{Sym}^2 \mathcal{A} \rightarrow \det(\mathcal{V}_5) \otimes (\mathcal{V}_6/\mathcal{V}_5)^{\otimes 2}$$

defined by (35) induces a family of quadratic forms

$$\mathbf{q} : \mathcal{V}_6 \otimes \text{Sym}^2 \mathcal{W} \rightarrow \det(\mathcal{V}_5) \otimes (\mathcal{V}_6/\mathcal{V}_5)^{\otimes 2}.$$

Proof. We will proceed in two steps. First, we show that $\mathbf{q}_{\mathcal{A}}$ induces a family \mathbf{q}' of quadratic forms on \mathcal{W}' . Since $\mathcal{A} \rightarrow \mathcal{W}'$ is surjective, it is enough to show that its kernel bundle is contained in the kernel of $\mathbf{q}_{\mathcal{A}}$. This is obvious, since the kernel of that map is contained in the kernel of λ_3 (by definition of \mathcal{W}') which in turn is contained in the kernel of $\mathbf{q}_{\mathcal{A}}$ by (35).

We then show that the family of quadratic forms \mathbf{q}' on \mathcal{W}' induces a family of quadratic forms on \mathcal{W} given by the Hecke transform of \mathbf{q}' as defined in Lemma 2.11. We set

$$\mathcal{E} := \mathcal{W}' \quad \mathcal{K} := \mathcal{W}''/\mathcal{W}'(-2E), \quad \mathcal{V} := \mathcal{V}_6 \otimes \det(\mathcal{V}_5)^\vee \otimes (\mathcal{V}_6/\mathcal{V}_5)^\vee.$$

Restricting the sequence $0 \rightarrow \mathcal{W}' \rightarrow \mathcal{W}'' \rightarrow \mathcal{W}''/\mathcal{W}' \rightarrow 0$ to $2E$, we obtain

$$0 \rightarrow \mathcal{K} \rightarrow \mathcal{W}'_{2E} \rightarrow \mathcal{W}''_{2E} \rightarrow \mathcal{W}''/\mathcal{W}' \rightarrow 0,$$

thus the line bundle \mathcal{K} on $2E$ is the kernel of the map $\mathcal{W}'_{2E} \rightarrow \mathcal{W}''_{2E}$. In particular, it is a line subbundle of \mathcal{E}_{2E} . Moreover, \mathcal{K} is contained in the kernel of the map $\mathcal{W}' \rightarrow \wedge^2 \mathcal{V}_5 \otimes (\mathcal{V}_6/\mathcal{V}_5)$ restricted to $2E$, hence, by the definition of $\mathbf{q}_{\mathcal{A}}$ in (35), it is contained in the kernel of \mathbf{q}' .

We are therefore in the set-up of Lemma 2.11. By definition of \mathscr{W} , there is an exact sequence

$$0 \rightarrow \mathscr{W}' \rightarrow \mathscr{W} \rightarrow \mathscr{K}_E(E) \rightarrow 0.$$

Comparing with (8), we see that the bundle $\mathscr{W} \simeq \tilde{\mathscr{E}}$ can be identified with the Hecke transform of \mathscr{W}' with respect to \mathscr{K} , and Lemma 2.11 provides it with a family of quadratic forms. □

We can now prove the main result of this section.

Proposition 4.6. *The collection $(S, \mathscr{W}, \mathscr{V}_6, \mathscr{V}_5, \mu, \mathbf{q})$ constructed above is a family of smooth normalized GM data of dimension n . Its special locus coincides scheme-theoretically with the Cartier divisor E .*

Proof. To show that $(S, \mathscr{W}, \mathscr{V}_6, \mathscr{V}_5, \mu, \mathbf{q})$ is a family of GM data, we only have to verify (18). Since the family of quadratic forms \mathbf{q} is induced by $\mathbf{q}_{\mathscr{A}}$, it is enough to check that $\mathbf{q}_{\mathscr{A}}(v)(a_1, a_2) = v \wedge \mu(\nu(a_1)) \wedge \mu(\nu(a_2))$ for $v \in \mathscr{V}_5$. But this follows from the equality $\lambda_4(v \wedge \xi) = -v \wedge \lambda_3(\xi)$ for $v \in \mathscr{V}_5$ and $\xi \in \bigwedge^3 \mathscr{V}_6$, and the commutativity of (33).

The statement about the special locus is also clear, since by construction, the map μ is a composition $\mathscr{W} \rightarrow \mathscr{W}'' \hookrightarrow \bigwedge^2 \mathscr{V}_5 \otimes (\mathscr{V}_6/\mathscr{V}_5)$, where the second map is a fiberwise monomorphism and the degeneration scheme of the first map is equal to E .

It remains to show that the family of GM data is smooth, that is, that for each geometric point s of S , the GM intersection corresponding to the GM data at the point s is smooth.

If $s \notin E$, then $\mathscr{W}_s = \mathscr{W}'_s$ is the image of the map $\mathscr{A}_s \xrightarrow{(\lambda_3)_s} \bigwedge^2 \mathscr{V}_{5,s} \otimes (\mathscr{V}_6/\mathscr{V}_5)_s$ and the quadratic form \mathbf{q} on it is induced by the form $\mathbf{q}_{\mathscr{A}}$ on \mathscr{A}_s . Therefore, by [6, Theorem 3.6], the corresponding GM intersection is just the ordinary GM variety associated with the Lagrangian subspace $\mathscr{A}_s \subset \bigwedge^3 \mathscr{V}_{6,s}$, which has no decomposable vectors, and the hyperplane $\mathscr{V}_{5,s} \subset \mathscr{V}_{6,s}$. It is smooth by [6, Theorem 3.16].

Now assume $s \in E$. For brevity, we write V_6, V_5, A, W', W, W'' and so on for the fibers of the corresponding vector bundles at the geometric point s . We also choose a trivialization for V_6/V_5 to get rid of it in the formulas. Consider the restriction

$$A \twoheadrightarrow W' \rightarrow W \rightarrow W'' \hookrightarrow \bigwedge^2 V_5 \tag{38}$$

to s of the sequence (37). Denote by K_A, K' , and K the respective kernels of the first three maps in (38). We have $\dim(K_A) = 5 - n$ and $\dim(K') = \dim(K) = 1$. Since the rank of the composition of the maps in (38) is $n + 4$ (because s is a point of the special locus), it follows that K' is equal to the kernel of the composition $W' \rightarrow W \rightarrow \bigwedge^2 V_5$. Therefore, the map $\mu : W \rightarrow \bigwedge^2 V_5$ is injective on $W_0 = \nu(A)$.

In particular, we have a canonical direct sum decomposition

$$W = K \oplus W_0. \tag{39}$$

Note that $(W_0, V_6, V_5, \mu|_{W_0}, \mathbf{q}|_{W_0})$ is a GM data set corresponding to a smooth GM variety X_0 of dimension $n - 1$: this follows from [6, Theorems 3.6 and 3.16].

The direct sum decomposition (39) coincides with the direct sum decomposition of Proposition 2.12(a) (its construction is the same). Therefore, by Proposition 2.12(b), the decomposition is orthogonal with respect to the quadrics $\mathbf{q}(v)$ for all $v \in V_6$. Furthermore, the subspace K is contained in the kernel of the quadric $\mathbf{q}(v)$ for all $v \in V_5$, since (18) holds and K is the kernel of μ . It follows that

$$M_X = \bigcap_{v \in V_5} \{\mathbf{q}(v) = 0\} \subset \mathbf{P}(W)$$

is the cone with vertex $\mathbf{P}(K) \subset \mathbf{P}(W)$ over the Grassmannian hull $M_{X_0} \subset \mathbf{P}(W_0)$ of X_0 (see [6, Sec. 2.4]) and, if $v_0 \in V_6 \setminus V_5$, the last quadric $\mathbf{q}(v_0)$ can be written as a sum $\mathbf{q}(v_0) = \mathbf{q}_1 \oplus \mathbf{q}_0$, where $\mathbf{q}_1 \in \text{Sym}^2 K^\vee$ and $\mathbf{q}_0 \in \text{Sym}^2 W_0^\vee$ is the equation of X_0 in M_{X_0} .

Let us now show that \mathbf{q}_1 , which by Proposition 2.12(b) is the residual quadric of $\mathbf{q}'(v_0)$, is nonzero.

For a general vector $v_0 \in V_6 \setminus V_5$, we have $A \cap (v_0 \wedge \wedge^2 V_6) = 0$ (see Remark 5.3). Consider the family of quadratic forms $\mathbf{q}(v_0)$ in a small neighborhood S^0 of s in S . Upon shrinking S^0 , we may assume that the vector bundle \mathcal{V}_6 is trivial with fiber V_6 , that

$$\wedge^3 V_6 \otimes \mathcal{O} = \wedge^3 \mathcal{V}_5 \oplus (v_0 \wedge \wedge^2 V_6) \otimes \mathcal{O}$$

is a Lagrangian direct sum decomposition, and that $\mathcal{A} \cap (v_0 \wedge \wedge^2 V_6) \otimes \mathcal{O} = 0$ at all points of S^0 . By [6, Lemma C.5], we obtain

$$\text{Coker}(\mathbf{q}_{\mathcal{A}}(v_0) : \mathcal{A} \rightarrow \mathcal{A}^\vee) \simeq \text{Coker}(\varphi^\vee : \wedge^2 \mathcal{V}_5^\vee \rightarrow \mathcal{A}^\vee)$$

(where we assume that the line bundles $\det(\mathcal{V}_5)$ and $\mathcal{V}_6/\mathcal{V}_5$ are trivial on S^0). Since the kernel of the epimorphism $\mathcal{A} \rightarrow \mathcal{W}'$ is contained in the kernels of both $\mathbf{q}_{\mathcal{A}}(v_0)$ and φ , we can cancel it out and obtain

$$\text{Coker}(\mathbf{q}'(v_0) : \mathcal{W}' \rightarrow \mathcal{W}'^\vee) \simeq \text{Coker}(\wedge^2 \mathcal{V}_5^\vee \rightarrow \mathcal{W}'^\vee).$$

The rightmost map factors as an epimorphism $\wedge^2 \mathcal{V}_5^\vee \rightarrow \mathcal{W}''^\vee$ followed by the dual of the map $\mathcal{W}' \rightarrow \mathcal{W}''$. Dualizing the sequence $0 \rightarrow \mathcal{W}' \rightarrow \mathcal{W}'' \rightarrow \mathcal{W}''/\mathcal{W}' \rightarrow 0$ and taking into account that $\mathcal{W}''/\mathcal{W}'$ is a line bundle on the Cartier divisor $2E$ (by construction of the bundles \mathcal{W}' and \mathcal{W}'' at the beginning of Sec. 4.2), we deduce that $\text{Coker}(\wedge^2 \mathcal{V}_5^\vee \rightarrow \mathcal{W}''^\vee)$ is a line bundle on $2E$, hence so is the sheaf $\text{Coker}(\mathbf{q}'(v_0) : \mathcal{W}' \rightarrow \mathcal{W}'^\vee)$. By Lemma 2.5, the discriminant $\text{Disc}(\mathbf{q}'(v_0))$ is equal to $2E$ and by Lemma 2.9, the residual quadric \mathbf{q}_1 is nonzero.

In particular, the GM intersection X corresponding to the point s is the double covering of M_{X_0} branched over X_0 , that is, the special GM variety associated with X_0 (see [6, Lemma 2.33]). It is a smooth GM variety of dimension n . \square

4.3. Compositions of the constructions

We show that the constructions introduced in Secs. 4.1 and 4.2 are mutually inverse. We keep the assumption $n \leq 5$.

Let S be a scheme and let $E \subset S$ be an effective Cartier divisor. Denote by

$$\mathfrak{M}_n^{\text{GM}}(S, E) = \{(S, \mathscr{W}, \mathscr{V}_6, \mathscr{V}_5, \mu, \mathbf{q}) \in \mathfrak{M}_n^{\text{GM}}(S) \mid S_{\text{GM, spe}} = E\},$$

$$\mathfrak{M}_{n+5}^{\text{Lag}}(S, E) = \{(S, \mathscr{V}_6, \mathscr{V}_5, \mathscr{A}) \in \mathfrak{M}_{n+5}^{\text{Lag}}(S) \mid S_{\text{Lag, spe}} = E\}$$

the subgroupoids of $\mathfrak{M}_n^{\text{GM}}(S)$ and $\mathfrak{M}_{n+5}^{\text{Lag}}(S)$ (defined in Secs. 3.1 and 3.5) formed by all families of smooth normalized GM data of dimension n (respectively, by all families of Lagrangian data of rank $n + 5$ avoiding decomposable vectors) over S whose special locus is E .

Proposition 4.7. *Assume $n \in \{3, 4, 5\}$. For any effective Cartier divisor $E \subset S$, the morphism of stacks \mathbf{a} defined in Proposition 4.1 induces an equivalence of groupoids*

$$\mathfrak{M}_n^{\text{GM}}(S, E) \xrightarrow{\sim} \mathfrak{M}_{n+5}^{\text{Lag}}(S, 2E).$$

Proof. By Proposition 4.1, the morphism of stacks \mathbf{a} doubles the special locus, hence induces a functor between the groupoids $\mathfrak{M}_n^{\text{GM}}(S, E) \rightarrow \mathfrak{M}_{n+5}^{\text{Lag}}(S, 2E)$. Let us show that the construction of Sec. 4.2 defines its quasi-inverse functor. This construction is clearly functorial, so it remains to consider its compositions with \mathbf{a} .

Let us start with a family of smooth normalized GM data $(S, \mathscr{W}, \mathscr{V}_6, \mathscr{V}_5, \mu, \mathbf{q})$ with special locus E and let $(S, \mathscr{V}_6, \mathscr{V}_5, \mathscr{A})$ be the family of Lagrangian data obtained by applying the morphism \mathbf{a} . Its special locus is $2E$ by Proposition 4.1. Applying the construction of Sec. 4.2 to $(S, \mathscr{V}_6, \mathscr{V}_5, \mathscr{A})$, we must show that the family of GM data that we get is isomorphic to the original one.

The first step of the construction of Sec. 4.2 is the factorization (37) of the morphism φ defined by (27). Comparing it with the factorization of Lemma 4.4 and using the uniqueness of such a factorization (Proposition 2.8), we deduce that the bundles \mathscr{W}' , \mathscr{W} , \mathscr{W}'' , and the maps ν and μ defined by this factorization agree with those in the lemma. It remains to show that the quadratic forms \mathbf{q} agree. This follows from the compatibility of Lemma 4.3 and the uniqueness of the induced quadratic form.

Conversely, let us start with a family of Lagrangian data $(S, \mathscr{V}_6, \mathscr{V}_5, \mathscr{A})$. We produce a family $(S, \mathscr{W}, \mathscr{V}_6, \mathscr{V}_5, \mu, \mathbf{q})$ of smooth normalized GM data by the constructions of Sec. 4.2 and apply the functor \mathbf{a} . In other words, we consider the diagram (28) and our goal is to show that the cohomology bundle $\text{Ker}(f_2)/\text{Im}(f_1)$ of its upper row is isomorphic to the Lagrangian subbundle $\mathscr{A} \subset \bigwedge^3 \mathscr{V}_6$ we started with.

For this, we consider the map

$$f_4 : \mathscr{V}_6 \otimes \mathscr{A} \otimes (\mathscr{V}_6/\mathscr{V}_5)^\vee \rightarrow \bigwedge^3 \mathscr{V}_5 \oplus (\mathscr{V}_6 \otimes \mathscr{W} \otimes (\mathscr{V}_6/\mathscr{V}_5)^\vee)$$

$$v \otimes a \mapsto (\lambda_4(v \wedge a), v \otimes \nu(a)).$$

We have

$$f_2 \circ f_4(v \otimes a)(w') = \lambda_4(v \wedge a) \wedge \mu(w') + \mathbf{q}(v)(\nu(a), w').$$

For $w' = \nu(a')$, we have $\mu(w') = \mu(\nu(a')) = \lambda_3(a')$, hence the right side equals

$$-\mathbf{q}_{\mathcal{A}}(v)(a, a') + \mathbf{q}(v)(\nu(a), \nu(a')) = 0,$$

since \mathbf{q} is induced by $\mathbf{q}_{\mathcal{A}}$ via the map ν . This means that the composition

$$\mathcal{V}_6 \otimes \mathcal{A} \otimes \mathcal{A} \otimes (\mathcal{V}_6/\mathcal{V}_5)^\vee \xrightarrow{\nu} \mathcal{V}_6 \otimes \mathcal{A} \otimes \mathcal{W} \otimes (\mathcal{V}_6/\mathcal{V}_5)^\vee \xrightarrow{f_2 \circ f_4} \det(\mathcal{V}_6)$$

vanishes. Since ν is surjective on the complement of a Cartier divisor, it follows from Lemma 2.7 that $f_2 \circ f_4 = 0$. Therefore, the map f_4 factors through the kernel of f_2 .

Furthermore, the restriction of f_4 to $\mathcal{V}_5 \otimes \mathcal{A} \otimes (\mathcal{V}_6/\mathcal{V}_5)^\vee$ can be rewritten as

$$f_4(v \otimes a) = (\lambda_4(v \wedge a), v \otimes \nu(a)) = (-v \wedge \lambda_3(a), v \otimes \nu(a)) = f_1(v \wedge \nu(a)),$$

hence the composition $\mathcal{V}_6 \otimes \mathcal{A} \otimes (\mathcal{V}_6/\mathcal{V}_5)^\vee \xrightarrow{f_4} \text{Ker}(f_2) \rightarrow \text{Ker}(f_2)/\text{Im}(f_1)$ factors as

$$\mathcal{V}_6 \otimes \mathcal{A} \otimes (\mathcal{V}_6/\mathcal{V}_5)^\vee \xrightarrow{\lambda} \mathcal{A} \longrightarrow \text{Ker}(f_2)/\text{Im}(f_1).$$

Finally, by the commutativity of (33), we have

$$f_3 \circ f_4(v \otimes a) = f_3(\lambda_4(v \wedge a), v \otimes \nu(a)) = \lambda_4(v \wedge a) + v \wedge \mu(\nu(a)) = \lambda(v)a.$$

Together with the above observation, this means that the composition

$$\mathcal{A} \rightarrow \text{Ker}(f_2)/\text{Im}(f_1) \xrightarrow{f_3} \wedge^3 \mathcal{V}_6$$

is the embedding of \mathcal{A} . Since both \mathcal{A} and $\text{Ker}(f_2)/\text{Im}(f_1)$ are Lagrangian subbundles in $\wedge^3 \mathcal{V}_6$, they are isomorphic. \square

Remark 4.8. The same argument proves that there is an equivalence $\mathfrak{M}_{2,\text{ord,ss}}^{\text{GM}}(S, E) \xrightarrow{\sim} \mathfrak{M}_7^{\text{Lag}}(S, 2E)$ of groupoids, where $\mathfrak{M}_{2,\text{ord,ss}}^{\text{GM}} \subset \mathfrak{M}_{2,\text{ord}}^{\text{GM}}$ is the substack defined by (25).

We finish this section by stating a combination of the above results (including Theorem 3.7) which is a simplified version of Proposition 4.7.

Corollary 4.9. *Let $(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$ be a family of Lagrangian data of rank $n + 5$, avoiding decomposable vectors, and such that its special locus is a double Cartier divisor $S_{\text{Lag, spe}} = 2E$. There is a unique family of smooth polarized GM varieties $(S, \mathcal{X} \rightarrow S, \mathcal{H})$ such that $(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$ is obtained from the corresponding family of GM data by the morphism \mathbf{a} .*

This corollary will be used later for constructing interesting families of GM varieties.

5. Descriptions as Global Quotients

We describe the moduli stacks $\mathfrak{M}_n^{\text{GM}}$ and $\mathfrak{M}_n^{\text{Lag}}$ (defined in Secs. 3.1 and 3.5) as global quotients stacks and derive a description of their coarse moduli spaces as the corresponding GIT quotients.

5.1. EPW sextics

Let V_6 be a 6-dimensional \mathbf{k} -vector space and let $A \subset \wedge^3 V_6$ be a Lagrangian subspace for the $\det(V_6)$ -valued symplectic form defined by wedge product.

Definition 5.1. For any integer ℓ , we set

$$Y_A^{\geq \ell} := \{[v] \in \mathbf{P}(V_6) \mid \dim(A \cap (v \wedge \wedge^2 V_6)) \geq \ell\} \tag{40}$$

and endow it with a scheme structure as in [15, Sec. 2]. The locally closed subsets

$$Y_A^\ell := Y_A^{\geq \ell} \setminus Y_A^{\geq \ell+1} \tag{41}$$

of $\mathbf{P}(V_6)$ form the Eisenbud–Popescu–Walter (EPW) stratification and the sequence of inclusions

$$\mathbf{P}(V_6) = Y_A^{\geq 0} \supset Y_A^{\geq 1} \supset Y_A^{\geq 2} \supset \dots$$

is called the EPW sequence. When the scheme $Y_A := Y_A^{\geq 1}$ is not the whole space $\mathbf{P}(V_6)$, it is a sextic hypersurface [15, (1.8)] called an EPW sextic. The scheme $Y_A^{\geq 2}$ is nonempty and has everywhere dimension ≥ 2 [15, (2.9)].

The following theorem gathers various results of O’Grady’s (see [6, Theorem B.2]; all these results were proved for $\mathbf{k} = \mathbf{C}$ but, by the Lefschetz principle, they extend to any field \mathbf{k} of characteristic zero).

Theorem 5.2 (O’Grady). *Let $A \subset \wedge^3 V_6$ be a Lagrangian subspace. If A contains no decomposable vectors, that is, if $\mathbf{P}(A) \cap \text{Gr}(3, V_6) = \emptyset$, then*

- (a) Y_A is an integral normal sextic hypersurface in $\mathbf{P}(V_6)$;
- (b) $Y_A^{\geq 2} = \text{Sing}(Y_A)$ is an integral normal Cohen–Macaulay surface of degree 40;
- (c) $Y_A^{\geq 3} = \text{Sing}(Y_A^{\geq 2})$ is finite and smooth, and is empty for A general;
- (d) $Y_A^{\geq 4}$ is empty.

Remark 5.3. It follows that if the Lagrangian subspace A contains no decomposable vectors, we have $A \cap (v_0 \wedge \wedge^2 V_6) = 0$ for v_0 general in V_6 . We used this observation in the proof of Proposition 4.6.

If $A \subset \wedge^3 V_6$ is a Lagrangian subspace, its orthogonal $A^\perp \subset \wedge^3 V_6^\vee$ is also a Lagrangian subspace. In the dual projective space $\mathbf{P}(V_6^\vee) = \text{Gr}(5, V_6)$, the EPW sequence for A^\perp can be described in terms of A as

$$Y_{A^\perp}^{\geq \ell} = \{V_5 \in \text{Gr}(5, V_6) \mid \dim(A \cap \wedge^3 V_5) \geq \ell\}. \tag{42}$$

The canonical identification $\text{Gr}(3, V_6) \simeq \text{Gr}(3, V_6^\vee)$ induces an isomorphism between the intersections $\mathbf{P}(A) \cap \text{Gr}(3, V_6)$ and $\mathbf{P}(A^\perp) \cap \text{Gr}(3, V_6^\vee)$ [17, (2.82)]. In particular,

a Lagrangian A contains no decomposable vectors if and only if the same holds for A^\perp .

We will not need this fact, but if A contains no decomposable vectors, the hypersurfaces Y_A and Y_{A^\perp} are projective duals [6, Proposition B.3; 16, Corollary 3.6].

If $\mathbf{k} = \mathbf{C}$ and A contains no decomposable vectors, O’Grady defined in [18, Sec. 1.2] a canonical double cover $\tilde{Y}_A \rightarrow Y_A$ (called the **double EPW sextic**). This construction was generalized in [8, Theorem 5.2] to other EPW strata; it works over an arbitrary field \mathbf{k} of characteristic different from 2 and provides canonical double coverings

$$\tilde{Y}_A^{\geq 0} \rightarrow Y_A^{\geq 0}, \quad \tilde{Y}_A^{\geq 1} \rightarrow Y_A^{\geq 1}, \quad \tilde{Y}_A^{\geq 2} \rightarrow Y_A^{\geq 2}$$

branched over $Y_A^{\geq 1}$, $Y_A^{\geq 2}$, and $Y_A^{\geq 3}$, respectively (the first of these is the usual double covering of $\mathbf{P}(V_6)$ branched over the EPW sextic hypersurface). We denote the quotient stacks of these coverings by their natural involutions by

$$\hat{Y}_A^{\geq \ell} := \tilde{Y}_A^{\geq \ell} / \mu_2. \tag{43}$$

They come with natural maps

$$\rho_A^\ell : \hat{Y}_A^{\geq \ell} \longrightarrow Y_A^{\geq \ell}.$$

For $\ell = 0$, we obtain the root stack of $\mathbf{P}(V_6)$ with respect to the EPW sextic hypersurface.

Consider the natural action of the group $\mathrm{PGL}(V_6)$ on the Lagrangian Grassmannian $\mathrm{LGr}(\wedge^3 V_6)$ and its natural linearization in the line bundle $\mathcal{O}(2)$ (note that the line bundle $\mathcal{O}(1)$ does not admit a linearization). O’Grady showed in [20] that the GIT quotient

$$\overline{\mathbf{M}}^{\mathrm{EPW}} := \mathrm{LGr}(\wedge^3 V_6) // \mathrm{PGL}(V_6)$$

is a coarse GIT moduli space for double EPW sextics. The following lemma will be crucial for us.

Lemma 5.4 (O’Grady). *The hypersurface*

$$\Sigma := \{A \in \mathrm{LGr}(\wedge^3 V_6) \mid A \text{ has decomposable vectors}\}$$

is $\mathrm{PGL}(V_6)$ -invariant and its complement

$$\mathrm{LGr}_{\mathrm{adv}}(\wedge^3 V_6) = \mathrm{LGr}(\wedge^3 V_6) \setminus \Sigma$$

is affine and consists of stable points. In particular, for any $A \in \mathrm{LGr}_{\mathrm{adv}}(\wedge^3 V_6)$, the stabilizer $\mathrm{PGL}(V_6)_A \subset \mathrm{PGL}(V_6)$ is finite.

Proof. The stability statement was proved in [20, Corollary 2.5.1] (over $\mathbf{k} = \mathbf{C}$, but stability is defined over the algebraic closure [14, Definition 1.7] and by the Lefschetz principle, stability over \mathbf{C} implies stability over any algebraically closed field of characteristic zero) and the rest is easy. \square

The hypersurface $\Sigma \subset \mathbf{LGr}(\wedge^3 V_6)$ has degree 42, the degree of $\mathbf{Gr}(3, V_6)$. We denote by Σ^{EPW} the image of Σ in $\overline{\mathbf{M}}^{\text{EPW}}$. Its complement

$$\mathbf{M}^{\text{EPW}} := \overline{\mathbf{M}}^{\text{EPW}} \setminus \Sigma^{\text{EPW}} = \mathbf{LGr}_{\text{adv}}(\wedge^3 V_6) // \mathbf{PGL}(V_6) \tag{44}$$

is affine; it is a coarse moduli space for EPW sextics Y_A such that A has no decomposable vectors.

5.2. The moduli stack of Lagrangian data

We deal here with the easier case of the moduli stack of Lagrangian data.

For each integer $n \in \{2, 3, 4, 5\}$, we consider the following relative versions of some EPW strata (we changed the number ℓ in (42) to $n = 5 - \ell$):

$$S_n := \{(A, V_5) \in \mathbf{LGr}_{\text{adv}}(\wedge^3 V_6) \times \mathbf{P}(V_6^\vee) \mid \dim(A \cap \wedge^3 V_5) = 5 - n\}$$

(one could also define S_n for $n \leq 1$, but it is empty) and

$$\overline{S}_n := \{(A, V_5) \in \mathbf{LGr}_{\text{adv}}(\wedge^3 V_6) \times \mathbf{P}(V_6^\vee) \mid \dim(A \cap \wedge^3 V_5) \in \{5 - n, 6 - n\}\}.$$

The subscheme S_{n-1} of \overline{S}_n is closed and S_n is its open complement. The scheme \overline{S}_n is locally closed in $\mathbf{LGr}_{\text{adv}}(\wedge^3 V_6) \times \mathbf{P}(V_6^\vee)$ and in $\mathbf{LGr}(\wedge^3 V_6) \times \mathbf{P}(V_6^\vee)$. In particular, it is a quasiprojective scheme. We will need the following result.

Lemma 5.5. *For $n \geq 2$, the scheme S_n is smooth and*

$$\dim(S_n) = 60 - \frac{1}{2}(5 - n)(6 - n).$$

For $n \in \{2, 5\}$, the scheme \overline{S}_n is smooth; for $n \in \{3, 4\}$, it is normal and $\text{Sing}(\overline{S}_n) = S_{n-1}$.

Proof. The fiber of the projection $\overline{S}_n \rightarrow \mathbf{LGr}_{\text{adv}}(\wedge^3 V_6)$ over a Lagrangian subspace A with no decomposable vectors is the union $Y_{A^\perp}^{5-n} \cup Y_{A^\perp}^{6-n}$ of strata of the dual EPW stratification associated with A , hence Theorem 5.2 applies. \square

Consider now the action of $\mathbf{PGL}(V_6)$ on the product $\mathbf{LGr}(\wedge^3 V_6) \times \mathbf{P}(V_6^\vee)$. As we noted above, the line bundle $\mathcal{O}(2, 0)$ has a natural linearization. It is clear that $\mathcal{O}(0, 6)$ also admits a linearization. Consequently, for any $m \in \mathbf{Z}$, the line bundle $\mathcal{O}(2m, 6)$ admits a $\mathbf{PGL}(V_6)$ -linearization.

Corollary 5.6. *Take $n \in \{2, 3, 4, 5\}$. For sufficiently large m , the subschemes $\overline{S}_n \subset \mathbf{LGr}(\wedge^3 V_6) \times \mathbf{P}(V_6^\vee)$ and $S_n \subset \overline{S}_n$ consist of $\mathbf{PGL}(V_6)$ -stable points for the $\mathcal{O}(2m, 6)$ -linearization.*

Proof. This follows from Lemma 5.4 and [14, Proposition 2.18] applied to morphisms $\overline{S}_n \rightarrow \mathbf{LGr}_{\text{adv}}(\wedge^3 V_6)$ and $S_n \rightarrow \mathbf{LGr}_{\text{adv}}(\wedge^3 V_6)$. \square

The action of $\mathbf{PGL}(V_6)$ on \overline{S}_n also induces an action of $\mathbf{GL}(V_6)$. The canonical morphism

$$\overline{S}_n / \mathbf{GL}(V_6) \longrightarrow \overline{S}_n / \mathbf{PGL}(V_6)$$

of global quotient stacks is a \mathbb{G}_m -gerbe, because the center $\mathbb{G}_m \subset \mathrm{GL}(V_6)$ acts trivially on $\overline{\mathcal{S}}_n$. In fact, this morphism is the rigidification for the natural embedding of \mathbb{G}_m into the automorphism groups of objects of the stack $\overline{\mathcal{S}}_n/\mathrm{GL}(V_6)$. Recall also that the stack $\mathfrak{M}_{n+5}^{\mathrm{Lag}}$ is the rigidification of the stack $\widetilde{\mathfrak{M}}_{n+5}^{\mathrm{Lag}}$ (Remark 3.16).

Proposition 5.7. *For each $n \in \{2, 3, 4, 5\}$, the moduli stack $\mathfrak{M}_{n+5}^{\mathrm{Lag}}$ of families of Lagrangian data of rank $n + 5$ avoiding decomposable vectors is the global quotient stack*

$$\mathfrak{M}_{n+5}^{\mathrm{Lag}} \simeq \overline{\mathcal{S}}_n/\mathrm{PGL}(V_6).$$

In particular, it is a separated Deligne–Mumford stack of finite presentation over \mathbb{Q} . Its special locus is also a global quotient stack

$$\mathfrak{M}_{n+5, \mathrm{spe}}^{\mathrm{Lag}} \simeq S_{n-1}/\mathrm{PGL}(V_6).$$

The stack $\mathfrak{M}_{n+5}^{\mathrm{Lag}}$ is smooth for $n \in \{2, 5\}$; for $n \in \{3, 4\}$, it is singular along the substack $\mathfrak{M}_{n+5, \mathrm{spe}}^{\mathrm{Lag}}$.

Proof. We first prove that the stack of families of linearized Lagrangian data $\widetilde{\mathfrak{M}}_n^{\mathrm{Lag}}$ is isomorphic to the quotient stack $\overline{\mathcal{S}}_n/\mathrm{GL}(V_6)$ by constructing morphisms in both directions between these stacks.

The scheme $\overline{\mathcal{S}}_n$ comes with the tautological family of Lagrangian data on the trivial bundle $\mathcal{V}_6 = V_6 \otimes \mathcal{O}_{\overline{\mathcal{S}}_n}$ (the Lagrangian subbundle is pulled back from $\mathrm{LGr}(\wedge^3 V_6)$ and the subbundle \mathcal{V}_5 is pulled back from $\mathbf{P}(V_6^\vee) \simeq \mathrm{Gr}(5, V_6)$). The definition of $\overline{\mathcal{S}}_n$ ensures that this family of Lagrangian data has rank $n + 5$ and avoids decomposable vectors. Hence, it induces a morphism $\overline{\mathcal{S}}_n \rightarrow \widetilde{\mathfrak{M}}_{n+5}^{\mathrm{Lag}}$. The morphism is $\mathrm{GL}(V_6)$ -equivariant, hence factors through a map from the quotient stack $\overline{\mathcal{S}}_n/\mathrm{GL}(V_6)$ to $\widetilde{\mathfrak{M}}_{n+5}^{\mathrm{Lag}}$.

Let us construct the inverse. Let S be a scheme and let $(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$ be a family of linearized Lagrangian data of rank $n + 5$, avoiding decomposable vectors. Consider the $\mathrm{GL}(V_6)$ -torsor $f : \widetilde{S} \rightarrow S$ associated with the vector bundle \mathcal{V}_6 , so that the pullback of \mathcal{V}_6 to \widetilde{S} comes with a canonical trivialization $f^*\mathcal{V}_6 \simeq V_6 \otimes \mathcal{O}_{\widetilde{S}}$. The pullbacks of the bundles \mathcal{A} and \mathcal{V}_5 can be considered, respectively, as a Lagrangian subbundle $f^*\mathcal{A} \hookrightarrow \wedge^3 V_6 \otimes \mathcal{O}_{\widetilde{S}}$ and as a corank-1 subbundle $f^*\mathcal{V}_5 \hookrightarrow V_6 \otimes \mathcal{O}_{\widetilde{S}}$. Moreover, these subbundles are $\mathrm{GL}(V_6)$ -equivariant. Together they provide a $\mathrm{GL}(V_6)$ -equivariant map

$$\widetilde{S} \longrightarrow \mathrm{LGr}(\wedge^3 V_6) \times \mathbf{P}(V_6^\vee).$$

As the family $(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$ has rank $n + 5$ and avoids decomposable vectors, the map factors through the subscheme $\overline{\mathcal{S}}_n$. Since this map is $\mathrm{GL}(V_6)$ -equivariant, it induces a map

$$S = \widetilde{S}/\mathrm{GL}(V_6) \rightarrow \overline{\mathcal{S}}_n/\mathrm{GL}(V_6).$$

This construction defines a morphism of stacks $\widetilde{\mathfrak{M}}_{n+5}^{\mathrm{Lag}} \rightarrow \overline{\mathcal{S}}_n/\mathrm{GL}(V_6)$.

It is easy to see that the morphisms we constructed above are mutually inverse, hence define an isomorphism of stacks. Moreover, this isomorphism is compatible with the embeddings of $\mathbb{G}_m(S)$ into the automorphism groups of objects of the stacks $\widetilde{\mathfrak{M}}_{n+5}^{\text{Lag}}$ and $\overline{\mathfrak{S}}_n/\text{GL}(V_6)$. Therefore, the rigidifications of these stacks, $\mathfrak{M}_{n+5}^{\text{Lag}}$ and $\overline{\mathfrak{S}}_n/\text{PGL}(V_6)$, are also isomorphic.

Since $\overline{\mathfrak{S}}_n$ is quasiprojective and, by Corollary 5.6, consists of $\text{PGL}(V_6)$ -stable points, the stack $\overline{\mathfrak{S}}_n/\text{PGL}(V_6)$ is a separated Deligne–Mumford stack of finite presentation over \mathbf{Q} . It is clear that under the isomorphism $\mathfrak{M}_{n+5}^{\text{Lag}} \simeq \overline{\mathfrak{S}}_n/\text{PGL}(V_6)$, the special locus of $\mathfrak{M}_{n+5}^{\text{Lag}}$ corresponds to the substack $\mathfrak{S}_{n-1}/\text{PGL}(V_6)$. The description of the singular locus of $\overline{\mathfrak{S}}_n/\text{PGL}(V_6)$ follows immediately from Lemma 5.5. \square

5.3. The moduli stack of GM varieties

We now describe the moduli stack of smooth GM varieties (which we identify with the moduli stack of smooth normalized GM data).

As before, consider the product $\text{LGr}(\wedge^3 V_6) \times \mathbf{P}(V_6^\vee)$. Set $\mathcal{V}_6 := V_6 \otimes \mathcal{O}$ and let $\mathcal{V}_5 \subset \mathcal{V}_6$ and $\mathcal{A} \subset \wedge^3 \mathcal{V}_6$ be the pullbacks of the tautological subbundles from $\mathbf{P}(V_6^\vee)$ and $\text{LGr}(\wedge^3 V_6)$, respectively. Note that $\wedge^3 \mathcal{V}_6 / \wedge^3 \mathcal{V}_5$ is isomorphic to $\wedge^2 \mathcal{V}_5 \otimes (\mathcal{V}_6 / \mathcal{V}_5)$ via the map λ_3 , so that the subscheme $\mathfrak{S}_n \subset \text{LGr}_{\text{adv}}(\wedge^3 V_6) \times \mathbf{P}(V_6^\vee)$ is just the rank- $(n + 5)$ degeneracy locus of the morphism φ defined by (27). In particular, the scheme $\overline{\mathfrak{S}}_n$ is a Lagrangian intersection locus (as defined in [8, Sec. 4]) for the Lagrangian subbundles

$$\mathcal{A} \hookrightarrow \wedge^3 \mathcal{V}_6 \quad \text{and} \quad \wedge^3 \mathcal{V}_5 \hookrightarrow \wedge^3 \mathcal{V}_6.$$

Following [8, Sec. 4], we denote by \mathcal{C}_n the cokernel sheaf of φ on $\overline{\mathfrak{S}}_n$. By definition of $\overline{\mathfrak{S}}_n$, the rank of \mathcal{C}_n is $5 - n$. Consider the reflexive hull of its top exterior power

$$\mathcal{R}_n := (\wedge^{5-n} \mathcal{C}_n)^{\vee\vee}. \tag{45}$$

For $n \leq 4$, this is a rank-1 reflexive sheaf on $\overline{\mathfrak{S}}_n$, and, for $n = 5$, we have $\mathcal{R}_5 \simeq \mathcal{O}_{\overline{\mathfrak{S}}_5}$. Furthermore, it was shown in the proof of [8, Theorem 4.2] that if \mathcal{L}_n is the line bundle

$$\mathcal{L}_n := (\det(V_6)^{15-n} \otimes \det(\mathcal{A}^\vee) \otimes \det(\mathcal{V}_5^\vee)^{\otimes 6})|_{\overline{\mathfrak{S}}_n}, \tag{46}$$

there is a natural morphism

$$\mathbf{m}_n : \mathcal{R}_n \otimes \mathcal{R}_n \longrightarrow \mathcal{L}_n$$

which, for $n \leq 4$, identifies \mathcal{L}_n with the reflexive hull $(\mathcal{R}_n \otimes \mathcal{R}_n)^{\vee\vee}$, and, for $n = 5$, is just a global section of \mathcal{L}_5 (in fact, $\mathbf{m}_5 = \det(\varphi)$).

Lemma 5.8. *The subscheme $B(\mathbf{m}_n) \subset \overline{\mathfrak{S}}_n$, defined by the ideal image of the map \mathbf{m}_n twisted by \mathcal{L}_n^{-1} , is equal to $\mathfrak{S}_{n-1} \subset \overline{\mathfrak{S}}_n$.*

Proof. By [8, Definition 2.8], the subscheme $B(\mathbf{m}_n)$ coincides (locally over $\overline{\mathfrak{S}}_n$) with the branch locus of the double covering of $\overline{\mathfrak{S}}_n$ associated with the reflexive

sheaf \mathcal{R}_n and the morphism \mathbf{m}_n . Since, by Lemma 5.5, the schemes S_n are smooth of expected dimensions, [8, Corollary 4.8] identifies the branch locus with the scheme S_{n-1} . □

We are therefore in the set-up of Appendix A. Accordingly, we consider the root stack

$$\widehat{S}_n \rightarrow \overline{S}_n$$

of $(\mathcal{R}_n, \mathbf{m}_n)$. For $n = 5$, the stack \widehat{S}_5 is isomorphic to $\sqrt{(\mathcal{L}_5, \det(\varphi))/\overline{S}_5}$, the root stack with respect to the hypersurface $S_4 \subset \overline{S}_5$ in the sense of [2, Sec. B.2]. For $n \leq 4$, the stacky locus S_{n-1} of \widehat{S}_n has codimension $6 - n \geq 2$; in this sense \widehat{S}_n is a *generalized* root stack.

We have the following property.

Lemma 5.9. *The stack \widehat{S}_n is a smooth separated Deligne–Mumford stack. The action of the group $\mathrm{PGL}(V_6)$ on \overline{S}_n lifts to an action on \widehat{S}_n such that the morphism $\widehat{S}_n \rightarrow \overline{S}_n$ is $\mathrm{PGL}(V_6)$ -equivariant.*

Proof. To show that \widehat{S}_n is a smooth and separated Deligne–Mumford stack, it is enough, in view of Proposition A.2, to check that the double étale cover of \overline{S}_n associated with the morphism \mathbf{m}_n and a square root of \mathcal{L}_n (which exists locally over \overline{S}_n) is smooth. This follows from [8, Lemma 3.6 and Proposition 3.7], since $\mathrm{LGr}_{\mathrm{adv}}(\wedge^3 V_6) \times \mathbf{P}(V_6^\vee)$ is smooth and S_n is smooth of expected codimension (Lemma 5.5).

To show that the $\mathrm{PGL}(V_6)$ action on \overline{S}_n lifts to \widehat{S}_n , recall from (A.3) that the stack \widehat{S}_n can be defined as the quotient stack

$$\widehat{S}_n = \widehat{\widehat{S}}_n / \mathbb{G}_m, \quad \text{where } \widehat{\widehat{S}}_n := \mathrm{Spec}_{\overline{S}_n}(\mathcal{O}_{\overline{S}_n}[\mathcal{L}_n^{\pm 1}, \mathcal{R}_n]), \tag{47}$$

the sheaf of algebras $\mathcal{O}_{\overline{S}_n}[\mathcal{L}_n^{\pm 1}, \mathcal{R}_n]$ is defined in (A.2), and the \mathbb{G}_m -action corresponds to its grading. The sheaves \mathcal{R}_n and \mathcal{L}_n and the morphism \mathbf{m}_n are $\mathrm{GL}(V_6)$ -equivariant and the center $\mathbb{G}_m \subset \mathrm{GL}(V_6)$ acts on them with respective weights $3(5 - n)$ and $6(5 - n)$ by (45) and (46). Therefore, the group

$$\mathbb{G}_n := \mathrm{GL}(V_6) / \mu_{3(5-n)} \tag{48}$$

acts on the sheaf of algebras $\mathcal{O}_{\overline{S}_n}[\mathcal{L}_n^{\pm 1}, \mathcal{R}_n]$ defined in (A.2), hence also on its relative spectrum $\widehat{\widehat{S}}_n$, in such a way that the action of its center $\mathbb{G}_m / \mu_{3(5-n)} \simeq \mathbb{G}_m$ corresponds to the grading of the algebra. Therefore, the stack \widehat{S}_n carries an action of the quotient group

$$(\mathrm{GL}(V_6) / \mu_{3(5-n)}) / (\mathbb{G}_m / \mu_{3(5-n)}) \simeq \mathrm{GL}(V_6) / \mathbb{G}_m = \mathrm{PGL}(V_6)$$

and the map $\widehat{S}_n \rightarrow \overline{S}_n$ is $\mathrm{PGL}(V_6)$ -equivariant. □

The argument of the proof of the lemma also has the following useful consequence.

Corollary 5.10. *There is an isomorphism of stacks $\widehat{S}_n/\mathrm{PGL}(V_6) \simeq \widehat{S}_n/G_n$.*

We are now ready to prove the main result of this section.

Theorem 5.11. *For $n \in \{3, 4, 5\}$, the stack of smooth polarized GM varieties $\mathfrak{M}_n^{\mathrm{GM}}$ is isomorphic to the global quotient stack*

$$\mathfrak{M}_n^{\mathrm{GM}} \simeq \widehat{S}_n/\mathrm{PGL}(V_6) \simeq \widehat{S}_n/G_n.$$

In particular, it is a smooth separated Deligne–Mumford stack of finite presentation over \mathbf{Q} .

Proof. The first step is the construction of a morphism of stacks

$$\widehat{S}_n/G_n \simeq \widehat{S}_n/\mathrm{PGL}(V_6) \longrightarrow \mathfrak{M}_n^{\mathrm{GM}}.$$

This is equivalent to the construction of a G_n -equivariant family of smooth GM varieties over \widehat{S}_n , which is accomplished below by a combination of several constructions described earlier.

The natural morphism $\widehat{S}_n \rightarrow \overline{S}_n$ can be factored as the composition

$$\widehat{S}_n \xrightarrow{f} \overline{\overline{S}}_n := \mathrm{Spec}_{\overline{S}_n}(\mathcal{O}_{\overline{S}_n}[\mathcal{L}_n^{\pm 1}]) = \mathrm{Spec}_{\overline{S}_n}\left(\bigoplus_{i \in \mathbf{Z}} \mathcal{L}_n^i\right) \xrightarrow{\theta} \overline{S}_n,$$

where $\theta : \overline{\overline{S}}_n \rightarrow \overline{S}_n$ is the \mathbb{G}_m -torsor associated with the line bundle \mathcal{L}_n and f is the double cover associated by [8, Proposition 2.5] with the reflexive sheaf $\theta^*\mathcal{R}_n$ and the natural morphism

$$\theta^*\mathcal{R}_n \otimes \theta^*\mathcal{R}_n \xrightarrow{\theta^*(\mathbf{m}_n)} \theta^*\mathcal{L}_n \simeq \mathcal{O}.$$

The sheaf $\theta^*\mathcal{R}_n$ is the reflexive sheaf associated with the Lagrangian intersection of the subbundles $\theta^*\mathcal{A}$ and $\theta^*(\wedge^3 \mathcal{V}_5)$ on the scheme $\overline{\overline{S}}_n$. Therefore, by [8, Lemma 3.6, Proposition 3.7], the double cover \widehat{S}_n is smooth.

Recall from [8, Definition 2.8] the notions of branch and ramification loci, $B(f)$ and $R(f)$, for the double cover f . By Lemma 5.8 and [8, Corollary 4.8], we have

$$B(f) = \theta^{-1}(S_{n-1}), \quad R(f) \simeq \theta^{-1}(S_{n-1}) \subset \widehat{S}_n,$$

and the preimage $f^{-1}(B(f))$ is the first-order infinitesimal neighborhood of $R(f)$.

Denote by

$$\mathbf{S} := \mathrm{Bl}_{R(f)}(\widehat{S}_n) \xrightarrow{\beta} \widehat{S}_n$$

the blow up of the scheme \widehat{S}_n along $R(f)$ and let E be its exceptional divisor. The preimage of the subscheme S_{n-1} under the map $\beta \circ f \circ \theta$ is the Cartier divisor $2E$.

The following diagram collects the stacks and morphisms that we constructed:

$$\begin{array}{ccccc}
 \mathbf{S} & \xrightarrow{\beta} & \widehat{\widehat{S}}_n & \xrightarrow[2:1]{f} & \overline{\overline{S}}_n \\
 & & \downarrow \mathbb{G}_m & & \downarrow \theta \\
 & & \widehat{S}_n & \xrightarrow{\text{root}} & \overline{S}_n \longrightarrow \mathrm{LGr}_{\mathrm{adv}}(\wedge^3 V_6) \times \mathbf{P}(V_6^\vee).
 \end{array}$$

The labels “2 : 1”, \mathbb{G}_m , and “root” in the diagram mean that the corresponding arrows are a double covering, \mathbb{G}_m -torsors, and a root stack, respectively.

On $\overline{\overline{S}}_n$, we have the tautological family $(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$ of Lagrangian data described in Proposition 5.7. Its pullback to the blow up \mathbf{S} is a family of Lagrangian data on \mathbf{S} of rank $n + 5$ avoiding decomposable vectors. Its special locus is the preimage of S_{n-1} , that is, the Cartier divisor $2E$. Since this divisor is a double, the construction of Sec. 4.2 applies: by Proposition 4.7, there exists a family of smooth normalized GM data $(\mathbf{S}, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ with special locus E . We claim that this family is the pullback with respect to the blow up morphism $\beta : \mathbf{S} \rightarrow \widehat{\widehat{S}}_n$. Since this is the blow up of a smooth scheme along a smooth center, it is enough to check that the bundles \mathcal{W} , \mathcal{V}_6 , and \mathcal{V}_5 restrict trivially to the fibers of the exceptional divisor E .

For the bundles \mathcal{V}_6 and \mathcal{V}_5 , this is obvious, since they are by construction pullbacks from $\widehat{\widehat{S}}_n$. For \mathcal{W} , it is a bit more complicated. This bundle is constructed in Lemma 4.5 and, according to Proposition 2.12, the restriction of \mathcal{W} to E is a direct sum

$$\mathcal{W}|_E \simeq (\mathcal{W}'_E / \mathcal{K}_E) \oplus \mathcal{K}_E(E).$$

The first summand $\mathcal{W}'_E / \mathcal{K}_E$ is isomorphic to the image of the restriction to E of the pullback to \mathbf{S} of the map $\varphi : \mathcal{A} \rightarrow \wedge^2 \mathcal{V}_5 \otimes (\mathcal{V}_6 / \mathcal{V}_5)$. In particular, it is trivial on the fibers of E . The second summand comes by Proposition 2.12 with a natural isomorphism

$$\mathbf{q}_1 : \mathrm{Sym}^2(\mathcal{K}_E(E)) \xrightarrow{\sim} \mathcal{V}_6 / \mathcal{V}_5.$$

Its target is a line bundle trivial on the fibers of E , hence so is its source. Finally, the fibers of E are projective spaces, hence a line bundle on a fiber, whose square is trivial, is trivial itself. Thus, $\mathcal{K}_E(E)$ is trivial on the fibers of E and so is \mathcal{W} .

We conclude that there is a family of GM data on $\widehat{\widehat{S}}_n$ whose pullback to \mathbf{S} is the family of GM data obtained from the family of Lagrangian data $(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$ by the construction of Sec. 4.2. Let us check that it is G_n -equivariant. The family $(S, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$ is G_n -equivariant as a family of Lagrangian data (note, however, that it is not equivariant as a family of linearized Lagrangian data; see Remark 3.16) because, in Definition 3.15 of a morphism of Lagrangian data, we ask for isomorphisms between *projectivizations* of the appropriate bundles. The construction of Sec. 4.2 is natural, hence the resulting family of GM data on the blow up \mathbf{S} of $\widehat{\widehat{S}}_n$ is also G_n -equivariant.

Finally, the pullback functor for the blow up $\beta : \mathbf{S} \rightarrow \widehat{\mathbf{S}}_n$ is fully faithful, hence the resulting GM data on $\widehat{\mathbf{S}}_n$ is G_n -equivariant (again, it is not equivariant as a family of linearized GM data; see Remark 3.4). Consequently, we obtain a family of smooth normalized GM data on the quotient stack $\widehat{\mathbf{S}}_n/G_n \simeq \widehat{\mathbf{S}}_n/\mathrm{PGL}(V_6)$, which gives the desired map

$$\widehat{\mathbf{S}}_n/\mathrm{PGL}(V_6) \rightarrow \mathfrak{M}_n^{\mathrm{GM}}.$$

Let us construct the morphism in the opposite direction. Recall that $\mathfrak{M}_n^{\mathrm{GM}}$ is a smooth Deligne–Mumford stack by Proposition 3.2. Let

$$S \rightarrow \mathfrak{M}_n^{\mathrm{GM}}$$

be an étale covering by a smooth scheme S , let $(S, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ be the corresponding family of smooth normalized GM data, and let \widetilde{S} be the $\mathrm{GL}(V_6)$ -torsor associated with the rank-6 bundle \mathcal{V}_6 , so that the pullback of \mathcal{V}_6 to \widetilde{S} is the trivial bundle $V_6 \otimes \mathcal{O}_{\widetilde{S}}$.

Let $(\widetilde{S}, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$ be the family of Lagrangian data obtained from $(\widetilde{S}, \mathcal{W}, \mathcal{V}_6, \mathcal{V}_5, \mu, \mathbf{q})$ by the construction of Proposition 4.1. Since \mathcal{V}_6 is trivial, we obtain a morphism

$$\widetilde{S} \rightarrow \mathrm{LGr}(\wedge^3 V_6) \times \mathbf{P}(V_6^\vee)$$

such that the bundles \mathcal{A} and \mathcal{V}_5 are the pullbacks of the tautological bundles. Since $(\widetilde{S}, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$ has rank $n+5$ and avoids decomposable vectors, this morphism factors through $\overline{\mathbf{S}}_n$. Let us show that it also factors through the stack $\widehat{\mathbf{S}}_n \rightarrow \overline{\mathbf{S}}_n$.

If $n = 5$, the stack $\widehat{\mathbf{S}}_n$ is the root stack of the section $\det(\varphi)$ of the line bundle \mathcal{L}_5 , so, by [2, Sec. B.1], it is enough to check that the pullback of the ideal generated by $\det(\varphi)$ is a square. Since this ideal defines the Lagrangian-special locus for the family $(\widetilde{S}, \mathcal{V}_6, \mathcal{V}_5, \mathcal{A})$, it is, by Proposition 4.1, the square of the ideal defining the GM-special locus on \widetilde{S} . The universal property of the root stack gives the required factorization.

If $n < 5$, we apply Proposition A.6. Its assumptions are satisfied because \widetilde{S} is smooth and the locus $B(\mathbf{m}_n)$ associated with the map \mathbf{m}_n is equal to S_{n-1} by Lemma 5.8, so, by Proposition 4.1, its preimage in \widetilde{S} is set-theoretically equal to the GM special locus in \widetilde{S} , which by Lemma 3.13 has codimension at least 2 since $n < 5$.

Therefore, we obtain a morphism $\widetilde{S} \rightarrow \widehat{\mathbf{S}}_n$. This morphism is $\mathrm{GL}(V_6)$ -equivariant by construction (where the group $\mathrm{GL}(V_6)$ acts on $\widehat{\mathbf{S}}_n$ via its quotient $\mathrm{PGL}(V_6)$). Since the center $\mathbb{G}_m \subset \mathrm{GL}(V_6)$ acts trivially on $\widehat{\mathbf{S}}_n$, this morphism factors through a $\mathrm{PGL}(V_6)$ -equivariant morphism $\widetilde{S}/\mathbb{G}_m \rightarrow \widehat{\mathbf{S}}_n$. Passing to quotients by $\mathrm{PGL}(V_6)$, we obtain a morphism

$$S = \widetilde{S}/\mathrm{GL}(V_6) = (\widetilde{S}/\mathbb{G}_m)/\mathrm{PGL}(V_6) \rightarrow \widehat{\mathbf{S}}_n/\mathrm{PGL}(V_6).$$

Furthermore, if we replace the étale covering $S \rightarrow \mathfrak{M}_n^{\text{GM}}$ by another étale covering $S' \rightarrow S \rightarrow \mathfrak{M}_n^{\text{GM}}$, the morphisms $S \rightarrow \widehat{S}_n/\text{PGL}(V_6)$ and $S' \rightarrow \widehat{S}_n/\text{PGL}(V_6)$ are compatible. Therefore, we obtain a morphism

$$\mathfrak{M}_n^{\text{GM}} \rightarrow \widehat{S}_n/\text{PGL}(V_6)$$

which is inverse to the one constructed before.

In view of Proposition 3.2, the only thing that remains to be proved is the separatedness of the stack $\mathfrak{M}_n^{\text{GM}}$. This follows from the fact that the scheme \overline{S}_n provides a covering of the stack $\overline{S}_n/\text{PGL}(V_6)$ in the smooth topology. The morphism $\widehat{S}_n \rightarrow \overline{S}_n$ induced by the morphism

$$\widehat{S}_n/\text{PGL}(V_6) \rightarrow \overline{S}_n/\text{PGL}(V_6) \tag{49}$$

on this covering is proper by Corollary A.4 hence, by [23, Lemma 06TZ], so is the morphism (49). Consequently, the separatedness of $\overline{S}_n/\text{PGL}(V_6)$ (proved in Proposition 5.7) implies the separatedness of $\mathfrak{M}_n^{\text{GM}} \simeq \widehat{S}_n/\text{PGL}(V_6)$. \square

One immediate consequence of Theorem 5.11 is the following.

Corollary 5.12. *For $n \in \{3, 4, 5\}$, the stack $\mathfrak{M}_{n,\text{ord}}^{\text{GM}}$ of ordinary smooth GM varieties of dimension n is isomorphic to the quotient stack $S_n/\text{PGL}(V_6)$. Similarly, the stack $\mathfrak{M}_{2,\text{ord,ss}}^{\text{GM}}$ of ordinary strongly smooth GM surfaces is isomorphic to the quotient stack $S_2/\text{PGL}(V_6)$. These stacks are smooth separated Deligne–Mumford stacks of finite presentation over \mathbf{Q} .*

Proof. The first part follows from Theorem 5.11. The second part follows from Remark 4.8 and Proposition 5.7, since $S_1 = \emptyset$, hence $\overline{S}_2 = S_2$. \square

Theorem 5.11 does not describe the stack $\mathfrak{M}_6^{\text{GM}}$ of smooth GM varieties of dimension 6. However, since every such variety is special, we have the following result.

Corollary 5.13. *We have an equality of stacks $\mathfrak{M}_6^{\text{GM}} = \mathfrak{M}_{6,\text{spe}}^{\text{GM}}$. Therefore, $\mathfrak{M}_6^{\text{GM}}$ is a μ_2 -gerbe over $\mathfrak{M}_{5,\text{ord}}^{\text{GM}} \simeq S_5/\text{PGL}(V_6)$.*

Proof. The first assertion is obvious and the second follows from Lemma 3.13 and Corollary 5.12. \square

5.4. Coarse moduli spaces

We use the global quotient descriptions from previous sections to describe the coarse moduli spaces of GM varieties.

Recall that for any $m \in \mathbf{Z}$, the line bundle $\mathcal{O}(2m, 6)$ on $\text{LGr}(\Lambda^3 V_6) \times \mathbf{P}(V_6^\vee)$ admits a $\text{PGL}(V_6)$ -linearization. This line bundle also admits a $\text{GL}(V_6)$ -linearization and, since for $n \in \{3, 4, 5\}$, the subgroup $\mu_{3(5-n)} \subset \text{GL}(V_6)$ acts trivially on it, this

linearization induces a G_n -linearization (where $G_n = \mathrm{GL}(V_6)/\mu_{3(5-n)}$ was defined in (48)).

Corollary 5.14. *For each $n \in \{2, 3, 4, 5\}$ and for $m \gg 0$, the scheme \widehat{S}_n consists of G_n -stable points for the $\mathcal{O}(2m, 6)$ -linearization.*

Proof. As in Corollary 5.6, this follows from Lemma 5.4 and [14, Proposition 2.18] applied to the morphism $\widehat{S}_n \rightarrow \mathrm{LGr}_{\mathrm{adv}}(\wedge^3 V_6)$. □

In the next theorem, we prove that the stacks $\mathfrak{M}_n^{\mathrm{GM}}$, $\mathfrak{M}_{n,\mathrm{ord}}^{\mathrm{GM}}$, $\mathfrak{M}_{n,\mathrm{spe}}^{\mathrm{GM}}$, and $\mathfrak{M}_{n+5}^{\mathrm{Lag}}$ all admit coarse moduli spaces, which we denote by $\mathbf{M}_n^{\mathrm{GM}}$, $\mathbf{M}_{n,\mathrm{ord}}^{\mathrm{GM}}$, $\mathbf{M}_{n,\mathrm{spe}}^{\mathrm{GM}}$, and $\mathbf{M}_{n+5}^{\mathrm{Lag}}$, respectively, and we describe them as GIT quotients.

Theorem 5.15. (a) *For $n \in \{3, 4, 5, 6\}$, the respective coarse moduli spaces $\mathbf{M}_n^{\mathrm{GM}}$ and $\mathbf{M}_{n+5}^{\mathrm{Lag}}$ of the stacks $\mathfrak{M}_n^{\mathrm{GM}}$ and $\mathfrak{M}_{n+5}^{\mathrm{Lag}}$ are both isomorphic to the quasiprojective GIT quotient*

$$\mathbf{M}_n^{\mathrm{GM}} \simeq \mathbf{M}_{n+5}^{\mathrm{Lag}} \simeq \overline{S}_n // \mathrm{PGL}(V_6)$$

taken with respect to the natural linearization of the line bundle $\mathcal{O}(2m, 6)$ for sufficiently large m .

(b) *For $n \in \{3, 4, 5\}$, the respective coarse moduli spaces $\mathbf{M}_{n,\mathrm{ord}}^{\mathrm{GM}}$ and $\mathbf{M}_{2,\mathrm{ord,ss}}^{\mathrm{GM}}$ of the stacks $\mathfrak{M}_{n,\mathrm{ord}}^{\mathrm{GM}}$ and $\mathfrak{M}_{2,\mathrm{ord,ss}}^{\mathrm{GM}}$ of ordinary GM varieties are isomorphic to the GIT quotients*

$$\mathbf{M}_{n,\mathrm{ord}}^{\mathrm{GM}} \simeq S_n // \mathrm{PGL}(V_6), \quad \mathbf{M}_{2,\mathrm{ord,ss}}^{\mathrm{GM}} \simeq S_2 // \mathrm{PGL}(V_6).$$

(c) *For $n \in \{3, 4, 5, 6\}$, the coarse moduli space $\mathbf{M}_{n,\mathrm{spe}}^{\mathrm{GM}}$ of the stack $\mathfrak{M}_{n,\mathrm{spe}}^{\mathrm{GM}}$ of special GM varieties is isomorphic to the GIT quotient*

$$\mathbf{M}_{n,\mathrm{spe}}^{\mathrm{GM}} \simeq S_{n-1} // \mathrm{PGL}(V_6).$$

Proof. We first prove part (a) for $n \in \{3, 4, 5\}$. Since, by Corollary 5.14, the scheme \widehat{S}_n consists of stable points for the G_n -linearization of the bundle $\mathcal{O}(2m, 6)$, the morphism

$$\widehat{S}_n / G_n \rightarrow \widehat{S}_n // G_n$$

to the corresponding GIT quotient is a tame moduli space [4, Theorem 13.6], hence is a coarse moduli space [4, Remark 7.3]. It remains to recall that $\mathfrak{M}_n^{\mathrm{GM}} \simeq \widehat{S}_n / G_n$ by Theorem 5.11. Similarly, the GIT quotient $\overline{S}_n // \mathrm{PGL}(V_6)$ is the coarse moduli space for the stack $\overline{S}_n / \mathrm{PGL}(V_6)$, which by Proposition 5.7 is isomorphic to $\mathfrak{M}_{n+5}^{\mathrm{Lag}}$. So, it remains to identify the GIT quotients $\widehat{S}_n // G_n$ and $\overline{S}_n // \mathrm{PGL}(V_6)$.

For this note that, since the group $G_m / \mu_{3(5-n)}$ acts on the algebra $\mathcal{O}_{\overline{S}_n}[\mathcal{L}_n^{\pm 1}, \mathcal{R}_n]$ via its grading, we have

$$\widehat{S}_n // (G_m / \mu_{3(5-n)}) = \mathrm{Spec}_{\overline{S}_n}(\mathcal{O}_{\overline{S}_n}[\mathcal{L}_n^{\pm 1}, \mathcal{R}_n]) // (G_m / \mu_{3(5-n)}) \simeq \mathrm{Spec}_{\overline{S}_n}(\mathcal{O}_{\overline{S}_n}) \simeq \overline{S}_n.$$

Moreover, we have $\mathrm{PGL}(V_6) \simeq G_n / (\mathbb{G}_m / \mu_{3(5-n)})$. Therefore,

$$\widehat{S}_n // G_n \simeq (\widehat{S}_n // (\mathbb{G}_m / \mu_{3(5-n)})) // (G_n / (\mathbb{G}_m / \mu_{3(5-n)})) \simeq \overline{S}_n // \mathrm{PGL}(V_6).$$

This proves part (a) for $n \in \{3, 4, 5\}$.

The proof of part (b) is completely analogous, using Corollary 5.12 instead of Theorem 5.11.

Let us prove part (c). By Lemma 3.13 and Remark 3.14, the automorphism group scheme of each object of the stack $\mathfrak{M}_{n,\mathrm{spe}}^{\mathrm{GM}}$ contains the constant group scheme μ_2 and the morphisms of stacks $\mathfrak{M}_{n,\mathrm{spe}}^{\mathrm{GM}} \rightarrow \mathfrak{M}_{n-1,\mathrm{ord}}^{\mathrm{GM}}$ for $n \geq 4$, and $\mathfrak{M}_{3,\mathrm{spe}}^{\mathrm{GM}} \rightarrow \mathfrak{M}_{2,\mathrm{ord,ss}}^{\mathrm{GM}}$ for $n = 3$, are the μ_2 -rigidifications. Therefore, by [2, Theorem C.1.1(4)], they have the same coarse moduli space and we conclude by part (b).

Finally, part (a) for $n = 6$ follows from Corollary 5.13 and part (c). \square

The coarse moduli space for smooth GM sixfolds (and for smooth ordinary GM fivefolds), which according to the above results is the GIT quotient $S_5 / \mathrm{PGL}(V_6)$, can also be constructed directly by following Mumford’s proof for hypersurfaces in the projective space. Moreover, this approach gives the additional information that this moduli space is affine.

Proposition 5.16. *The coarse moduli space for smooth ordinary GM fivefolds and for smooth special GM sixfolds is affine.*

Proof. The argument is classical [14, Proposition 4.2]. A smooth ordinary GM fivefold is by definition a smooth hypersurface of degree 2 in $G := \mathrm{Gr}(2, V_5)$. Inside the projective space $\mathbf{P}(H^0(G, \mathcal{O}_G(2)))$, the subset of points corresponding to sections whose zero locus in G is singular is a hypersurface. This hypersurface is ample, hence its complement $\mathbf{P}(H^0(G, \mathcal{O}_G(2)))^0$ is affine and $\mathrm{SL}(V_5)$ -invariant. The action of the reductive group $\mathrm{SL}(V_5)$ on this affine set is linearizable and since the automorphism group of any smooth ordinary GM fivefold is finite [6, Proposition 3.21(c)], the stabilizers are finite at points of $\mathbf{P}(H^0(G, \mathcal{O}_G(2)))^0$, which is therefore contained in the stable locus.

The coarse moduli space for smooth ordinary GM fivefolds is therefore a dense affine open subset of the projective irreducible 25-dimensional GIT quotient

$$\mathbf{P}(H^0(G, \mathcal{O}_G(2))) // \mathrm{SL}(V_5).$$

This proves the proposition. \square

The affineness properties can be also deduced from Theorem 5.15. Indeed, we have

$$\mathbf{M}_{5,\mathrm{ord}}^{\mathrm{GM}} \simeq \mathbf{M}_{6,\mathrm{spe}}^{\mathrm{GM}} \simeq S_5 // \mathrm{PGL}(V_6)$$

and the scheme $S_5 = (\mathrm{LGr}(\wedge^3 V_6) \times \mathbf{P}(V_6^\vee)) \setminus (\Sigma \cup \{\det(\varphi) = 0\})$ is affine since the divisor $\Sigma \cup \{\det(\varphi) = 0\}$ in $\mathrm{LGr}(\wedge^3 V_6) \times \mathbf{P}(V_6^\vee)$ is ample.

6. Applications

In this section, we work over $\mathbf{k} = \mathbf{C}$.

6.1. The period map

The coarse moduli space \mathbf{M}^{EPW} for double EPW sextics was constructed in (44). It is an affine integral scheme of dimension 20. The composition

$$\pi_n : \mathfrak{M}_n^{\text{GM}} \rightarrow \mathbf{M}_n^{\text{GM}} \xrightarrow{\sim} \overline{\mathcal{S}}_n // \text{PGL}(V_6) \rightarrow \text{LGr}_{\text{adv}}(\wedge^3 V_6) // \text{PGL}(V_6) = \mathbf{M}^{\text{EPW}}$$

defines a morphism from the stack $\mathfrak{M}_n^{\text{GM}}$ of smooth GM varieties, or from its coarse moduli space \mathbf{M}_n^{GM} , to the coarse moduli space \mathbf{M}^{EPW} .

The period map

$$\wp^{\text{EPW}} : \mathbf{M}^{\text{EPW}} \hookrightarrow \mathcal{D}$$

for double EPW sextics was constructed by O’Grady, with values in the appropriate period domain \mathcal{D} ; it is an open embedding by Verbitsky’s Torelli Theorem [19, Theorem 1.3].

Proposition 6.1. *Let $n \in \{4, 6\}$. The map*

$$\wp_n^{\text{GM}} := \wp^{\text{EPW}} \circ \pi_n : \mathfrak{M}_n^{\text{GM}} \rightarrow \mathcal{D} \tag{50}$$

is the period map for the middle cohomology of GM varieties of dimension n .

Proof. This follows from [7, Proposition 5.27]. □

Remark 6.2. Let $n \in \{3, 5\}$. By [7, Proposition 3.1], GM varieties of dimension n have intermediate Jacobians that are 10-dimensional principally polarized abelian varieties. This defines a period map $\wp_n^{\text{GM}} : \mathfrak{M}_n^{\text{GM}} \rightarrow \mathbf{A}_{10}$, where \mathbf{A}_{10} is the coarse moduli space for 10-dimensional principally polarized abelian varieties. In the forthcoming paper [9], we will prove that this map factors as

$$\mathfrak{M}_n^{\text{GM}} \xrightarrow{\pi_n} \mathbf{M}^{\text{EPW}} \xrightarrow{\wp^{\text{EPW}}} \mathcal{D} \longrightarrow \mathcal{D}/r_{\mathcal{D}} \dashrightarrow \mathbf{A}_{10},$$

where $r_{\mathcal{D}}$ is the involution of the domain \mathcal{D} defined by O’Grady in [16] (geometrically, it corresponds to passing from an EPW sextic to its dual EPW sextic). The dashed arrow is expected to be generically injective.

We can use Proposition 6.1 to describe the fibers of \wp_n^{GM} for $n \in \{4, 6\}$: they are the same as the fibers of π_n . The stacks $\widehat{Y}_{A^\perp}^{\geq \ell}$ were defined in (43) and the maps $\rho_{A^\perp}^\ell$ right after.

Corollary 6.3. *If $A \subset \wedge^3 V_6$ is a Lagrangian subspace with no decomposable vectors, there is an isomorphism of stacks*

$$\pi_4^{-1}([A]) \simeq (\rho_{A^\perp}^1)^{-1}(Y_{A^\perp} \setminus Y_{A^\perp}^3) / \text{PGL}(V_6)_A \subset \widehat{Y}_{A^\perp}^{\geq 1} / \text{PGL}(V_6)_A, \tag{51}$$

where $\text{PGL}(V_6)_A$ is the stabilizer of A in $\text{PGL}(V_6)$. Furthermore, the stack $\pi_6^{-1}([A])$ is a μ_2 -gerbe over $Y_{A^\perp}^0 / \text{PGL}(V_6)_A$. In particular, there are isomorphisms of coarse

moduli spaces

$$\pi_4^{-1}([A])_{\text{coarse}} \simeq (Y_{A^\perp} \setminus Y_{A^\perp}^3) // \text{PGL}(V_6)_A \quad \text{and} \quad \pi_6^{-1}([A])_{\text{coarse}} \simeq Y_{A^\perp}^0 // \text{PGL}(V_6)_A.$$

Proof. By definition of the scheme \overline{S}_4 , the fiber of the map $\overline{S}_4 \rightarrow \text{LGr}_{\text{adv}}(\wedge^3 V_6)$ over the point $[A]$ is the union $Y_{A^\perp} \setminus Y_{A^\perp}^3 = Y_{A^\perp}^1 \sqcup Y_{A^\perp}^2$ of two EPW strata, hence the fiber of the composition $\widehat{S}_4 \rightarrow \overline{S}_4 \rightarrow \text{LGr}_{\text{adv}}(\wedge^3 V_6)$ is isomorphic to $(\rho_{A^\perp}^1)^{-1}(Y_{A^\perp} \setminus Y_{A^\perp}^3) \subset \widehat{Y}_{A^\perp}^{\geq 1}$. Thus, the stack $\pi_4^{-1}([A])$ is isomorphic to the quotient stack $(\rho_{A^\perp}^1)^{-1}(Y_{A^\perp} \setminus Y_{A^\perp}^3) / \text{PGL}(V_6)_A$ and its coarse moduli space is $(Y_{A^\perp} \setminus Y_{A^\perp}^3) // \text{PGL}(V_6)_A$.

Similarly, Corollary 5.13 identifies the fiber $\pi_6^{-1}([A])$ with a μ_2 -gerbe over $Y_{A^\perp}^0 // \text{PGL}(V_6)_A$ and its coarse moduli space with $Y_{A^\perp}^0 // \text{PGL}(V_6)_A$. \square

6.2. Complete families of smooth GM varieties

Complete nonisotrivial families of smooth projective varieties are hard to find (especially those parameterized by rational curves) and are interesting for this reason. Using our results, one can construct such families of GM varieties, some parameterized by the projective line.

We start with a simple observation.

Lemma 6.4. *Let $(\mathcal{X} \rightarrow S, \mathcal{H})$ be a family of smooth GM varieties of dimension n over a proper reduced connected scheme S . The map $\pi_n : S \rightarrow \mathbf{M}^{\text{EPW}}$ is constant.*

Proof. This follows from the fact that \mathbf{M}^{EPW} is affine. \square

By Lemma 6.4, any family of smooth GM varieties of dimension n parameterized by a proper reduced connected scheme S corresponds to a fixed Lagrangian subspace $A \subset \wedge^3 V_6$ and varying Plücker hyperplanes $V_5 \subset V_6$. In other words, repeating the argument of Corollary 6.3, we see that such a family corresponds to a morphism

$$S \rightarrow \pi_n^{-1}([A]) = (\rho_{A^\perp}^{5-n})^{-1}(Y_{A^\perp}^{\geq 5-n} \setminus Y_{A^\perp}^{\geq 5-n}) / \text{PGL}(V_6)_A \subset \widehat{Y}_{A^\perp}^{\geq 5-n} / \text{PGL}(V_6)_A.$$

The following result can be used to construct such a map.

Proposition 6.5. *Let $n \in \{3, 4, 5, 6\}$ and let S be a connected reduced scheme. Assume that $f : S \rightarrow Y_{A^\perp}^{\geq 5-n}$ is a nonconstant morphism such that $f^{-1}(Y_{A^\perp}^{\geq 7-n}) = \emptyset$ and that*

$$f^{-1}(Y_{A^\perp}^{\geq 6-n}) \quad \text{is equal to } 2E \text{ for some Cartier divisor } E \text{ on } S. \tag{52}$$

Then there is a nonisotrivial family of smooth GM varieties $\mathcal{X} \rightarrow S$ of dimension n .

Proof. The family $\mathcal{X} \rightarrow S$ exists by Corollary 4.9: indeed, consider the family of Lagrangian data on S given by the trivial bundles $\mathcal{V}_6 = \wedge^3 V_6 \otimes \mathcal{O}_S$ and $\mathcal{A} = A \otimes \mathcal{O}_S$, and take for \mathcal{V}_5 the pullback of the tautological rank-5 bundle on $\mathbf{P}(V_6^\vee)$ via the map $S \xrightarrow{f} Y_{A^\perp}^{\geq 5-n} \hookrightarrow \mathbf{P}(V_6^\vee)$.

This family is not isotrivial, because the corresponding map from S to the coarse moduli space \mathbf{M}_n^{GM} is nonconstant: this map is the composition of f with the quotient morphism $Y_{A^\perp}^{\geq 5-n} \rightarrow Y_{A^\perp}^{\geq 5-n} // \text{PGL}(V_6)_A$; since the group $\text{PGL}(V_6)_A$ is finite and $\dim(f(S)) > 0$, this is clear. \square

It is not easy to find a map satisfying the condition (52). Sometimes, a double covering trick helps.

Example 6.6. Let $L \subset \mathbf{P}(V_6^\vee)$ be a line such that $L \not\subset Y_{A^\perp}^{\geq 1}$ and $L \cap Y_{A^\perp}^{\geq 2} = \emptyset$. Then $L \cap Y_{A^\perp}^{\geq 1}$ is a divisor of degree 6 (because $Y_{A^\perp}^{\geq 1}$ is a sextic hypersurface). Let $\tilde{L} \rightarrow L$ be the normalization of the double cover of L branched over $L \cap Y_{A^\perp}^{\geq 1}$. Then,

- if the intersection $L \cap Y_{A^\perp}^{\geq 1}$ is transverse, \tilde{L} is an integral curve of genus 2;
- if the intersection $L \cap Y_{A^\perp}^{\geq 1}$ has exactly one nonreduced point, and its multiplicity is 2 or 3, \tilde{L} is an integral curve of genus 1;
- in all other cases, each component of \tilde{L} is isomorphic to \mathbf{P}^1 .

A general line falls into the first case. A general line tangent to $Y_{A^\perp}^{\geq 1}$ at a general point falls into the second case. Bitangent lines to $Y_{A^\perp}^{\geq 1}$ (of which there is a 6-dimensional family) fall into the third case.

Applying Proposition 6.5 to any of these families, we obtain a family of smooth GM varieties of dimension 5 over \tilde{L} . It is not isotrivial by Proposition 6.5.

To construct families of GM varieties, one can also apply directly Corollary 6.3.

Example 6.7. Assume $Y_{A^\perp}^{\geq 3} = \emptyset$. There is a family of smooth GM fourfolds of maximal variation parameterized by the double EPW sextic \tilde{Y}_{A^\perp} . Indeed, by Corollary 6.3, there is a map

$$\tilde{Y}_{A^\perp} \longrightarrow \tilde{Y}_{A^\perp} / \mu_2 = \hat{Y}_{A^\perp} \longrightarrow \hat{Y}_{A^\perp} / \text{PGL}(V_6)_A = \pi_4^{-1}([A]) \hookrightarrow \mathfrak{M}_4^{\text{GM}}$$

and we obtain a family of GM fourfolds over \tilde{Y}_{A^\perp} by pulling back the universal family over $\mathfrak{M}_4^{\text{GM}}$.

Since any smooth double EPW sextic contains a uniruled divisor (the Gromov–Witten invariants computed in [21] include the degree of the divisor spanned by deformations of a rational curve of minimal degree on any smooth double EPW sextic, and this degree is nonzero), hence many rational curves, one obtains smooth nonisotrivial families of GM fourfolds parameterized by \mathbf{P}^1 .

Example 6.8. Assume $Y_{A^\perp}^{\geq 3} = \emptyset$. As in the previous example, we can pull back the universal family of GM threefolds by the composition (see (51) for the notation)

$$Y_{A^\perp}^{\geq 2} \xrightarrow{\sim} \hat{Y}_{A^\perp}^{\geq 2} \longrightarrow \hat{Y}_{A^\perp}^{\geq 2} / \text{PGL}(V_6)_A \hookrightarrow \hat{S}_3 / \text{PGL}(V_6) \simeq \mathfrak{M}_3^{\text{GM}}$$

and obtain a family of smooth ordinary GM threefolds with maximal variation parameterized by the projective surface $Y_{A^\perp}^{\geq 2}$.

When A is general, the cotangent bundle of $Y_{A^\perp}^{\geq 2}$ is globally generated [5, Corollary 7.3] hence this surface contains no rational curves. Any Lagrangian A with no decomposable vectors such that $Y_{A^\perp}^{\geq 2}$ contains a rational curve and $Y_{A^\perp}^{\geq 3} = \emptyset$ would give rise to a smooth nonisotrivial families of GM threefolds parameterized by \mathbf{P}^1 , but we do not know any such Lagrangian.

Acknowledgments

We are grateful to Ariyan Javanpeykar and to Alex Perry for interesting discussions and extremely useful comments on preliminary drafts of this paper. A. K. is also grateful to Sergey Gorchinskiy for sharing his understanding of stacks.

The work of Alexander Kuznetsov is supported by the Russian Science Foundation under grant 19-11-00164 and performed in Steklov Mathematical Institute of Russian Academy of Sciences.

Appendix A. The Generalized Root Construction

We discuss a generalization of the root stack construction of [2] which is also a particular case of the *canonical stack* construction, as defined (under another name) in [24, Note 2.9, Proof of Proposition 2.8] and developed in [12].

Let S be a normal irreducible scheme. Let \mathcal{R} be a reflexive sheaf of rank 1 on S such that the reflexive hull

$$\mathcal{L} := (\mathcal{R} \otimes \mathcal{R})^{\vee\vee}$$

is a line bundle. If $\mathcal{L} \simeq \mathcal{M}^{\otimes 2}$ for some line bundle \mathcal{M} on S , there is a scheme

$$\tilde{S} := \text{Spec}(\mathcal{O}_S \oplus (\mathcal{M}^{-1} \otimes \mathcal{R})) \tag{A.1}$$

equipped with a map $\tilde{\rho} : \tilde{S} \rightarrow S$ which is finite of degree 2 and étale over the locally free locus of \mathcal{R} [8, Proposition 2.5], and an involution τ of \tilde{S} over S . Let

$$\hat{S} := \tilde{S}/\mu_2(\tau)$$

be the quotient stack with respect to the μ_2 -action on \tilde{S} generated by τ . There is a natural map $\hat{\rho} : \hat{S} \rightarrow S$ which is an isomorphism over the locally free locus of \mathcal{R} ; over $\text{Sing}(\mathcal{R})$, it is a nilpotent thickening of a μ_2 -gerbe over $\text{Sing}(\mathcal{R})$.

We want to show that the construction that produces the stack \hat{S} from S is more natural in a sense than the construction of the double covering. In particular, it does not require the existence (hence nor the choice) of a square root \mathcal{M} of \mathcal{L} .

The construction is very simple. Slightly generalizing the above set-up, we assume that \mathcal{R} is a reflexive sheaf of rank 1 on S such that the sheaf $(\mathcal{R} \otimes \mathcal{R})^{\vee\vee}$ is locally free and let

$$\mathbf{m} : \mathcal{R} \otimes \mathcal{R} \longrightarrow \mathcal{L}$$

be a nonzero morphism to a line bundle \mathcal{L} . Consider the quasicohherent sheaf

$$\begin{aligned} \mathcal{O}_S[\mathcal{L}^{\pm 1}, \mathcal{R}] &:= \bigoplus_{i \in \mathbf{Z}} (\mathcal{L}^i \oplus (\mathcal{L}^i \otimes \mathcal{R})) \\ &\simeq \dots \oplus \mathcal{L}^{-1} \oplus (\mathcal{L}^{-1} \otimes \mathcal{R}) \oplus \mathcal{O}_S \oplus \mathcal{R} \oplus \mathcal{L} \oplus (\mathcal{L} \otimes \mathcal{R}) \oplus \dots \end{aligned} \tag{A.2}$$

with the \mathbf{Z} -grading defined by

$$\deg(\mathcal{L}^i) = 2i \quad \text{and} \quad \deg(\mathcal{L}^i \otimes \mathcal{R}) = 2i + 1.$$

Lemma A.1. *The morphism \mathbf{m} induces on $\mathcal{O}_S[\mathcal{L}^{\pm 1}, \mathcal{R}]$ a commutative associative \mathcal{O}_S -algebra structure.*

Proof. There is a natural associative algebra structure on the sheaf

$$\mathcal{O}_S[\mathcal{R}] := \mathcal{O}_S \oplus \mathcal{R} \oplus (\mathcal{R} \otimes \mathcal{R})^{\vee\vee} \oplus (\mathcal{R} \otimes \mathcal{R} \otimes \mathcal{R})^{\vee\vee} \oplus (\mathcal{R} \otimes \mathcal{R} \otimes \mathcal{R} \otimes \mathcal{R})^{\vee\vee} \oplus \dots$$

(the associativity follows from the functoriality of the reflexive hull). It is also commutative since the automorphism of $(\mathcal{R} \otimes \mathcal{R})^{\vee\vee}$ induced by the transposition of $\mathcal{R} \otimes \mathcal{R}$ is an automorphism of a reflexive sheaf which is the identity on the locally free locus of \mathcal{R} , hence is itself the identity. Finally, the morphism \mathbf{m} induces a morphism $\mathcal{O}_S[(\mathcal{R} \otimes \mathcal{R})^{\vee\vee}] \rightarrow \mathcal{O}_S[\mathcal{L}] \hookrightarrow \mathcal{O}_S[\mathcal{L}^{\pm 1}]$ of commutative associative algebras and we have

$$\mathcal{O}_S[\mathcal{L}^{\pm 1}, \mathcal{R}] = \mathcal{O}_S[\mathcal{R}] \otimes_{\mathcal{O}_S[(\mathcal{R} \otimes \mathcal{R})^{\vee\vee}]} \mathcal{O}_S[\mathcal{L}^{\pm 1}],$$

because $(\mathcal{R}^{\otimes 2m})^{\vee\vee} \simeq ((\mathcal{R} \otimes \mathcal{R})^{\vee\vee})^{\otimes m}$ and $(\mathcal{R}^{\otimes (2m+1)})^{\vee\vee} \simeq ((\mathcal{R} \otimes \mathcal{R})^{\vee\vee})^{\otimes m} \otimes \mathcal{R}$. □

Consider the quotient stack

$$\widehat{S} := (\text{Spec}_S(\mathcal{O}_S[\mathcal{L}^{\pm 1}, \mathcal{R}])) / \mathbb{G}_m \tag{A.3}$$

for the \mathbb{G}_m -action corresponding to the grading defined above. We will call this stack the root stack of $(\mathcal{R}, \mathbf{m})$.

Proposition A.2. *Let \mathcal{M} be a line bundle on S with an isomorphism $\mathcal{L} \simeq \mathcal{M}^{\otimes 2}$. Consider the morphism*

$$(\mathcal{M}^{-1} \otimes \mathcal{R}) \otimes (\mathcal{M}^{-1} \otimes \mathcal{R}) \simeq \mathcal{M}^{-2} \otimes (\mathcal{R} \otimes \mathcal{R}) \xrightarrow{\mathbf{m}} \mathcal{M}^{-2} \otimes \mathcal{L} \simeq \mathcal{O}_S$$

and the associated double covering $\widetilde{S} \rightarrow S$ defined by (A.1). There is a natural isomorphism of stacks $\widetilde{S}/\mu_2 \simeq \widehat{S}$.

Proof. Consider the sheaf of commutative algebras

$$\mathcal{O}_S[\mathcal{M}^{\pm 1}, \mathcal{R}] := \mathcal{O}_S[\mathcal{L}^{\pm 1}, \mathcal{R}] \oplus (\mathcal{M} \otimes \mathcal{O}_S[\mathcal{L}^{\pm 1}, \mathcal{R}]) \tag{A.4}$$

with multiplication induced by the multiplication in the algebra $\mathcal{O}_S[\mathcal{L}^{\pm 1}, \mathcal{R}]$ and the isomorphism $\mathcal{M} \otimes \mathcal{M} \xrightarrow{\simeq} \mathcal{L} \hookrightarrow \mathcal{O}_S[\mathcal{L}^{\pm 1}, \mathcal{R}]$. This algebra carries a natural

$((\mathbf{Z}/2) \oplus \mathbf{Z})$ -grading induced by the \mathbf{Z} -grading of $\mathcal{O}_S[\mathcal{L}^{\pm 1}, \mathcal{R}]$ and

$$\deg(\mathcal{M}) = (1, 1).$$

This grading corresponds to a $(\mu_2 \times \mathbb{G}_m)$ -action on $\mathrm{Spec}_S(\mathcal{O}_S[\mathcal{M}^{\pm 1}, \mathcal{R}])$.

By the definition (A.4) of the algebra $\mathcal{O}_S[\mathcal{M}^{\pm 1}, \mathcal{R}]$, the μ_2 -action on $\mathrm{Spec}_S(\mathcal{O}_S[\mathcal{M}^{\pm 1}, \mathcal{R}])$ is free and the invariant part is equal to $\mathcal{O}_S[\mathcal{L}^{\pm 1}, \mathcal{R}]$. We get an étale double covering

$$\mathrm{Spec}_S(\mathcal{O}_S[\mathcal{M}^{\pm 1}, \mathcal{R}]) \longrightarrow \mathrm{Spec}_S(\mathcal{O}_S[\mathcal{L}^{\pm 1}, \mathcal{R}]).$$

On the other hand, forgetting the $(\mathbf{Z}/2)$ -grading and keeping the \mathbf{Z} -grading, we see that the i th component of the algebra is isomorphic to $\mathcal{M}^i \otimes (\mathcal{O}_S \oplus (\mathcal{M}^{-1} \otimes \mathcal{R}))$. Therefore, the corresponding \mathbb{G}_m -action on $\mathrm{Spec}_S(\mathcal{O}_S[\mathcal{M}^{\pm 1}, \mathcal{R}])$ is also free and the invariant part is equal to $\mathcal{O}_S \oplus (\mathcal{M}^{-1} \otimes \mathcal{R})$. We get a \mathbb{G}_m -torsor

$$\mathrm{Spec}_S(\mathcal{O}_S[\mathcal{M}^{\pm 1}, \mathcal{R}]) \longrightarrow \mathrm{Spec}_S(\mathcal{O}_S \oplus (\mathcal{M}^{-1} \otimes \mathcal{R})) = \tilde{S}.$$

Combining these maps, we obtain a diagram

$$\begin{array}{ccc} \mathrm{Spec}_S(\mathcal{O}_S[\mathcal{M}^{\pm 1}, \mathcal{R}]) & \xrightarrow{\mu_2} & \mathrm{Spec}_S(\mathcal{O}_S[\mathcal{L}^{\pm 1}, \mathcal{R}]) \\ \downarrow \mathbb{G}_m & & \\ \tilde{S} & & \end{array}$$

where the horizontal arrow is a μ_2 -torsor and the vertical arrow is a \mathbb{G}_m -torsor. It induces a μ_2 -torsor

$$\tilde{S} \rightarrow \mathrm{Spec}_S(\mathcal{O}_S[\mathcal{L}^{\pm 1}, \mathcal{R}]) / \mathbb{G}_m = \hat{S}.$$

It follows that $\hat{S} \simeq \tilde{S} / \mu_2$ and since $\deg(\mathcal{M}^{-1} \otimes \mathcal{R}) = (1, 0) \in (\mathbf{Z}/2) \oplus \mathbf{Z}$, the action of μ_2 on \tilde{S} is induced by the involution of the double covering $\tilde{S} \rightarrow S$. \square

Remark A.3. In the case where $\mathcal{R} = \mathcal{O}_S$ and the morphism $\mathbf{m} : \mathcal{R} \otimes \mathcal{R} \rightarrow \mathcal{L}$ is given by a global section s of the line bundle \mathcal{L} , the stack \hat{S} coincides with the usual root stack $\sqrt{(\mathcal{L}, s)} / S$ defined in [2, Sec. B.2]: this follows from Proposition A.2 applied (étale locally) to the double covering \tilde{S} of S branched over the zero locus of s .

We now discuss some properties of the root stack \hat{S} .

Corollary A.4. *The natural morphism $\hat{\rho} : \hat{S} \rightarrow S$ is proper.*

Proof. The question is local over S , so we may assume we are in the set-up of Proposition A.2 and $\hat{S} = \tilde{S} / \mu_2$. Then \tilde{S} is proper over S by (A.1), hence so is \hat{S} by [23, Lemma 0CQK]. \square

Consider the subscheme $B(\mathbf{m}) \subset S$ defined by the ideal image of the map

$$\mathcal{L}^{-1} \otimes \mathcal{R} \otimes \mathcal{R} \xrightarrow{\mathbf{m}} \mathcal{L}^{-1} \otimes \mathcal{L} = \mathcal{O}_S. \tag{A.5}$$

Proposition A.2 implies the main properties of the root stack \hat{S} .

Corollary A.5. *The natural morphism $\hat{\rho} : \hat{S} \rightarrow S$ is an isomorphism over the complement of $B(\mathbf{m}) \subset S$ and is a nilpotent thickening of a μ_2 -gerbe over $B(\mathbf{m})$.*

Proof. Set $Z := B(\mathbf{m})$. Over $S \setminus Z$, we have an isomorphism $\mathcal{R} \otimes \mathcal{R} \simeq \mathcal{L}$, hence \mathcal{R} is invertible. The double covering $\tilde{S} \rightarrow S$ (which exists locally over S) is therefore étale over $S \setminus Z$, hence its quotient stack $\hat{S} \rightarrow S$ is an isomorphism over $S \setminus Z$.

On the other hand, over Z , the multiplication in the algebra defining \tilde{S} is zero, hence there is a natural embedding $Z \rightarrow \tilde{S}$ over $Z \subset S$, and the schematic preimage of $Z \subset S$ in \tilde{S} is a nilpotent thickening of $Z \subset \tilde{S}$. The μ_2 -action on $Z \subset \tilde{S}$ is trivial, hence gives a μ_2 -gerbe $Z/\mu_2 \hookrightarrow \tilde{S}/\mu_2$ over Z and the preimage of Z in \hat{S} is its nilpotent thickening. □

The following property of the stack \hat{S} is quite useful. It is similar to the universal property of canonical smooth Deligne–Mumford stacks proved in [11, Theorem 4.6] (see also [3, Lemma 2.4.1]).

Proposition A.6. *Let $\hat{S} \xrightarrow{\hat{\rho}} S$ be the root stack defined by (A.3) and let $B(\mathbf{m}) \subset S$ be the subscheme defined by (A.5). Let T be a smooth scheme and let $f : T \rightarrow S$ be a morphism such that $\text{codim}_T(f^{-1}(B(\mathbf{m}))) \geq 2$. There is a unique factorization*

$$f : T \rightarrow \hat{S} \xrightarrow{\hat{\rho}} S.$$

Proof. Consider the scheme

$$T \times_S \text{Spec}_S(\mathcal{O}_S[\mathcal{L}^{\pm 1}, \mathcal{R}]) \simeq \text{Spec}_T(\mathcal{O}_T[f^*\mathcal{L}^{\pm 1}, f^*\mathcal{R}]).$$

The sheaf $\mathcal{M} := (f^*\mathcal{R})^{\vee\vee}$ is a rank-1 reflexive sheaf on a smooth scheme T , hence is a line bundle. Therefore, there is a natural epimorphism

$$f^*\mathcal{R} \twoheadrightarrow \mathcal{M} \otimes \mathcal{I},$$

where \mathcal{I} is an ideal sheaf such that the support of the quotient sheaf \mathcal{O}/\mathcal{I} has codimension at least 2. Furthermore, the morphism $\mathbf{m} : \mathcal{R} \otimes \mathcal{R} \rightarrow \mathcal{L}$ induces the morphism $f^*\mathbf{m} : f^*\mathcal{R} \otimes f^*\mathcal{R} \rightarrow f^*\mathcal{L}$ which factors through the tensor product of the reflexive hulls

$$f^*\mathcal{R} \otimes f^*\mathcal{R} \rightarrow \mathcal{M} \otimes \mathcal{M} \rightarrow f^*\mathcal{L},$$

and is an isomorphism away from $f^{-1}(B(\mathbf{m}))$ and the support of \mathcal{O}/\mathcal{I} , that is, in codimension 1. Since T is smooth, it follows that

$$f^*\mathcal{L} \simeq \mathcal{M}^{\otimes 2}.$$

Therefore, we have a natural morphism of graded \mathcal{O}_T -algebras

$$\mathcal{O}_T[f^*\mathcal{L}^{\pm 1}, f^*\mathcal{R}] \twoheadrightarrow \mathcal{O}_T[\mathcal{M}^{\pm 2}, \mathcal{M} \otimes \mathcal{I}] \hookrightarrow \mathcal{O}_T[\mathcal{M}^{\pm 2}, \mathcal{M}] = \mathcal{O}_T[\mathcal{M}^{\pm 1}].$$

It induces a morphism

$$\begin{aligned} \mathrm{Spec}_T(\mathcal{O}_T[\mathcal{M}^{\pm 1}]) &\rightarrow \mathrm{Spec}_T(\mathcal{O}_T[f^* \mathcal{L}^{\pm 1}, f^* \mathcal{R}]) \\ &= T \times_S \mathrm{Spec}_S(\mathcal{O}_S[\mathcal{L}^{\pm 1}, \mathcal{R}]) \rightarrow \mathrm{Spec}_S(\mathcal{O}_S[\mathcal{L}^{\pm 1}, \mathcal{R}]) \end{aligned}$$

compatible with the \mathbb{G}_m -actions corresponding to the gradings of the algebras. Since the source is a \mathbb{G}_m -torsor over T , passing to the quotients by \mathbb{G}_m , we obtain a morphism $T \rightarrow \widehat{S}$. By construction, the composition $T \rightarrow \widehat{S} \rightarrow S$ is equal to f and the constructed morphism is unique with this property. \square

References

- [1] D. Abramovich, A. Corti and A. Vistoli, Twisted bundles and admissible covers, Special issue in honor of Steven L. Kleiman, *Comm. Algebra* **31** (2003) 3547–3618.
- [2] D. Abramovich, T. Graber and A. Vistoli, Gromov–Witten theory of Deligne–Mumford stacks, *Amer. J. Math.* **130** (2008) 1337–1398.
- [3] D. Abramovich and A. Vistoli, Compactifying the space of stable maps, *J. Amer. Math. Soc.* **15** (2002) 27–75.
- [4] J. Alper, Good moduli spaces for Artin stacks, *Ann. Inst. Fourier (Grenoble)* **63** (2013) 2349–2402.
- [5] O. Debarre, A. Iliev and L. Manivel, On the period map for prime Fano threefolds of degree 10, *J. Algebraic Geom.* **21** (2012) 21–59.
- [6] O. Debarre and A. Kuznetsov, Gushel–Mukai varieties: classification and birationalities, *Algebr. Geom.* **5** (2018) 15–76.
- [7] O. Debarre and A. Kuznetsov, Gushel–Mukai varieties: linear spaces and periods, *Kyoto J. Math.* **59**(4) (2019) 897–953.
- [8] O. Debarre and A. Kuznetsov, Double covers of quadratic degeneracy and Lagrangian intersection loci, *Math. Ann.*, to appear.
- [9] O. Debarre and A. Kuznetsov, Gushel–Mukai varieties: intermediate Jacobians, in preparation.
- [10] D. Eisenbud, *Commutative Algebra with a View Toward Algebraic Geometry*, Graduate Texts in Mathematics, Vol. 150 (Springer-Verlag, New York, 1995).
- [11] B. Fantechi, É. Mann and F. Nironi, Smooth toric Deligne–Mumford stacks, *J. Reine Angew. Math.* **648** (2010) 201–244.
- [12] A. Geraschenko and M. Satriano, A “bottom up” characterization of smooth Deligne–Mumford stacks, *Int. Math. Res. Not. IMRN* **21** (2017) 6469–6483.
- [13] A. Kuznetsov and A. Perry, Derived categories of Gushel–Mukai varieties, *Compos. Math.* **154** (2018) 1362–1406.
- [14] D. Mumford, J. Fogarty and F. Kirwan, *Geometric Invariant Theory*, 3rd edn., *Ergebnisse der Mathematik und ihrer Grenzgebiete*, Vol. 34 (Springer-Verlag, Berlin, 1994).
- [15] K. O’Grady, Irreducible symplectic 4-folds and Eisenbud–Popescu–Walter sextics, *Duke Math. J.* **134** (2006) 99–137.
- [16] K. O’Grady, Dual double EPW-sextics and their periods, *Pure Appl. Math. Q.* **4** (2008) 427–468.
- [17] K. O’Grady, EPW-sextics: taxonomy, *Manuscripta Math.* **138** (2012) 221–272.
- [18] K. O’Grady, Double covers of EPW-sextics, *Michigan Math. J.* **62** (2013) 143–184.
- [19] K. O’Grady, Periods of double EPW-sextics, *Math. Z.* **280** (2015) 485–524.

- [20] K. O’Grady, Moduli of double EPW–sextics, *Mem. Amer. Math. Soc.* **240** (2016) 1136.
- [21] G. Oberdieck, Gromov–Witten invariants of the Hilbert schemes of points of a K3 surface, *Geom. Topol.* **22** (2018) 323–437.
- [22] V. G. Sarkisov, On conic bundle structures, *Izv. Akad. Nauk SSSR Ser. Mat.* **46** (1982) 371–408, 432, English transl.: *Math. USSR Izv.* **20** (1983) 355–390.
- [23] The Stacks Project Authors, The Stacks Project (2018), <https://stacks.math.columbia.edu>.
- [24] A. Vistoli, Intersection theory on algebraic stacks and on their moduli spaces, *Invent. Math.* **97** (1989) 613–670.