



Geometry/Functional Analysis

The Knaster problem and the geometry of high-dimensional cubes

Boris S. Kashin^a, Stanislaw J. Szarek^{b,c}

^a Steklov Mathematical Institute, 8 Gubkina Street, 117966, GSP1, Moscow, Russia

^b Équipe d'analyse fonctionnelle, B.C. 186, Université Paris VI, 4, place Jussieu, 75252 Paris, France

^c Department of Mathematics, Case Western Reserve University, Cleveland, OH 44106-7058, USA

Received 20 March 2003; accepted 22 April 2003

Presented by Mikhaël Gromov

Abstract

We study questions of the following type: *Given positive semi-definite matrix \mathcal{G} , does there exist a sequence of vectors in \mathbb{R}^n whose Grammian equals to \mathcal{G} and which has some specified additional properties (typically related to the sup norm)?* In particular, we show that the answer to the 1947 Knaster problem about real functions on spheres is negative for sufficiently large dimensions. *To cite this article: B.S. Kashin, S.J. Szarek, C. R. Acad. Sci. Paris, Ser. I 336 (2003).*

© 2003 Académie des sciences. Published by Éditions scientifiques et médicales Elsevier SAS. All rights reserved.

Résumé

Le problème de Knaster et la géométrie des cubes en grande dimension. Nous étudions des questions du type suivant : Soit \mathcal{G} une matrice positive semi-définie, existe-t-il une suite de vecteurs dans \mathbb{R}^n dont la matrice de Gram est égale à \mathcal{G} et qui possède certaines propriétés supplémentaires (typiquement liées à la norme sup) ? En particulier, nous montrons que la réponse au problème de Knaster datant de 1947 et concernant les fonctions réelles sur les sphères est négative en dimension suffisamment grande. *Pour citer cet article : B.S. Kashin, S.J. Szarek, C. R. Acad. Sci. Paris, Ser. I 336 (2003).*

© 2003 Académie des sciences. Published by Éditions scientifiques et médicales Elsevier SAS. All rights reserved.

Version française abrégée

Nous considérons l'espace euclidien \mathbb{R}^n muni du produit scalaire $\langle \cdot, \cdot \rangle$ et de la norme induite $|\cdot|$. Comme d'habitude, $S^{n-1} := \{x \in \mathbb{R}^n : |x| = 1\}$ est la sphère unité et $B^n := \{x \in \mathbb{R}^n : |x| \leq 1\}$ la boule unité. Pour $x = (x_j)_{j=1}^n \in \mathbb{R}^n$, nous posons $\|x\|_\infty := \max_{1 \leq j \leq n} |x_j|$ et $\|x\|_1 := \sum_{j=1}^n |x_j|$. La même notation $\|\cdot\|_p$ sera utilisée pour la norme sur tout espace $L_p(\mu)$. Nous écrirons L_p pour $L_p(0, 1)$ et ℓ_p^n pour $(\mathbb{R}^n, \|\cdot\|_p)$. Pour une suite $\mathcal{Z} = (z_1, \dots, z_p)$ dans \mathbb{R}^n , on note $\mathcal{G}_{\mathcal{Z}} := [\langle z_i, z_k \rangle]_{i,k=1}^p$ sa matrice de Gram.

Dans cette Note nous considérons des problèmes du type suivant : quelles sont les conditions sur \mathcal{Z} pour garantir l'existence d'une suite $\mathcal{F} = (f_1, \dots, f_p)$ dans un espace $L_2(\mu)$ vérifiant $\mathcal{G}_{\mathcal{F}} = \mathcal{G}_{\mathcal{Z}}$ et possédant en outre certaines propriétés supplémentaires fixées à l'avance ? Des telles questions apparaissent naturellement dans

E-mail addresses: kashin@mi.ras.ru (B.S. Kashin), szarek@ccr.jussieu.fr (S.J. Szarek).

diverses branches de l'analyse, de la géométrie et des mathématiques appliquées, et motivent les deux résultats décrits ici. Le premier est lié à une question fondamentale concernant les systèmes orthogonaux de fonctions ; il résout aussi un problème qui est apparu indépendamment en théorie du contrôle. Le deuxième résultat résout un problème sur les fonctions réelles sur les sphères posé par Knaster en 1947.

Le point de départ du premier résultat est la question de Olevskiĭ ([11], p. 58) liée aux prolongements de suites de fonctions à des suites orthogonales, qui est équivalente au problème suivant :

Existe-t-il une constante $C > 0$ telle que, pour tout $p \in \mathbb{N}$ et pour toute matrice $p \times p$ positive semi-définie \mathcal{G} vérifiant $0 \leq \mathcal{G} \leq I$, on peut trouver une suite $\mathcal{F} = (f_1, \dots, f_p)$ dans L_∞ telle que $\mathcal{G}_\mathcal{F} = \mathcal{G}$ et que $\max_{1 \leq j \leq p} \|f_j\|_\infty \leq C$?

Ce problème a déjà été considéré (et résolu par l'affirmative sous des hypothèses supplémentaires assez fortes) dans [9]. Nous signalons que des affirmations similaires sont étroitement liées à plusieurs résultats profonds d'analyse fonctionnelle. Par exemple, il découle des théorèmes de Grothendieck et de Pietsch que pour toute suite $\mathcal{V} = (v_1, \dots, v_p)$ dans B^n on peut trouver $\mathcal{F} = (f_1, \dots, f_p)$ dans L_∞ vérifiant $\|f_j\|_\infty \leq \sqrt{\pi/2}$ pour $j = 1, \dots, p$ et telle que $\Delta := \mathcal{G}_\mathcal{F} - \mathcal{G}_\mathcal{V} \geq 0$ (voir, par exemple, [12], Theorem 5.10). Par ailleurs, on voit assez facilement (par exemple grâce au Lemme 3 ci-après) que, en général, on ne peut pas atteindre $\Delta = 0$ si les normes $\|f_j\|_\infty$ restent majorées par une constante universelle. La question suivante (posée également par Megretski [8]) est alors naturelle : « peut-on parvenir, sous les mêmes conditions, à Δ diagonal ? » Le théorème qui suit permet d'y répondre négativement (une démonstration détaillée en sera présentée ailleurs).

Théorème 1A. Soit $p \in \mathbb{N}$. Il existe des vecteurs v_1, \dots, v_p dans un espace euclidien avec $|v_j| \leq 1$, $j = 1, \dots, p$, tels que pour toutes les fonctions $f_1, \dots, f_p \in L_\infty$ vérifiant $\langle f_i, f_k \rangle = \langle v_i, v_k \rangle$ pour $1 \leq i < k \leq p$, on a la minoration $\max_{1 \leq j \leq p} \|f_j\|_\infty \geq c(\log p)^{1/4}$, où $c > 0$ est une constante universelle.

Notons que la réponse est différente si on remplace dans ce qui précède les expressions quadratiques par des expressions bilinéaires. Par exemple, l'affirmation suivante est vraie (et équivalente à la célèbre inégalité de Grothendieck, cf. [3] et voir l'Appendice pour les détails) :

Il existe une constante $C > 0$ telle que, pour tout $p, n \in \mathbb{N}$ et pour toutes suites $(u_1, \dots, u_p), (v_1, \dots, v_p)$ dans B^n , on peut trouver des suites $(f_1, \dots, f_p), (g_1, \dots, g_p)$ dans L_∞ vérifiant $\langle f_i, g_k \rangle = \langle u_i, v_k \rangle$, $\|f_i\|_\infty \leq C$ et $\|g_k\|_\infty \leq C$ pour tout $1 \leq i, k \leq p$.

Le deuxième résultat de cette Note concerne le problème de Knaster, posé en 1946 dans *Le Nouveau Livre Écossais* et en 1947 dans [6] :

Étant donnée une fonction continue $F : S^{n-1} \rightarrow \mathbb{R}^m$ et un ensemble de $p = n - m + 1$ points q_1, \dots, q_p dans S^{n-1} , existe-t-il une rotation $U \in \text{SO}(n)$ telle que $F(Uq_1) = \dots = F(Uq_p)$?

Des cas particuliers du problème sont implicitement présents dans des questions antérieures de Steinhaus et Rademacher, voir [4].

Il a depuis été découvert que la réponse au problème est négative pour $m > 2$ et pour certains valeurs de n pour $m = 2$ (voir [7] pour le premier contre-exemple et [1] pour les résultats les plus récents et une bibliographie). Par ailleurs, dans le cas central des fonctions à valeurs réelles (c.a.d., $m = 1$ et, par conséquent, $p = n$) il y a eu des résultats positifs partiels (voir [2,4,14]). Néanmoins, nous montrons ici que la réponse au problème de Knaster est également négative dans ce cas pour n suffisamment grand (même pour des fonctions convexes). Plus précisément, nous prouvons

Théorème 2A. Soit $p, n \in \mathbb{N}$ vérifiant $n \leq p \lfloor \log(p/2) \rfloor / 32$. Alors il existe $\mathcal{V} = \{v_1, \dots, v_p\} \subset S^{p-1}$ tel que pour toute isométrie $\sigma : \ell_2^p \rightarrow \ell_2^n$, la suite $(\|\sigma v_i\|_\infty)_{i=1}^p$ n'est pas constante.

Nous incluons une démonstration complète (bien que non optimisée).

We consider the Euclidean space \mathbb{R}^n endowed with the scalar product $\langle \cdot, \cdot \rangle$ and the induced norm $|\cdot|$. As usual, $S^{n-1} := \{x \in \mathbb{R}^n : |x| = 1\}$ is the unit sphere and $B^n := \{x \in \mathbb{R}^n : |x| \leq 1\}$ is the unit ball. For $x = (x_j)_{j=1}^n \in \mathbb{R}^n$ we set $\|x\|_\infty := \max_{1 \leq j \leq n} |x_j|$ and $\|x\|_1 := \sum_{j=1}^n |x_j|$. The same notation $\|\cdot\|_p$ shall be used for the norm on any $L_p(\mu)$ -space. We shall write L_p for $L_p(0, 1)$ and ℓ_p^n for $(\mathbb{R}^n, \|\cdot\|_p)$. For a sequence $\mathcal{Z} = (z_1, \dots, z_p)$ in \mathbb{R}^n , $\mathcal{G}_{\mathcal{Z}} := [\langle z_i, z_k \rangle]_{i,k=1}^p$ is its Gram matrix.

In this Note we consider problems of the following type: what are the conditions on \mathcal{Z} so that there exists a sequence of elements $\mathcal{F} = (f_1, \dots, f_p)$ in an $L_2(\mu)$ -space for which $\mathcal{G}_{\mathcal{F}} = \mathcal{G}_{\mathcal{Z}}$ and which has some additional prescribed properties? (Note that, by an elementary argument, the condition $\mathcal{G}_{\mathcal{F}} = \mathcal{G}_{\mathcal{Z}}$ is equivalent to existence of a linear isometry σ from the linear span of \mathcal{Z} into $L_2(\mu)$ such that $\sigma(z_j) = f_j$ for $j = 1, \dots, p$.) Such questions arise naturally in various areas of analysis, geometry and applied mathematics, and serve as a motivation for two results presented here. The first one is related to a fundamental question about orthogonal systems and solves a problem that arose independently in control theory. The second result answers, in the negative and for sufficiently large dimensions, a problem about functions on spheres posed by Knaster in 1947.

The starting point of the first result is the following question: given sequence $\Phi = (\varphi_j)_{j=1}^p$ of functions in $L_2(-1, 0)$ (where $p \in \mathbb{N} \cup \{\infty\}$), when can we extend it to an orthonormal sequence on $L_2(-1, 1)$? An elementary argument leads to the Schur criterion: $I - \mathcal{G}_\Phi \geq 0$. If it is satisfied, one can choose as the extensions of φ_j 's any sequence $\Psi = (\psi_j)_{j=1}^p$ in $L_2(0, 1)$ for which $\mathcal{G}_\Psi = I - \mathcal{G}_\Phi$. Olevskii ([11], p. 58) asked whether it is possible to additionally require that the functions ψ_j be uniformly bounded. This question makes sense also for finite systems and can be rephrased as

Does there exist $C > 0$ such that, for any $p \in \mathbb{N}$ and for any positive semi-definite $p \times p$ matrix \mathcal{G} verifying $0 \leq \mathcal{G} \leq I$, there is a sequence $\mathcal{F} = (f_1, \dots, f_p)$ in L_∞ such that $\mathcal{G}_{\mathcal{F}} = \mathcal{G}$ and $\max_{1 \leq j \leq p} \|f_j\|_\infty \leq C$?

This problem has been investigated already in [9] and solved, in the affirmative, under some additional (rather strong) assumptions. We point out that statements similar to the one above are closely related to several deep facts from functional analysis. For example, it follows from the Grothendieck theorem and the Pietsch factorization theorem that, given sequence $\mathcal{V} = (v_1, \dots, v_p)$ in B^n there is an $\mathcal{F} = (f_1, \dots, f_p)$ in L_∞ with $\|f_j\|_\infty \leq \sqrt{\pi/2}$ for $j = 1, \dots, p$ and such that $\mathcal{G}_{\mathcal{V}} \leq \mathcal{G}_{\mathcal{F}}$ (see, e.g., [12], Theorem 5.10). Consequently, $\mathcal{G}_{\mathcal{V}} = \mathcal{G}_{\mathcal{F}} - \Delta$, where Δ is positive semi-definite. However, as can be easily seen (e.g., from Lemma 3 below), one cannot, in general, have $\mathcal{G}_{\mathcal{V}} = \mathcal{G}_{\mathcal{F}}$ with $\|f_j\|_\infty$ bounded by a universal constant. A natural “next best try” is to aim for $\mathcal{G}_{\mathcal{V}} = \mathcal{G}_{\mathcal{F}} - \Delta$ with Δ diagonal; this question was also posed by Megretski (see [8]). However, this is not possible, either; we have

Theorem 1. *Given $p \in \mathbb{N}$ there exist vectors v_1, \dots, v_p in a Euclidean space with $|v_j| \leq 1$ for $j = 1, \dots, p$ such that whenever $f_1, \dots, f_p \in L_\infty$ verify $\langle f_i, f_k \rangle = \langle v_i, v_k \rangle$ for all $1 \leq i < k \leq p$, then*

$$\max_{1 \leq j \leq p} \|f_j\|_\infty \geq c(\log p)^{1/4},$$

where $c > 0$ is a universal constant.

Sketch of the construction. Let $d \in \mathbb{N}$ and let u_1, \dots, u_m be 1-net of S^{d-1} (i.e., if every $u \in S^{d-1}$ is within 1 of one of the u_j 's); by a standard volumetric argument, it is possible to achieve that with $m \leq 2(4/\sqrt{3})^d$. The sequence (v_j) is constructed by repeating each u_j s times, where s is exponential in d . One shows then that the conditions $\langle f_i, f_k \rangle = \langle v_i, v_k \rangle$ for $i \neq k$ and $\|f_j\|_\infty \leq K$ for $1 \leq j \leq p := ms$ are inconsistent if $K/(\log p)^{1/4}$ is small enough. The details of the argument will be presented elsewhere. \square

Let us point out that if we replace quadratic expressions by bilinear ones in the considerations above, the situation is quite different. For example, the following statement is true, and is easily seen to be equivalent to the famous Grothendieck inequality. (Cf. [3] and see Appendix for details and a derivation of the statement from the existence of very large nearly Euclidean subspaces of ℓ_1^n .)

There exists $C > 0$ such that, for any $p, n \in \mathbb{N}$ and for any sequences (u_1, \dots, u_p) , (v_1, \dots, v_p) in B^n there exist sequences (f_1, \dots, f_p) , (g_1, \dots, g_p) in $L_\infty(0, 1)$ such that $\langle f_i, g_k \rangle = \langle u_i, v_k \rangle$, $\|f_i\|_\infty \leq C$ and $\|g_k\|_\infty \leq C$ for all $1 \leq i, k \leq p$.

The remainder of the note will be devoted to the Knaster problem, for which we shall provide a complete, even if not fully optimized argument. The problem, as stated in the *New Scottish Book* in 1946 and in the 1947 note [6], asks

Given a continuous function $F : S^{n-1} \rightarrow \mathbb{R}^m$ and a configuration of $p = n - m + 1$ points q_1, \dots, q_p in S^{n-1} , is there a rotation $U \in \text{SO}(n)$ such that $F(Uq_1) = \dots = F(Uq_p)$?

Special cases of the problem go back to earlier inquiries by Steinhaus and Rademacher, cf. [4]. It has been since determined that the answer to the Knaster problem is negative for $m > 2$ and for some values of n for $m = 2$; see [7] for the first such result and [1] for an update and references. On the other hand, in the central case of real-valued functions (i.e., $m = 1$ and $p = n$) positive partial results were obtained: $n = 2$ is elementary, $n = 3$ was shown in [2]; the answer is also positive if the points q_1, \dots, q_p form an orthonormal sequence [4,14] or, more generally, an orbit of an Abelian group of rotations. Let us also recall that the case $m = 1$ of the problem, if true, would provide an alternate proof of Dvoretzky theorem on almost spherical sections of convex bodies with very good dependence on parameters (cf. [10]). However, we show here that, for sufficiently large n , the answer to the Knaster problem is negative also for $m = 1$.

We start the proof by introducing some more notation. If $d, n \in \mathbb{N}$ with $d \leq n$, $\mathcal{V} \subset \mathbb{R}^d$ and $\sigma : \mathbb{R}^d \rightarrow \mathbb{R}^n$, we shall say that σ is a *Knaster embedding* of \mathcal{V} into ℓ_∞^n if σ is an isometry of ℓ_2^d into ℓ_2^n and $\|\sigma(\cdot)\|_\infty$ is constant on \mathcal{V} . (One could also require that σ is an isometry just on the linear span of \mathcal{V} .) We then have

Theorem 2. *Given $p \in \mathbb{N}$, there exists $\mathcal{V} = \{v_1, \dots, v_p\} \subset S^{p-1}$ such that there is no Knaster embedding σ of \mathcal{V} into ℓ_∞^n with $n \leq p \lfloor \log(p/2) \rfloor / 32$. In particular, the sequence $(\|\sigma v_i\|_\infty)_{i=1}^p$ cannot be constant if $p = n$ and n is sufficiently large.*

The answer to the Knaster problem is thus negative (for large n) even for convex functions. However, it is conceivable that one may still obtain a proof of Dvoretzky theorem via a Knaster problem-like argument. Our example leaves open the possibility that the following is true

Given a continuous real-valued function F on S^{n-1} and a configuration of points v_1, \dots, v_p in S^{d-1} there is an isometry $\sigma : \mathbb{R}^d \rightarrow \mathbb{R}^n$ such that $F(\sigma v_1) = \dots = F(\sigma v_p)$ provided $n \geq pd$ (resp., if pd/n is small enough, or if $p = d$ and p^2/n is small enough).

In fact, this statement *does hold* if $F = \|\cdot\|_\infty$, as can be checked following our analysis. We separated the roles of p and d in the above since in fact in our example the set \mathcal{V} is concentrated on a small subspace of \mathbb{R}^p , specifically of dimension $d = O(\log p)$. Of course, a small perturbation of the set \mathcal{V} yields a basis of \mathbb{R}^p with no Knaster embedding into ℓ_∞^n . However, the condition number of that basis is very large and, in view of [14], such basis *cannot* be orthonormal.

For the proof of Theorem 2 we need two lemmas. The first of them is well known.

Lemma 3. *Let $d \in \mathbb{N}$ and let E be a d -dimensional subspace of L_2 and $S_E := \{f \in E, \|f\|_2 = 1\}$ its sphere. Then $\max\{\|f\|_\infty : f \in S_E\} \geq \sqrt{d}$,*

Consequently, if $\delta > 0$ and if S is any δ -net of S_E , then $\max\{\|f\|_\infty : f \in S\} \geq (1 - \delta^2/2)\sqrt{d}$. If E is a d -dimensional subspace of ℓ_2^n , the corresponding lower bounds are respectively $\sqrt{d/n}$ and $(1 - \delta^2/2)\sqrt{d/n}$.

Sketch of the proof. Let $(\varphi_j)_{j=1}^d$ be an orthonormal basis of E . Then $\mathcal{S}_\varphi := (\sum_{j=1}^d |\varphi_j|^2)^{1/2}$ is, modulo easily remedied measurability issues, the pointwise supremum of $|f|$ over $f \in S$. On the other hand, it is directly verified

that $\sqrt{d} = \|\mathcal{S}_\varphi\|_2 \leq \|\mathcal{S}_\varphi\|_\infty$. If S is a δ -net of S_E , then the convex hull of S contains $(1 - \delta^2/2)S_E$, which yields the second estimate. The above argument works for L_2 and L_∞ on any probability space; the variant for ℓ_2^n and ℓ_∞^n follows by rescaling. \square

Next, let $w_1, \dots, w_N, \dots, w_{2N} \in S^1$ be consecutive vertices of a regular $2N$ -gon (in particular $w_{N+j} = -w_j$ for $j = 1, \dots, N$) and set $\mathcal{P}_N := \{w_1, \dots, w_N\}$. We then have

Lemma 4. *Let $n, N \in \mathbb{N}$ and let $\sigma : \mathbb{R}^2 \rightarrow \mathbb{R}^n$ be a Knaster embedding of \mathcal{P}_N into ℓ_∞^n . Let A be the common value of $\|\sigma(w_j)\|_\infty$, $j = 1, \dots, N$. Then $A \leq 2/\sqrt{N}$.*

Versions of the above statement can be proved for dimensions higher than 2. They show that for a class of “tight near-embeddings” of S^{d-1} into ℓ_∞^n the lower bound from Lemma 3 gives in fact the correct order. This implies that, for small d , such embedding cannot be “random” since a “typical” value of $\|\cdot\|_\infty$ on S^{d-1} is of order $\sqrt{(1 + \log n)/n}$.

We shall postpone the proof of Lemma 4 for a moment and show how the two lemmas imply Theorem 2. Set $d := \lfloor \log(p/2) \rfloor$ and let \mathcal{S} be a set of cardinality $m < (4/\sqrt{3})^d < p/2$ such that $\mathcal{S} \cup (-\mathcal{S})$ is a 1-net of S^{d-1} (as indicated earlier, the existence of \mathcal{S} follows by a standard volumetric argument; the bound 3^d that usually appears in more general statements would work here also). Next, let $N := p - m$ (note that $N > p/2$) and consider the set \mathcal{P}_N of Lemma 4. Identifying \mathbb{R}^d and \mathbb{R}^2 with appropriate d - and 2-dimensional subspaces of \mathbb{R}^p allows to think of all these sets as subsets of S^{p-1} and to define $\mathcal{V} = \{v_1, \dots, v_p\} := \mathcal{S} \cup \mathcal{P}_N$. Now let $n \in \mathbb{N}$ and let σ be any isometry of ℓ_2^p (or the linear span of \mathcal{V}) into ℓ_2^n . Since $\sigma(\mathcal{S}) \cup (-\sigma(\mathcal{S}))$ is a 1-net in the sphere of the d -dimensional space $E = \sigma(\mathbb{R}^d)$, it follows from Lemma 3 that

$$\max_{1 \leq i \leq p} \|\sigma v_i\|_\infty \geq \frac{1}{2} \sqrt{\frac{d}{n}}.$$

On the other hand, Lemma 4 implies that

$$\min_{1 \leq i \leq p} \|\sigma v_i\|_\infty \leq \frac{2}{\sqrt{N}} < 2 \sqrt{\frac{2}{p}}.$$

Thus the sequence $(\|\sigma v_i\|_\infty)$ cannot be constant if $n \leq pd/32 = p\lfloor \log(p/2) \rfloor / 32$, which proves Theorem 2. An examination of the above argument shows that the smallest value $p = n$ for which it yields a counterexample to the Knaster problem is of order 10^{12} . A more careful (but still using only volumetric methods for estimating sizes of nets) calculation along the same lines allows to reduce this to approximately 3.2×10^4 . We expect that working with nets obtained by more efficient constructions and otherwise fine-tuning the argument may give a counterexample for n of order 10^2 . It would be of interest to narrow the gap between these values and those corresponding to positive results (to date, $n \leq 3$).

Proof of Lemma 4. Set $\sigma_1 := \sigma/A$. Then σ_1 is necessarily of the form $\sigma_1(x) = (\langle x, y_s \rangle)_{s=1}^n$ for some $y_1, \dots, y_n \in \mathbb{R}^2$ verifying (a) $|\langle w_i, y_s \rangle| \leq 1$ for $i = 1, \dots, N$ and $s = 1, \dots, n$; (b) for any $i = 1, \dots, N$ there exists $s_i \in \{1, \dots, n\}$ such that $|\langle w_i, y_{s_i} \rangle| = 1$. Let Q be the convex hull of $\mathcal{P}_N \cup (-\mathcal{P}_N)$ and Q° its polar. Then Q° is a regular $2N$ -gon circumscribed around the unit circle. The condition (a) above is equivalent to “ $y_s \in Q^\circ$ for $s = 1, \dots, n$.” The condition (b) says that for every side of Q° there is an s such that either y_s or $-y_s$ belong to that side, namely s_i for the side tangent to the unit circle at w_i . Since a point may belong to at most two sides of Q° (it does when it is a vertex), there must be at least $k := \lceil N/2 \rceil$ distinct points among the y_{s_i} ’s. Note that, by (b), $|y_{s_i}| \geq 1$ for $i = 1, \dots, N$.

Let us now calculate the Hilbert–Schmidt norm $\|\sigma_1\|_{HS} := (\text{tr}(\sigma_1^* \sigma_1))^{1/2}$ in two ways. First, $\|\sigma_1\|_{HS} = (\sum_{s=1}^n |y_s|^2)^{1/2}$, which, by the observations above, is $\geq (N/2)^{1/2}$. Next, σ_1 being a multiple of an isometry of \mathbb{R}^2 , $\|\sigma_1\|_{HS} = \sqrt{2}/A$. Comparing the two quantities we obtain $A \leq 2/\sqrt{N}$, as required. \square

Appendix

As is well known ([5], see also [13]), for any $\theta \in (0, 1)$ there is a constant $c(\theta) > 0$ such that, for any $N \in \mathbb{N}$, nearly all (in the sense of the invariant measure on the corresponding Grassmannian) $\lceil \theta N \rceil$ -dimensional subspaces $E \subset \mathbb{R}^N$ verify the property $\forall x \in E \quad c(\theta)|x| \leq \|x\|_1/\sqrt{N} \leq |x|$. By duality, the above condition is equivalent to $\forall x \in E \exists y \in E^\perp \quad \|x + y\|_\infty \leq c(\theta)^{-1}|x|/\sqrt{N}$. Applying this with $\theta = 2/3$ yields, for each $n \in \mathbb{N}$, an orthogonal decomposition $F_0 \oplus F_1 \oplus F_2$ of \mathbb{R}^{3n} with $\dim F_i = n$, $i = 0, 1, 2$, such that the above properties hold for $N = 3n$ and for all subspaces of the form $E = F_i + F_k$, $i, k \in \{0, 1, 2\}$ (cf. [12], Corollary 7.4).

Let now (u_1, \dots, u_p) , (v_1, \dots, v_p) be sequences in $B^n \subset \mathbb{R}^n$, which we shall identify (as an inner product space) with F_0 . Applying the preceding with $E = F_0 + F_2$ we obtain a sequence $(y_1, \dots, y_p) \in F_1 = (F_0 + F_2)^\perp$ such that $f_i := u_i + y_i$ verify $\|f_i\|_\infty \leq c(2/3)^{-1}/\sqrt{3n}$ for $i = 1, \dots, p$. Similarly, there are z_k 's in F_2 such that $g_k := v_k + z_k$, $k = 1, \dots, p$, satisfy the same estimate. Since clearly $\langle f_i, g_k \rangle = \langle u_i, v_k \rangle$ for all $1 \leq i, k \leq p$, it is now enough to identify \mathbb{R}^{3n} with, say, the appropriate space of step functions on $(0, 1)$ to obtain the required assertion with $C = c(2/3)^{-1}$. Note that the argument above yields $f_i, g_k \in \ell_\infty^N$ with $N = O(n)$, as opposed to $N = O(p^2)$ which follows from the Caratheodory theorem.

The fact that the Grothendieck inequality is related to existence of near-Euclidean subspaces of L_1 -spaces has been well known to experts in the area (see, e.g., [3] and its references). The argument above lays out a specific very direct and conceptually simple link.

Acknowledgements

Most of this research was done while the first named author was visiting University Paris VI under the support from CNRS (France). The second named author was partially supported by a grant from the NSF (USA). The authors thank G. Pisier for communicating to them the inquiries contained in [8] and G. Aubrun for help with the French version.

References

- [1] W. Chen, Counterexamples to Knaster's conjecture, *Topology* 37 (2) (1998) 401–405.
- [2] E.E. Floyd, Real-valued mappings of spheres, *Proc. Amer. Math. Soc.* 6 (1955) 957–959.
- [3] J.E. Gilbert, T.J. Leih, Factorization, tensor products, and bilinear forms in Banach space theory, in: *Notes in Banach Spaces*, University Texas Press, Austin, TX, 1980, pp. 182–305.
- [4] S. Kakutani, A proof that there exists a circumscribing cube around any bounded closed convex set in \mathbb{R}^3 , *Ann. of Math.* 43 (1942) 739–741.
- [5] B.S. Kashin, The widths of certain finite-dimensional sets and classes of smooth functions, *Izv. Akad. Nauk SSSR Ser. Mat.* 41 (1977) 334–351 (in Russian).
- [6] B. Knaster, Problem 4, *Colloq. Math.* 30 (1947) 30–31.
- [7] V.V. Makeev, Some properties of continuous mappings of spheres and problems in combinatorial geometry, in: *Geometric Questions in the Theory of Functions and Sets*, Kalinin. Gos. Univ., Kalinin, 1986, pp. 75–85 (in Russian).
- [8] A. Megretski, Relaxations of quadratic programs in operator theory and system analysis, in: *Systems, Approximation, Singular Integral Operators, and Related Topics* (Bordeaux, 2000), in: *Oper. Theory Adv. Appl.*, Vol. 129, Birkhäuser, Basel, 2001, pp. 365–392.
- [9] D. Menshoff, Sur les séries de fonctions orthogonales bornées dans leur ensembles, *Mat. Sb.* 3 (45) (1938) 103–120.
- [10] V.D. Milman, A few observations on the connections between local theory and some other fields, in: *Geometric Aspects of Functional Analysis* (1986/87), in: *Lecture Notes in Math.*, Vol. 1317, Springer-Verlag, Berlin, 1988, pp. 283–289.
- [11] A.M. Olevskii, *Fourier Series with Respect to General Orthogonal Systems*, Springer-Verlag, Berlin, 1975.
- [12] G. Pisier, in: *Factorization of Linear Operators and Geometry of Banach Spaces*, in: *CBMS Regional Conf. Ser. in Math.*, Vol. 60, American Mathematical Society, Providence, RI, 1986.
- [13] S.J. Szarek, On Kashin's almost Euclidean orthogonal decomposition of ℓ_1^n , *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys.* 2 (1978) 691–694.
- [14] H. Yamabe, Z. Yujobo, On the continuous function defined on a sphere, *Osaka Math. J.* 2 (1) (1950) 19–22.