

DERIVED CATEGORIES OF FANO THREEFOLDS

ALEXANDER KUZNETSOV

To V.A. Iskovskikh on the occasion of his 70th birthday

ABSTRACT. We consider the structure of the derived categories of coherent sheaves on Fano threefolds with Picard number 1 and describe a strange relation between derived categories of different threefolds. In the Appendix we discuss how the ring of algebraic cycles of a smooth projective variety is related to the Grothendieck group of its derived category.

1. INTRODUCTION

A smooth proper connected algebraic variety V is a Fano variety if the anticanonical class $-K_V$ on V is ample. In dimension 1 the only Fano variety is the projective line \mathbb{P}^1 . In dimension 2 the Fano varieties are known under the name of del Pezzo surfaces. There are 10 deformation classes of these — the projective plane with up to 8 blown up points (in generic position) and the quadric.

An ambitious program of classification of Fano threefolds was initiated by G. Fano in the beginning of 20-th century and was mostly accomplished by V. Iskovskikh in 1979 [Is]. The final stroke of brush was added by Mukai and Umemura in 1983 [MU]. In higher dimensions only some pieces of the classification are known.

There are many interconnections between Fano varieties, which help in the classification problems. For example, a hyperplane section of a Fano n -fold V is a Fano variety of dimension $n - 1$ if the anticanonical class of V is sufficiently large. This is most helpful for classification of Fano varieties with large anticanonical class. On the other hand, there are many birational transformations between different Fano varieties of the same dimension. This also is very useful. For example, the original approach of Fano developed by Iskovskikh was based on these kind of interconnections (the double projection from a line is one of the most important).

The goal of the present paper is to indicate that there are interconnections between some Fano threefolds on a higher level, the level of derived categories. On one side we consider a Fano threefold of index 2 and of degree d , and on the other side a Fano threefold of index 1 and degree $4d + 2$. Then we find in both derived categories an exceptional pair of vector bundles and consider the arising semiorthogonal decompositions. The crucial observation is that the nontrivial components of these decompositions are equivalent.

Actually, this is a very rough formulation. It is well known that in general Fano varieties have nontrivial moduli spaces, so the components of the derived categories which we consider in general vary. So, to be more precise one should say that there is a correspondence in the product of the moduli spaces of both types of Fano threefolds, points of which correspond to Fano threefolds with equivalent nontrivial components of derived categories. Investigation of the structure of this correspondence is an interesting question. We conjecture that this correspondence is dominant over the both moduli spaces and give some speculations about its structure.

1991 *Mathematics Subject Classification.* 14M15, 18E30.

I was partially supported by RFFI grants 05-01-01034, 07-01-00051 and 07-01-92211, INTAS 05-1000008-8118, the Russian Science Support Foundation, and gratefully acknowledge of the support of the Pierre Deligne fund based on his 2004 Balzan prize in mathematics.

Although it is not so easy to prove that this relation holds (actually, we can check this only for $d = 3, 4, 5$, while the cases $d = 1, 2$ are still under the question), it is much more difficult to understand *why* such a relation takes place. We believe that any progress in this direction will be very useful for understanding of the structure of Fano varieties and for the classification in higher dimensions.

2. CLASSIFICATION OF FANO THREEFOLDS

An excellent modern survey of the classification of Fano varieties is given in [IP]. Let us briefly remind those parts of the classification which are the most important for us. We will work over an algebraically closed field k of zero characteristic.

The most important discrete invariant of a Fano variety is its Picard lattice $\text{Pic } V$ which comes with the intersection form and with a distinguished element (the canonical class). In this paper we will be mostly concerned with the (most important) case $\text{Pic } V = \mathbb{Z}$. In this case the distinguished element is represented by a positive integer i_V such that

$$K_V = -i_V H,$$

where H is the positive generator of $\text{Pic } V$, and the intersection form is completely determined by a positive integer

$$d_V = H^{\dim V}.$$

These invariants are known as the **index** and the **degree** of V respectively.

The most general result concerning the index is the following

Theorem 2.1 ([Fu]). *If V is a Fano variety of index i_V then $i_V \leq \dim V + 1$. Moreover*

- if $i_V = \dim V + 1$ then $V = \mathbb{P}^n$;
- if $i_V = \dim V$ then $V = Q^n \subset \mathbb{P}^{n+1}$.

In particular, for threefolds we have

Corollary 2.2. *If V is a Fano threefold then $i_V \leq 4$. Moreover*

- if $i_V = 4$ then $V = \mathbb{P}^3$;
- if $i_V = 3$ then $V = Q^3 \subset \mathbb{P}^4$.

Fano threefolds of index 2 are also known as **del Pezzo threefolds** (since their hyperplane sections are del Pezzo surfaces).

Theorem 2.3. *Let V be a Fano threefold with $\text{Pic } V = \mathbb{Z}$ of index $i_V = 2$ and of degree d_V . Then*

$$1 \leq d_V \leq 5$$

and for each $1 \leq d \leq 5$ there exists a unique deformation class of Fano threefolds Y_d with $\text{Pic } Y_d = \mathbb{Z}$ of index 2 and of degree d . They have the following explicit description:

- $Y_5 = \text{Gr}(2, 5) \cap \mathbb{P}^6 \subset \mathbb{P}^9$ is a linear section of codimension 3 of the Grassmannian $\text{Gr}(2, 5)$ in the Plücker embedding;
- $Y_4 = Q \cap Q' \subset \mathbb{P}^5$ is an intersection of two 4-dimensional quadrics;
- $Y_3 \subset \mathbb{P}^4$ is a cubic hypersurface;
- $Y_2 \rightarrow \mathbb{P}^3$ is a double covering ramified in a quartic;
- Y_1 is a hypersurface of degree 6 in the weighted projective space $\mathbb{P}(3, 2, 1, 1, 1)$.

Now let V be a Fano threefold of index 1. Its general anticanonical section $S \subset V$ is a K3-surface, which comes with a polarization $H_S = H|_S$. It follows that

$$d_V = H^3 = H_S^2 = 2g_V - 2$$

for some integer $g_V \geq 2$ which is known as the genus of V .

Theorem 2.4. *Let V be a Fano threefold with $\text{Pic } V = \mathbb{Z}$ of index $i_V = 1$ and of genus g_V . Then*

$$2 \leq g_V \leq 12, \quad g_V \neq 11$$

and for each g in this range there exists a unique deformation class of Fano threefolds X_{2g-2} with $\text{Pic } X_{2g-2} = \mathbb{Z}$ of index 1 and of genus g . They have the following explicit description:

- $X_{22} \subset \mathbb{P}^{13}$ is the zero locus of a global section of the vector bundle $\Lambda^2 \mathcal{U}^* \oplus \Lambda^2 \mathcal{U}^* \oplus \Lambda^2 \mathcal{U}^*$ on the Grassmannian $\text{Gr}(3, 7)$, where \mathcal{U} denotes the tautological rank 3 bundle;
- $X_{18} = \text{G}_2 \text{Gr}(2, 7) \cap \mathbb{P}^{11} \subset \mathbb{P}^{13}$, where $\text{G}_2 \text{Gr}(2, 7)$ is the minimal compact homogeneous space for the simple algebraic group of type G_2 , which can be realized as the zero locus of a global section of the vector bundle $\mathcal{U}^\perp(1)$ on the Grassmannian $\text{Gr}(2, 7)$;
- $X_{16} = \mathbb{P}^{10} \cap \text{SGr}(3, 6) \subset \mathbb{P}^{13}$ is a linear section of codimension 3 of the symplectic Lagrangian Grassmannian $\text{SGr}(3, 6)$ in the Plücker embedding;
- $X_{14} = \mathbb{P}^9 \cap \text{Gr}(2, 6) \subset \mathbb{P}^{14}$ is a linear section of codimension 5 of the Grassmannian $\text{Gr}(2, 6)$ in the Plücker embedding;
- $X_{12} = \mathbb{P}^8 \cap \text{OGr}_+(5, 10) \subset \mathbb{P}^{15}$ is a linear section of codimension 7 of the connected component of the orthogonal Lagrangian Grassmannian $\text{OGr}_+(5, 10)$ in the half-spinor embedding;
- $X_{10} = \mathbb{P}^7 \cap Q \cap \text{Gr}(2, 5) \subset \mathbb{P}^9$ is a quadric section of a linear section of codimension 2 of the Grassmannian $\text{Gr}(2, 5)$ in the Plücker embedding; or $X_{10} \rightarrow Y_5$ is a twofold covering ramified in a quadric;
- $X_8 = Q \cap Q' \cap Q'' \subset \mathbb{P}^6$ is an intersection of three 5-dimensional quadrics;
- $X_6 = Q \cap F_3 \subset \mathbb{P}^5$ is an intersection of a quadric and a cubic;
- $X_4 \subset \mathbb{P}^4$ is a quartic; or $X_4 \rightarrow Q$ is a double cover of a quadric $Q \subset \mathbb{P}^4$ ramified in the intersection of Q with a quartic;
- $X_2 \rightarrow \mathbb{P}^3$ is a double covering ramified in a sextic.

We will also need the following result of S. Mukai.

Theorem 2.5 ([M]). *Assume that the genus g of a Fano threefold X_{2g-2} can be represented as a product $g = r \cdot s$ of two integers. Then on X_{2g-2} there exists a unique stable vector bundle \mathcal{E}_r of rank r with $c_1(\mathcal{E}_r) = -H$ and $c_2(\mathcal{E}_r) = \frac{1}{2}H^2 + (r-s)L$, where L is the class of a line on X_{2g-2} . Moreover, \mathcal{E}_r is exceptional and $H^\bullet(X, \mathcal{E}_r) = 0$.*

The definition of exceptional bundles is given below (see Definition 3.2).

Remark 2.6. When applied to the Fano threefolds X_{2g-2} with $g \geq 6$ this Theorem gives the vector bundles which are the restrictions of the dual tautological bundles from the corresponding Grassmannians.

3. RELATION OF DERIVED CATEGORIES

For an algebraic variety V we denote by $\mathcal{D}^b(V)$ the bounded derived category of coherent sheaves on V . Recall that $\mathcal{D}^b(V)$ is triangulated. We will always assume that V is smooth and projective.

Definition 3.1 ([BK, BO1]). A semiorthogonal decomposition of a triangulated category \mathcal{T} is a sequence of full triangulated subcategories $\mathcal{C}_1, \dots, \mathcal{C}_n$ in \mathcal{T} such that $\text{Hom}_{\mathcal{T}}(\mathcal{C}_i, \mathcal{C}_j) = 0$ for $i > j$ and for every object $T \in \mathcal{T}$ there exists a chain of morphisms $0 = T_n \rightarrow T_{n-1} \rightarrow \dots \rightarrow T_1 \rightarrow T_0 = T$ such that the cone of the morphism $T_k \rightarrow T_{k-1}$ is contained in \mathcal{C}_k for each $k = 1, 2, \dots, n$.

We will write $\mathcal{T} = \langle \mathcal{C}_1, \mathcal{C}_2, \dots, \mathcal{C}_n \rangle$ for a semiorthogonal decomposition of a triangulated category \mathcal{T} with components $\mathcal{C}_1, \mathcal{C}_2, \dots, \mathcal{C}_n$.

Definition 3.2 ([B]). An object $F \in \mathcal{T}$ is called *exceptional* if $\mathrm{Hom}(F, F) = \mathbf{k}$ and $\mathrm{Ext}^p(F, F) = 0$ for all $p \neq 0$. A collection of exceptional objects (F_1, \dots, F_m) is called *exceptional* if $\mathrm{Ext}^p(F_l, F_k) = 0$ for all $l > k$ and all $p \in \mathbb{Z}$.

Proposition 3.3 ([BO1]). *Any exceptional collection F_1, \dots, F_m in $\mathcal{D}^b(V)$ gives a semiorthogonal decomposition*

$$\mathcal{D}^b(V) = \langle \mathcal{C}, F_1, \dots, F_m \rangle$$

where $\mathcal{C} = \langle F_1, \dots, F_m \rangle^\perp = \{F \in \mathcal{D}^b(V) \mid \mathrm{Ext}^\bullet(F_k, F) = 0 \text{ for all } 1 \leq k \leq m\}$ and all the other components are the subcategories of $\mathcal{D}^b(V)$ generated by F_k (each of these is equivalent to $\mathcal{D}^b(\mathbf{k})$, the derived category of \mathbf{k} -vector spaces).

Now let V be a Fano variety of index $i = i_V$. The following result is well known.

Lemma 3.4. *Let V be a Fano variety of index $i = i_V$. Then the collection $\mathcal{O}_V, \mathcal{O}_V(H), \dots, \mathcal{O}_V((i-1)H)$ in $\mathcal{D}^b(V)$ is exceptional.*

Proof: We note that $H^p(V, \mathcal{O}_V(-kH)) = 0$ for $1 \leq k \leq i-1$ and all p by the Kodaira vanishing theorem, hence $\mathrm{Ext}^\bullet(\mathcal{O}_V(lH), \mathcal{O}_V(kH)) = 0$ for $0 \leq k < l \leq i-1$. Similarly, $H^p(V, \mathcal{O}_V) = 0$ for all $p > 0$ by the Kodaira vanishing theorem, while $H^0(V, \mathcal{O}_V) = \mathbf{k}$ since V is connected. Therefore all line bundles on V are exceptional. \square

Corollary 3.5. *For any Fano variety V we have the following semiorthogonal decomposition*

$$\mathcal{D}^b(V) = \langle \mathcal{B}_V, \mathcal{O}_V, \mathcal{O}_V(H), \dots, \mathcal{O}_V((i-1)H) \rangle,$$

where $i = i_V$ is the index of V and $\mathcal{B}_V = \{F \in \mathcal{D}^b(V) \mid H^\bullet(V, F(-kH)) = 0 \text{ for all } 0 \leq k \leq i-1\}$.

In particular, for Fano threefolds Y_d of index 2 we obtain a semiorthogonal decomposition

$$\mathcal{D}^b(Y_d) = \langle \mathcal{B}_{Y_d}, \mathcal{O}_V, \mathcal{O}_V(H) \rangle.$$

For Fano threefolds X_{2g-2} of index 1 and even genus $g = 2t$ we consider the vector bundle \mathcal{E}_2 of rank 2 provided by Theorem 2.5. Applying this Theorem we deduce the following

Lemma 3.6. *Let $X = X_{2g-2}$ be a Fano threefold of index 1 and even genus $g = 2t$. Let $\mathcal{E} = \mathcal{E}_2$ be the vector bundle of rank 2 on X constructed in Theorem 2.5. Then $(\mathcal{E}, \mathcal{O}_X)$ is an exceptional pair on X and we have a semiorthogonal decomposition*

$$\mathcal{D}^b(X_{2g-2}) = \langle \mathcal{A}_{X_{2g-2}}, \mathcal{E}, \mathcal{O}_{X_{2g-2}} \rangle,$$

where $\mathcal{A}_{X_{2g-2}} = \{F \in \mathcal{D}^b(X_{2g-2}) \mid H^\bullet(X_{2g-2}, F) = \mathrm{Ext}^\bullet(\mathcal{E}, F) = 0\}$.

Now we can formulate the Conjecture.

Conjecture 3.7. *Let \mathcal{MF}_d^i be the moduli spaces of Fano threefolds of index i and degree d . Then there is a correspondence $Z_d \subset \mathcal{MF}_d^2 \times \mathcal{MF}_{4d+2}^1$ which is dominant over each factor and such that for any point $(Y_d, X_{4d+2}) \in Z_d$ there is an equivalence of categories*

$$\mathcal{A}_{X_{4d+2}} \cong \mathcal{B}_{Y_d}.$$

The main support for this conjecture is provided by the following results.

Theorem 3.8. *Conjecture 3.7 is true for $d = 3, 4, 5$.*

A proof of Theorem 3.8 will be given in the next section. It consists of a case-by-case analysis, see Corollaries 4.3, 4.6 and 4.9. And now we are going to compare the numerical Grothendieck groups of categories $\mathcal{A}_{X_{4d+2}}$ and \mathcal{B}_{Y_d} .

For a triangulated category \mathcal{T} we denote by $K_0(\mathcal{T})$ its Grothendieck group. It comes with a bilinear Euler form

$$\chi([F], [G]) = \sum_p (-1)^p \dim \operatorname{Ext}^p(F, G).$$

The numerical Grothendieck group $K_0(\mathcal{T})_{num}$ is defined as the quotient $K_0(\mathcal{T})_{num} := K_0(\mathcal{T}) / \ker \chi$. If $\mathcal{T} = \mathcal{D}^b(V)$ we write $K_0(V)$ and $K_0(V)_{num}$ instead of $K_0(\mathcal{D}^b(V))$ and $K_0(\mathcal{D}^b(V))_{num}$ for brevity.

Proposition 3.9. *For all $1 \leq d \leq 5$ there is an isomorphism of numerical Grothendieck groups*

$$K_0(\mathcal{A}_{X_{4d+2}})_{num} \cong K_0(\mathcal{B}_{Y_d})_{num}$$

compatible with the Euler bilinear forms.

Proof: A computation based on the Riemann–Roch Theorem.

Note that by 5.8 the Chern character map $\operatorname{ch} : K_0(X_{2g-2})_{num} \rightarrow \bigoplus_{p=0}^3 H^{2p}(X_{2g-2}, \mathbb{Q})$ identifies the numerical Grothendieck group $K_0(X_{2g-2})_{num}$ with the lattice generated by elements

$$\operatorname{ch}(\mathcal{O}_{X_{2g-2}}) = 1, \quad \operatorname{ch}(\mathcal{O}_H) = H - (g-1)L + \frac{g-1}{3}P, \quad \operatorname{ch}(\mathcal{O}_L) = L + \frac{1}{2}P, \quad \operatorname{ch}(\mathcal{O}_P) = P,$$

where P is the class of a point, and by Riemann–Roch the Euler form can be expressed as

$$\chi(u, v) = \chi_0(u^* \cap v),$$

where $u \mapsto u^*$ is the involution of $\bigoplus_{p=0}^3 H^{2p}(X_{2g-2}, \mathbb{Q})$ given by $(-1)^p$ -multiplication on $H^{2p}(X_{2g-2}, \mathbb{Q})$, and χ_0 is given by the formula

$$\chi_0(x + yH + zL + wP) = x + \frac{g+11}{6}y + \frac{1}{2}z + w.$$

On the other hand, we have $\operatorname{ch}(\mathcal{O}_{X_{2g-2}}) = 1$, and using Theorem 2.5 it is easy to compute

$$\operatorname{ch}(\mathcal{E}) = 2 - H + \frac{g-4}{2}L - \frac{g-10}{12}P.$$

It follows that

$$K_0(\mathcal{A}_{X_{2g-2}})_{num} = \left\langle 1, 2 - H + \frac{g-4}{2}L - \frac{g-10}{12}P \right\rangle^\perp = \left\langle 1 - \frac{g}{2}L + \frac{g-4}{4}P, H - \frac{3g-6}{2}L + \frac{7g-40}{12}P \right\rangle.$$

Computing the form χ on the base vectors we conclude that

$$K_0(\mathcal{A}_{X_{2g-2}})_{num} \cong \mathbb{Z}^2, \quad \chi_{\mathcal{A}} = \begin{pmatrix} 1 - g/2 & -g/2 \\ 3 - g & 1 - g \end{pmatrix}.$$

as a lattice with a bilinear form.

Similarly, by 5.8 the Chern character map $\operatorname{ch} : K_0(Y_d)_{num} \rightarrow \bigoplus_{p=0}^3 H^{2p}(Y_d, \mathbb{Q})$ identifies $K_0(Y_d)_{num}$ with the lattice generated by elements

$$\operatorname{ch}(\mathcal{O}_{Y_d}) = 1, \quad \operatorname{ch}(\mathcal{O}_H) = H - \frac{d}{2}L + \frac{d}{6}P, \quad \operatorname{ch}(\mathcal{O}_L) = L, \quad \operatorname{ch}(\mathcal{O}_P) = P,$$

and it is easy to see that χ_0 is given by the formula

$$\chi_0(x + yH + zL + wP) = x + \frac{d+3}{3}y + z + w.$$

On the other hand, we have $\text{ch}(\mathcal{O}_{Y_d}) = 1$, $\text{ch}(\mathcal{O}_{Y_d}(H)) = 1 + H + \frac{d}{2}L + \frac{d}{6}P$, so it follows that

$$K_0(\mathcal{B}_{Y_d})_{\text{num}} = \left\langle 1, 1 + H + \frac{d}{2}L + \frac{d}{6}P \right\rangle^\perp = \left\langle 1 - L, H - \frac{d}{2}L + \frac{d-6}{6}P \right\rangle.$$

Computing the form χ on the base vectors we conclude that

$$K_0(\mathcal{B}_{Y_d})_{\text{num}} \cong \mathbb{Z}^2, \quad \chi_{\mathcal{B}} = \begin{pmatrix} -1 & -1 \\ 1-d & -d \end{pmatrix}.$$

as a lattice with the bilinear form.

A direct check shows that for the map $\mathbb{Z}^2 \rightarrow \mathbb{Z}^2$ given by the matrix

$$A = \begin{pmatrix} 0 & 1 \\ -1 & -2 \end{pmatrix}$$

we have

$$A^T \cdot \chi_{\mathcal{B}} \cdot A = \begin{pmatrix} -d & -1-d \\ 1-2d & -1-2d \end{pmatrix}$$

which for $g = 2d + 2$ coincides with the matrix of $\chi_{\mathcal{A}}$. Thus the map A gives a required isomorphism $K_0(\mathcal{B}_{Y_d})_{\text{num}} \cong K_0(\mathcal{A}_{X_{4d+2}})_{\text{num}}$ compatible with the Euler forms. \square

4. CASES $d \geq 3$

In this section we prove Theorem 3.8 by a case-by-case analysis.

4.1. The case $d = 5$. Recall that the del Pezzo threefold Y_5 of degree 5 is rigid, so that the moduli space \mathcal{MF}_5^2 is a point. On the contrary, Fano threefolds X_{22} of genus 12 have a 6-dimensional moduli space \mathcal{MF}_{22}^1 . We will show that the correspondence $Z_5 \subset \mathcal{MF}_5^2 \times \mathcal{MF}_{22}^1$ is the whole product. In the other words, we are going to show that for any X_{22} and for the unique Y_5 there is an equivalence $\mathcal{A}_{X_{22}} \cong \mathcal{B}_{Y_5}$. For this we give an explicit description of both categories in question.

For the X_{22} the description is based on the following result. Let \mathcal{E}_2 and \mathcal{E}_3 be the vector bundles of rank 2 and 3 on $X = X_{22}$ provided by Theorem 2.5 for the factorizations $g_X = 12 = 2 \cdot 6 = 3 \cdot 4$. Let $W = H^0(X, \mathcal{E}_3^*)^* \cong \mathbb{k}^7$, so that $X \subset \text{Gr}(3, W)$.

Theorem 4.1 ([K1, K2]). *The bundles $W/\mathcal{E}_3 \otimes \mathcal{O}_X(-1), \mathcal{E}_3^* \otimes \mathcal{O}_X(-1), \mathcal{E}_2, \mathcal{O}_X$ form a full exceptional collection, so that*

$$\mathcal{D}^b(X_{22}) = \langle W/\mathcal{E}_3 \otimes \mathcal{O}_X(-1), \mathcal{E}_3^* \otimes \mathcal{O}_X(-1), \mathcal{E}_2, \mathcal{O}_X \rangle.$$

Moreover, $\text{Hom}(W/\mathcal{E}_3 \otimes \mathcal{O}_X(-1), \mathcal{E}_3^* \otimes \mathcal{O}_X(-1)) = \mathbb{k}^3$, $\text{Ext}^{\neq 0}(W/\mathcal{E}_3 \otimes \mathcal{O}_X(-1), \mathcal{E}_3^* \otimes \mathcal{O}_X(-1)) = 0$, so that

$$\mathcal{A}_{X_{22}} \cong \mathcal{D}^b(\mathbb{Q}_3),$$

where $\mathbb{Q}_3 = (\bullet \rightrightarrows \bullet)$ is the Kronecker quiver with 3 arrows.

On the other hand, for the Y_5 a description of the derived category was given by Orlov. Let \mathcal{U} be the restriction to $Y = Y_5$ of the tautological bundle from the Grassmannian $\text{Gr}(2, 5)$ to Y . Let $W' = H^0(Y, \mathcal{U}^*)^* \cong \mathbb{k}^5$, so that $Y \subset \text{Gr}(2, W')$.

Theorem 4.2 ([Or]). *The bundles $W'/\mathcal{U} \otimes \mathcal{O}_Y(-1), \mathcal{U}, \mathcal{O}_Y, \mathcal{O}_Y(1)$ form a full exceptional collection, so that*

$$\mathcal{D}^b(Y_5) = \langle W'/\mathcal{U} \otimes \mathcal{O}_Y(-1), \mathcal{U}, \mathcal{O}_Y, \mathcal{O}_Y(1) \rangle.$$

Moreover, $\text{Hom}(W'/\mathcal{U} \otimes \mathcal{O}_Y(-1), \mathcal{U}) = \mathbb{k}^3$, $\text{Ext}^{\neq 0}(W'/\mathcal{U} \otimes \mathcal{O}_Y(-1), \mathcal{U}) = 0$, so that $\mathcal{B}_{Y_5} \cong \mathcal{D}^b(\mathbb{Q}_3)$.

From these two results we immediately deduce the required equivalence.

Corollary 4.3. *For any Fano threefold X_{22} of genus 12 and for the unique del Pezzo threefold Y_5 of degree 5 there is an equivalence of categories $\mathcal{A}_{X_{22}} \cong \mathcal{D}^b(\mathbb{Q}_3) \cong \mathcal{B}_{Y_5}$.*

4.2. The case $d = 4$. Recall that the moduli space \mathcal{MF}_4^2 of del Pezzo threefolds Y_4 of degree 4 is isomorphic to the moduli space \mathcal{M}_2 of curves of genus 2. The isomorphism is constructed as follows. Let $Y = Y_4 = Q \cap Q' \subset \mathbb{P}^5$. Consider the pencil of quadrics $\{Q_\lambda\}_{\lambda \in \mathbb{P}^1}$ generated by Q and Q' . If Y is smooth then the generic Q_λ is smooth and there are precisely 6 distinct points $\lambda_1, \dots, \lambda_6 \in \mathbb{P}^1$ for which the quadric Q_λ is degenerate. Consider the twofold covering $C(Y) \rightarrow \mathbb{P}^1$ ramified at the points λ_i . Then $C(Y)$ is a smooth curve of genus 2.

Theorem 4.4 ([BO1, K6]). *The map $\mathcal{MF}_4^2 \rightarrow \mathcal{M}_2$, $Y \mapsto C(Y)$ is an isomorphism. Moreover, there is an equivalence $\mathcal{B}_{Y_4} \cong \mathcal{D}^b(C(Y_4))$.*

Our goal is to show that X_{18} threefolds behave in a similar fashion. Indeed, recall that by definition any $X = X_{18}$ is a linear section of codimension 2 in $\mathbb{G}_2\text{Gr}(2, 7)$. Let $\{\mathcal{X}_\lambda\}_{\lambda \in \mathbb{P}^1}$ be the pencil of hyperplane sections of $\mathbb{G}_2\text{Gr}(2, 7)$ passing through X . Since the projective dual of $\mathbb{G}_2\text{Gr}(2, 7)$ is a hypersurface of degree 6, it follows that there are precisely 6 distinct points $\lambda_1, \dots, \lambda_6 \in \mathbb{P}^1$ for which \mathcal{X}_λ is singular. Consider the twofold covering $C(X) \rightarrow \mathbb{P}^1$ ramified at the points λ_i . Then $C(X)$ is a smooth curve of genus 2. Thus we obtain a map $\mathcal{MF}_{18}^1 \rightarrow \mathcal{M}_2$, $X \mapsto C(X)$.

Theorem 4.5 ([K5]). *There is an equivalence $\mathcal{B}_{X_{18}} \cong \mathcal{D}^b(C(X_{18}))$.*

Combining these results we obtain the case $d = 4$. Let $Z_4 \subset \mathcal{MF}_4^2 \times \mathcal{MF}_{18}^1$ be the graph of the morphism $\mathcal{MF}_{18}^1 \rightarrow \mathcal{M}_2 \cong \mathcal{MF}_4^2$.

Corollary 4.6. *For any pair of threefolds $(Y_4, X_{18}) \in Z_4 \subset \mathcal{MF}_4^2 \times \mathcal{MF}_{18}^1$ we have an equivalence of categories $\mathcal{A}_{X_{18}} \cong \mathcal{B}_{Y_4}$.*

4.3. The case $d = 3$. While in the previous two cases we were able to describe the categories under the question explicitly, for $d = 3$ this is no longer possible. We can only prove an equivalence in this case.

Recall that by definition any $X = X_{14}$ is a linear section of codimension 5 in $\text{Gr}(2, 6)$. Let W be the six-dimensional vector space, so that $\text{Gr}(2, 6) = \text{Gr}(2, W) \subset \mathbb{P}(\Lambda^2 W)$. Then X can be described by a 5-dimensional subspace $A \subset \Lambda^2 W^*$ or, equivalently, by an injective map $\alpha : A \rightarrow \Lambda^2 W$, where A is a fixed vector space of dimension 5. Let us denote the corresponding threefold X_{14} by $X(\alpha)$.

On the other hand, consider the whole space $\mathbb{P}(\Lambda^2 W^*)$, the space of skew-symmetric forms on W . Consider the hypersurface therein consisting of degenerate skew-forms. It is well known that the equation of this hypersurface is given by the Pfaffian polynomial. It follows that it is a cubic hypersurface, which is denoted by $\text{Pf}(W)$ and is called the Pfaffian variety. Certainly, the Pfaffian variety is singular, its singular locus coincides with the set of all skew-forms of rank 2 on W , that is with the Grassmannian $\text{Gr}(2, W^*) \subset \text{Pf}(W) \subset \mathbb{P}(\Lambda^2 W^*)$. However, the codimension of the singular locus in $\mathbb{P}(\Lambda^2 W^*)$ is $14 - 8 = 6$, so for generic α the preimage of $\text{Pf}(W)$ in $\mathbb{P}(A)$ is a smooth cubic hypersurface, which we denote $Y(\alpha)$. Further, associating with a degenerate skew-form on W its kernel, defines a rank 2 subbundle $K \subset W \otimes \mathcal{O}$ on the smooth locus of $\text{Pf}(W)$. Let $E(\alpha) = \alpha^* K \otimes \mathcal{O}_{Y(\alpha)}(1)$, where in the right-hand-side we consider α as a map $Y(\alpha) \rightarrow \text{Pf}(W)$ and $\mathcal{O}_{Y(\alpha)}(1)$ is the restriction of $\mathcal{O}_{\mathbb{P}(A)}(1)$.

Theorem 4.7 ([K3, K9]). *The map $X(\alpha) \mapsto (Y(\alpha), E(\alpha))$ gives an isomorphism of the moduli space \mathcal{MF}_{14}^1 of Fano threefolds X_{14} of genus 8 and the moduli space $\mathcal{MFL}_3^2(2)$ of pairs (Y, E) , where Y is a smooth cubic threefold and E is a stable vector bundle on Y of rank 2, $c_1(E) = 0$, $c_2(E) = 2L$ with $H^1(Y, E(-1)) = 0$. For every α there is an equivalence of categories $\mathcal{B}_{X_{14}(\alpha)} \cong \mathcal{A}_{Y_3(\alpha)}$.*

Remark 4.8. The bundles E in the statement of the Theorem are known as instanton bundles of charge 2.

Let $Z_3 \subset \mathcal{MF}_3^2 \times \mathcal{MF}_{14}^1$ be the graph of the morphism $\mathcal{MF}_{14}^1 \cong \mathcal{MFT}_3^2(2) \rightarrow \mathcal{MF}_3^2$.

Corollary 4.9. *For any pair of threefolds $(Y_3, X_{14}) \in Z_3 \subset \mathcal{MF}_3^2 \times \mathcal{MF}_{14}^1$ we have an equivalence of categories $\mathcal{A}_{X_{14}} \cong \mathcal{B}_{Y_3}$.*

4.4. Geometrical correspondences. Actually the proof of Theorem 4.7 in [K3] gives more than just an equivalence of categories. It gives also a geometrical correspondence between $X(\alpha)$ and $Y(\alpha)$.

Let $\mathbb{P}_{X(\alpha)}(\mathcal{E})$ be the projectivization of the exceptional rank 2 bundle on $X(\alpha)$. Since \mathcal{E} is the restriction of the tautological bundle from the Grassmannian $\mathrm{Gr}(2, W)$, we have a canonical map $\mathbb{P}_{X(\alpha)}(\mathcal{E}) \rightarrow \mathbb{P}(W)$. On the other hand, one can check that we have an isomorphism $H^0(Y(\alpha), E(\alpha)^* \otimes \mathcal{O}_{Y(\alpha)}(1)) \cong W^*$, hence we have also a canonical map $\mathbb{P}_{Y(\alpha)}(E(\alpha)) \rightarrow \mathbb{P}(W)$.

Theorem 4.10 ([K3]). *The images of $\mathbb{P}_{X(\alpha)}(\mathcal{E})$ and $\mathbb{P}_{Y(\alpha)}(E(\alpha))$ in $\mathbb{P}(W)$ coincide with a quartic hypersurface $M \subset \mathbb{P}(W)$ singular along a curve $C \subset M$ of genus 26. The maps $\mathbb{P}_{X(\alpha)}(\mathcal{E}) \rightarrow M$ and $\mathbb{P}_{Y(\alpha)}(E(\alpha)) \rightarrow M$ are small contractions and induce isomorphisms over the complement of C . Moreover, the induced birational isomorphism $\mathbb{P}_{X(\alpha)}(\mathcal{E}) \dashrightarrow \mathbb{P}_{Y(\alpha)}(E(\alpha))$ is a flop.*

Remark 4.11. The hypersurface $M \subset \mathbb{P}(W)$ is known as the *da Palatini quartic*. The curve C parameterizes lines on $X(\alpha)$ and at the same time jumping lines for $E(\alpha)$ on $Y(\alpha)$ (that is lines $L \subset Y(\alpha)$ for which $E(\alpha)|_L \cong \mathcal{O}_L(1) \oplus \mathcal{O}_L(-1)$).

We expect that some generalization of this result should hold for other values of d . For example, let $\mathcal{MFB}_{4d+2}^1(t)$ be the moduli space of pairs (X, F) , where X is a Fano threefold of index 1 and degree $4d + 2$, and F is a stable vector bundle on X of rank 2 with $c_1(F) = -H$, $c_2(F) = (d + 2 + t)L$ (note that for $t = 0$ by Theorem 2.5 there is only one such bundle, the exceptional bundle \mathcal{E}_2 , hence $\mathcal{MFB}_{4d+2}^1(0) = \mathcal{MF}_{4d+2}^1$). Using the Riemann–Roch (see the proof of Proposition 3.9) one can check that for any d , any t and any $(X, F) \in \mathcal{MFB}_{4d+2}^1(t)$ we have

$$\dim H^0(X, F^*) = d + 3 - t,$$

so that we have a map $\mathbb{P}_X(F) \rightarrow \mathbb{P}^{d+2-t}$. Moreover, since $c_1(F^*) = H$, $c_2(F^*) = (d + 2 - t)L$, it follows that the degree of the image of $\mathbb{P}_X(F)$ in \mathbb{P}^{d+2-t} is

$$\deg \mathbb{P}_X(F) = c_1(F^*)^3 - 2c_1(F^*)c_2(F^*) = H^3 - 2(d + 2 - t)HL = (4d + 2) - 2(d + 2 - t) = 2d - 2 + 2t.$$

Similarly, let $\mathcal{MFL}_d^1(k)$ be the moduli space of pairs (Y, E) , where Y is a Fano threefold of index 2 and degree d , and E is an instanton bundle of charge k on Y , that is a stable vector bundle of rank 2 with $c_1(E) = 0$, $c_2(E) = kL$ and $H^1(Y, E(-1)) = 0$ (see [K3]). Using the Riemann–Roch (see the proof of Proposition 3.9) one can check that for any d , any k and any $(Y, E) \in \mathcal{MFL}_d^1(k)$ we have

$$\dim H^0(Y, E^*(1)) = 2d - 2k + 4,$$

so that we have a map $\mathbb{P}_Y(E) \rightarrow \mathbb{P}^{2d-2k+3}$. Moreover, since $c_1(E^*(1)) = 2H$, $c_2(E(1)) = H^2 + kL$, it follows that the degree of the image of $\mathbb{P}_Y(E)$ in $\mathbb{P}^{2d-2k+3}$ is

$$\deg \mathbb{P}_Y(E) = c_1(E^*(1))^3 - 2c_1(E^*(1))c_2(E^*(1)) = 8H^3 - 4H(H^2 + kL) = 8d - 4(d + k) = 4d - 4k.$$

Note that whenever $d + 1 = 2k - t$ the dimensions and the degrees coincide.

Conjecture 4.12. *For each $1 \leq d \leq 5$ there are integers $k, t \geq 0$ satisfying $d + 1 = 2k - t$ for which there is an isomorphism $\xi : \mathcal{MFL}_d^2(k) \cong \mathcal{MFB}_{4d+2}^1(t)$ such that for $(X, F) = \xi(Y, E)$ there is an isomorphism $h : H^0(Y, E^*(1)) \xrightarrow{\sim} H^0(X, F^*)$ and a birational isomorphism $\theta : \mathbb{P}_X(F) \dashrightarrow \mathbb{P}_Y(E)$ such that the*

diagram

$$\begin{array}{ccc}
 \mathbb{P}_X(F) & \overset{\theta}{\dashrightarrow} & \mathbb{P}_Y(E) \\
 \downarrow & & \downarrow \\
 \mathbb{P}(H^0(X, F^*)^*) & \xrightarrow[\cong]{h} & \mathbb{P}(H^0(Y, E^*(1))^*)
 \end{array}$$

commutes. Moreover, there is an equivalence $\mathcal{A}_X \cong \mathcal{B}_Y$, that is $Z_d \subset \mathcal{MF}_d^2 \times \mathcal{MF}_{4d+2}^1$ is the image of the graph of the isomorphism $\xi : \mathcal{MF}_d^2(k) \rightarrow \mathcal{MF}_{4d+2}^1(t)$.

Remark 4.13. For $d = 3$ by Theorem 4.10 we should take $k = 2$, $t = 0$. For $d = 5$ we expect $k = 4$, $t = 2$ will work.

5. APPENDIX. THE GROTHEDIECK GROUP AND ALGEBRAIC CYCLES

Let X be a smooth projective variety of dimension n . Let $A^p(X)$ denote the group of algebraic cycles on X of codimension p modulo rational equivalence. Let $A^\bullet(X) = \bigoplus_{p=0}^n A^p(X)$ be the Chow ring. Let $K_0(X)$ be the Grothendieck group of the category of coherent sheaves on X (equivalently, of the derived category $\mathcal{D}^b(X)$). Consider the Chern character map $\text{ch} : K_0(X) \rightarrow A^\bullet(X) \otimes \mathbb{Q}$. It is well known that ch induces an isomorphism of \mathbb{Q} -vector spaces $K_0(X) \otimes \mathbb{Q} \rightarrow A^\bullet(X) \otimes \mathbb{Q}$.

On the other hand, consider on both sides the numerical equivalence. Recall that an algebraic cycle $a \in A^p(X)$ is numerically equivalent to zero, if it lies in the kernel of the bilinear intersection form:

$$A^\bullet(X) \otimes A^\bullet(X) \longrightarrow A^\bullet(X) \xrightarrow{\text{pr}} A^n(X) \xrightarrow{\text{deg}} \mathbb{Z}.$$

In other words, if its intersection with any cycle in $A^{n-p}(X)$ is zero. Let $A^\bullet(X)_{\text{num}} = \bigoplus A^p(X)_{\text{num}}$ be the ring of algebraic cycles modulo the numerical equivalence. Note that any torsion class in $A^\bullet(X)$ is numerically trivial, hence $A^\bullet(X)_{\text{num}}$ is torsion free.

Similarly, a class $v \in K_0(X)$ is numerically equivalent to zero, if it lies in the kernel of the Euler bilinear form:

$$\chi : K_0(X) \otimes K_0(X) \longrightarrow \mathbb{Z}, \quad \chi([F], [G]) = \sum_i (-1)^i \dim \text{Ext}^i(F, G).$$

Let $K_0(X)_{\text{num}} := K_0(X) / \text{Ker } \chi$ be the numerical Grothendieck group.

The Riemann-Roch formula shows that the kernel of the Euler form coincides with the preimage under the Chern character map of the subring of $A^\bullet(X) \otimes \mathbb{Q}$ consisting of numerically trivial algebraic cycles. It follows that ch descends to a map $K_0(X)_{\text{num}} \rightarrow A^\bullet(X)_{\text{num}} \otimes \mathbb{Q}$ which we denote by ch as well, and induces an isomorphism of \mathbb{Q} -vector spaces $K_0(X)_{\text{num}} \otimes \mathbb{Q} \cong A^\bullet(X)_{\text{num}} \otimes \mathbb{Q}$.

For any p -cycle $Z = \sum a_i S_i$ we define $[\mathcal{O}_Z] := \sum a_i [\mathcal{O}_{S_i}] \in K_0(X)$. Note that $[\mathcal{O}_Z]$ really depends on the cycle Z , not only on its rational or numerical equivalence class. A little bit later we will show how one can get rid of this dependance (see Remark 5.3).

Definition 5.1. We will say that a smooth projective n -dimensional variety X is **AK-compatible** if for any collection of cycles Z_p^i on X , $0 \leq p \leq n$, $1 \leq i \leq m_p$ such that $\text{codim } Z_p^i = p$ and $\{Z_1^p, Z_2^p, \dots, Z_{m_p}^p\}$ is a basis in $A^p(X)_{\text{num}}$ the classes $[\mathcal{O}_{Z_p^i}]$, $0 \leq p \leq n$, $1 \leq i \leq m_p$ form a \mathbb{Z} -basis in $K_0(X)_{\text{num}}$.

If an algebraic variety X is AK-compatible, then one can easily describe its numerical Grothendieck group by choosing some bases in the groups of algebraic cycles and considering their structure sheaves. Certainly such a description may be useful in many cases. The goal of this section is to find some easily verifiable criterion for AK-compatibility.

We start with some preparations. Consider the following two filtrations on $K_0(X)_{num}$. The first one is induced by the codimension filtration on $A^\bullet(X)$:

$$F^p K_0(X)_{num} = \text{ch}^{-1}(\oplus_{q=p}^n A^q(X)_{num} \otimes \mathbb{Q}).$$

The second one is induced by the codimension of support:

$$S^p K_0(X)_{num} = \langle [G] \mid \text{codim supp}(G) \geq p \rangle,$$

where $\langle \ \rangle$ stands for the linear span. We will call the filtration F^\bullet the induced filtration, and S^\bullet the support filtration. Let $\text{gr}_F^p K_0(X)_{num}$ and $\text{gr}_S^p K_0(X)_{num}$ be the graded factors of these filtrations.

Note that for any $G \in \mathcal{D}^b(X)$ with $\text{codim supp}(G) \geq p$ we have $\text{ch}(G) \in \oplus_{q \geq p} A^q(X)_{num} \otimes \mathbb{Q}$, hence $S^p K_0(X)_{num} \subset F^p K_0(X)_{num}$ and we have the following commutative diagram

$$\begin{array}{ccccc} S^p K_0(X)_{num} & \hookrightarrow & F^p K_0(X)_{num} & \xrightarrow{\text{ch}} & \oplus_{q \geq p} A^q(X)_{num} \otimes \mathbb{Q} \\ \uparrow & & \uparrow & & \uparrow \\ S^{p+1} K_0(X)_{num} & \hookrightarrow & F^{p+1} K_0(X)_{num} & \xrightarrow{\text{ch}} & \oplus_{q \geq p+1} A^q(X)_{num} \otimes \mathbb{Q} \end{array}$$

Passing to the graded factors we obtain a chain of maps

$$\text{gr}_S^p K_0(X)_{num} \xrightarrow{i_p} \text{gr}_F^p K_0(X)_{num} \xrightarrow{\text{ch}_p} A^p(X)_{num} \otimes \mathbb{Q}$$

Here ch_p is the p -th coefficient of the Chern character and i_p is the map induced by the identity map of $K_0(X)_{num}$. Note that it follows that $\text{ch}_p : \text{gr}_F^p K_0(X)_{num} \otimes \mathbb{Q} \rightarrow A^p(X)_{num} \otimes \mathbb{Q}$ is an isomorphism. Let us show that the inverse map is defined over \mathbb{Z} .

Lemma 5.2. *There exists a linear map $\mathcal{O}_p : A^p(X)_{num} \rightarrow \text{gr}_F^p K_0(X)_{num}$, which is inverse to ch_p . Moreover, $\mathcal{O}_p(Z) = [\mathcal{O}_Z] \bmod F^{p+1} K_0(X)_{num}$.*

Proof: For each p -cycle Z on X define $\mathcal{O}_p(Z)$ as the image of the class of its structure sheaf in $\text{gr}_F^p K_0(X)_{num}$. Since

$$\text{ch}(\mathcal{O}_Z) = Z + \text{terms of degree larger than } p, \quad (\dagger)$$

we have $[\mathcal{O}_Z] \in F^p K_0(X)_{num}$ and $\text{ch}_p(\mathcal{O}_p(Z)) = Z$. Since ch_p is injective, it follows that \mathcal{O}_p is correctly defined and $\text{ch}_p \circ \mathcal{O}_p = \text{id}$. \square

Remark 5.3. As we see from this Lemma the class of $[\mathcal{O}_Z]$ in $\text{gr}_F^p K_0(X)_{num}$ only depends on the numerical class of Z .

Further, it is easy to see that for any coherent sheaf G supported in codimension p the p -th coefficient of the Chern character is integer, $\text{ch}_p(G) \in A^p(X)_{num}$. Indeed, if Z_1, \dots, Z_m are codimension p components of $\text{supp } G$ and ℓ_i is the length of G at generic point of Z_i then $\text{ch}_p(G) = \sum \ell_i Z_i$. Moreover, the same argument shows that the map $\text{ch}_p : \text{gr}_S^p K_0(X)_{num} \rightarrow A^p(X)_{num}$ is surjective. It follows that we have the following commutative diagram

$$\begin{array}{ccc} A^p(X)_{num} & \hookrightarrow & A^p(X)_{num} \otimes \mathbb{Q} \\ \uparrow \text{ch}_p & \searrow \mathcal{O}_p & \uparrow \text{ch}_p \\ \text{gr}_S^p K_0(X)_{num} & \xrightarrow{i_p} & \text{gr}_F^p K_0(X)_{num} \end{array}$$

Proposition 5.4. *The following properties for a smooth projective variety X are equivalent:*

- (i) $\forall 0 \leq p \leq n$ \mathcal{O}_p is an isomorphism;
- (ii) X is AK-compatible;

- (iii) $\forall 0 \leq p \leq n$ $S^p K_0(X)_{num} = F^p K_0(X)_{num}$;
- (iv) $\forall 0 \leq p \leq n$ $\text{ch}_p(\text{gr}_F^p K_0(X)_{num}) \subset A^p(X)_{num}$.

Proof: (i) \Rightarrow (ii): An evident induction argument shows that $\{[\mathcal{O}_{Z_q^i}]\}_{q \geq p}^{1 \leq i \leq m_q}$ is a basis in $F^p K_0(X)_{num}$.

(ii) \Rightarrow (iii): Assume that X is AK-compatible. Choose bases in all $A^p(X)_{num}$ as in Definition 5.1 and assume that $v = \sum a_q^i [\mathcal{O}_{Z_q^i}] \in F^p K_0(X)_{num}$ for some $a_q^i \in \mathbb{Z}$. Let q be the minimal integer such that $a_q^i \neq 0$ for some i and assume that $q < p$. Then applying ch_q we see that

$$0 = \text{ch}_q(v) = \sum_i a_q^i \text{ch}_q([\mathcal{O}_{Z_q^i}]) = \sum_i a_q^i \text{ch}_q(\mathcal{O}_q(Z_q^i)) = \sum_i a_q^i Z_q^i$$

which implies $a_q^i = 0$ for all i . So, it follows that $q \geq p$, hence $a_q^i = 0$ for all $q < p$. But then it is clear that $v \in S^p K_0(X)_{num}$. So we see that $F^p K_0(X)_{num} \subset S^p K_0(X)_{num}$.

(iii) \Rightarrow (iv): If $S^p K_0(X)_{num} = F^p K_0(X)_{num}$ then $\text{gr}_S^p K_0(X)_{num} = \text{gr}_F^p K_0(X)_{num}$ for all p , hence $\text{ch}_p(\text{gr}_F^p K_0(X)_{num}) = \text{ch}_p(\text{gr}_S^p K_0(X)_{num}) \subset A^p(X)_{num}$.

(iv) \Rightarrow (i): Since ch_p and \mathcal{O}_p are mutually inverse isomorphisms of $\text{gr}_F^p K_0(X)_{num} \otimes \mathbb{Q}$ and $A^p(X)_{num} \otimes \mathbb{Q}$ and preserve lattices $\text{gr}_F^p K_0(X)_{num}$ and $A^p(X)_{num}$, it follows that they induce isomorphisms of these lattices. \square

Our next goal is to give several sufficient conditions for AK-compatibility.

Lemma 5.5. *For any smooth projective variety X we have $\text{ch}_p(\text{gr}_F^p K_0(X)_{num}) \subset A^p(X)_{num}$ for $p = 0, 1, 2$ and $p = n$.*

Proof: Note that $\text{ch}_0(G)$ is the rank of G and $\text{ch}_1(G) = c_1(G)$, which implies the claim for $p = 0$ and $p = 1$. For $p = 2$ we have $\text{ch}_2(G) = c_1(G)^2/2 - c_2(G)$, so if $c_1(G) = \text{ch}_1(G) = 0$, then $\text{ch}_2(G) = -c_2(G) \in A^2(X)_{num}$. Finally, if $G \in F^n K_0(X)_{num}$ then by Riemann–Roch we have $\text{ch}_n(X) = \chi(\mathcal{O}_X, G) \in \mathbb{Z}$, where we have identified $A^n(X)_{num}$ with \mathbb{Z} via the degree map. This proves the claim for $p = n$. \square

Corollary 5.6. *If $\dim X \leq 3$ then X is AK-compatible.*

Another approach to AK-compatibility is given by the following

Lemma 5.7. *Assume that the intersection pairing $A^p(X)_{num} \otimes A^{n-p}(X)_{num} \rightarrow \mathbb{Z}$ induces an isomorphism $A^p(X)_{num} \rightarrow A^{n-p}(X)_{num}^*$. Then the map $\mathcal{O}_p : A^p(X)_{num} \rightarrow \text{gr}_F^p K_0(X)_{num}$ is an isomorphism. In particular, if the intersection pairing $A^p(X)_{num} \otimes A^{n-p}(X)_{num} \rightarrow \mathbb{Z}$ induces an isomorphism $A^p(X)_{num} \rightarrow A^{n-p}(X)_{num}^*$ for all p then X is AK-compatible.*

Proof: Let $Z, W \subset X$ be subschemes of codimension p and $(n-p)$ respectively. Then (\dagger) and Riemann–Roch imply that

$$Z \cdot W = (-1)^p \chi(\mathcal{O}_Z, \mathcal{O}_W),$$

hence we have a commutative diagram

$$\begin{array}{ccc} A^p(X)_{num} & \xrightarrow{\mathcal{O}_p} & \text{gr}_F^p K_0(X)_{num} \\ \downarrow \cdot & & \downarrow (-1)^p \chi \\ A^{n-p}(X)_{num}^* & \xleftarrow{\mathcal{O}_p^*} & \text{gr}_F^{n-p} K_0(X)_{num}^* \end{array}$$

Note that all the maps are finite index embeddings. So, if the left vertical arrow is an isomorphism then \mathcal{O}_p is also an isomorphism. \square

The results of this section allow to describe $K_0(X)_{num}$ for all Fano threefolds.

Corollary 5.8. *Let V be a Fano threefold with $\text{Pic } V \cong \mathbb{Z}$. Let H be the generator of $\text{Pic } V$, L a line on V , and P a point on V . Then $K_0(X)_{\text{num}} = \langle [\mathcal{O}_V], [\mathcal{O}_H], [\mathcal{O}_L], [\mathcal{O}_P] \rangle$.*

Proof: We can argue either by Corollary 5.6 or by Lemma 5.7 that V is AK-compatible. Hence by definition of AK-compatibility we obtain the required basis. \square

Remark 5.9. One can combine the results of Lemma 5.5 and Lemma 5.7 for the verification of AK-compatibility. In other words, if for an algebraic variety X the conditions of Lemma 5.7 are true for all $3 \leq p \leq n - 1$ then X is AK-compatible. Indeed, it is easy to see from the proof of Proposition 5.4 that the property (ivp) for each p implies the property (ip).

These considerations apply e.g. for cubic fourfolds. Indeed, for $p = 3$ the conditions of Lemma 5.7 are true, hence the cubic fourfold is AK-compatible.

REFERENCES

- [B] A. Bondal, *Representations of associative algebras and coherent sheaves*, (Russian) *Izv. Akad. Nauk SSSR Ser. Mat.* **53** (1989), no. 1, 25–44; translation in *Math. USSR-Izv.* **34** (1990), no. 1, 23–42.
- [BK] A. Bondal, M. Kapranov, *Representable functors, Serre functors, and reconstructions*, (Russian) *Izv. Akad. Nauk SSSR Ser. Mat.* **53** (1989), no. 6, 1183–1205, 1337; translation in *Math. USSR-Izv.* **35** (1990), no. 3, 519–541.
- [BO1] A. Bondal, D. Orlov, *Semiorthogonal decomposition for algebraic varieties*, preprint math.AG/9506012.
- [Fu] Fujita T., *On the structure of polarized varieties with A -genera zero*, *J. Fac. Sci. Univ Tokyo. Sec. IA.* **22**, 103–115.
- [Is] V. Iskovskikh, *Anticanonical models of three-dimensional algebraic varieties*, *Current problems in mathematics*, VINITI, Moscow, **12**, 59–157 (Russian); translation in *J. Soviet Math.* **13** (1980) 745–814.
- [IP] V. Iskovskikh, Yu. Prokhorov, *Fano varieties*, *Algebraic geometry*, V, 1–247, *Encyclopaedia Math. Sci.*, **47**, Springer, Berlin, 1999.
- [K1] Kuznetsov A., *An exceptional collection of vector bundles on V_{22} Fano threefolds*, *Vestnik MGU, Ser. 1, Mat. Mekh.*, 1996, no. 3, p. 41–44 (in Russian).
- [K2] Kuznetsov A., *Fano threefolds V_{22}* , preprint MPIM/97-24.
- [K3] Kuznetsov A., *Derived categories of cubic and V_{14} threefolds*, *Proc. V.A.Steklov Inst. Math.*, V. 246 (2004), p. 183–207; preprint math.AG/0303037.
- [K5] Kuznetsov A., *Hyperplane sections and derived categories*, *Izvestiya RAN: Ser. Mat.* **70:3** p. 23–128 (in Russian); translation in *Izvestiya: Mathematics* **70:3** p. 447–547.
- [K6] Kuznetsov A., *Derived categories of quadric fibrations and intersections of quadrics*, *Advances in Mathematics*, V. 218 (2008), N. 5, 1340–1369.
- [K8] Kuznetsov A., *Lefschetz decompositions and categorical resolutions of singularities*, *Selecta Mathematica*, V. 13 (2008), N. 4, 661–696.
- [K9] Kuznetsov A., *Homological projective duality for Grassmannians of lines*, preprint math.AG/0610957.
- [M] Mukai S., *Fano 3-folds*, *London Math. Soc. Lect. Note Ser.* **179**, 255–263.
- [MU] S. Mukai, H. Umemura, *Minimal rational threefolds*, *Algebraic geometry (Tokyo/Kyoto, 1982)*, 490–518, *Lecture Notes in Math.*, **1016**, Springer, Berlin, 1983.
- [Or] D. Orlov, *Exceptional set of vector bundles on the variety V_5* , (Russian) *Vestnik Moskov. Univ. Ser. I Mat. Mekh.* 1991, no. 5, 69–71; translation in *Moscow Univ. Math. Bull.* **46** (1991), no. 5, 48–50.
- [O1] D. Orlov, *Projective bundles, monoidal transformations, and derived categories of coherent sheaves*, (Russian) *Izv. Ross. Akad. Nauk Ser. Mat.* **56** (1992), no. 4, 852–862; translation in *Russian Acad. Sci. Izv. Math.* **41** (1993), no. 1, 133–141.

ALGEBRA SECTION, STEKLOV MATHEMATICAL INSTITUTE, 8 GUBKIN STR., MOSCOW 119991 RUSSIA

THE PONCELET LABORATORY, INDEPENDENT UNIVERSITY OF MOSCOW

E-mail address: akuznet@mi.ras.ru